

**NATURAL RESOURCES CONSERVATION SERVICE
NEW JERSEY
CONSERVATION PRACTICE STANDARD**

FIELD BORDER

(Ac.)

CODE 386

DEFINITION

A strip of permanent vegetation established at the edge or around the perimeter of a field.

PURPOSE

This practice may be applied to accomplish one or more of the following:

- Reduce erosion from wind and water
- Protect soil and water quality
- Manage pest populations
- Provide wildlife food and cover
- Increase carbon storage
- Improve air quality

CONDITIONS WHERE PRACTICE APPLIES

This practice is applied around the perimeter of fields. Its use can support or connect other buffer practices within and between fields. This practice may also apply to recreation land or other land uses where agronomic crops including forages are grown.

CRITERIA

General Criteria Applicable to All Purposes

Field borders shall be established around the field edges to the extent needed to meet the resource needs and producer objectives. Minimum field border widths shall be at least 15 feet.

The field borders shall be established to adapted species of permanent grass, legumes

and/or shrubs that accomplish the design objective and do not function as host for diseases of the field crop.

Plants selected for field borders will have the physical characteristics necessary to control wind and water erosion to tolerable levels on the field border area. Species shall be selected from Standards 342 Critical Area Planting and 512 Pasture and Hayland Planting. Native species that can fulfill the purposes of the Standard are preferred.

Seedbed preparation, seeding rates, dates, depths, fertility requirements, and planting methods will be consistent with approved local criteria and site conditions.

Ephemeral gullies and rills present in the planned border area will be eliminated as part of seedbed preparation. If present, ephemeral gullies and rills located immediately upslope from the planned border area need to be treated to ensure more of a sheet flow into the planned border area.

Additional Criteria to Reduce Erosion from Wind and Water

Field border establishment, in conjunction with other practices, will be timed so that the soil will be adequately protected during the critical erosion period(s).

Establish stiff-stemmed, upright grasses, grass/legumes or forbs to trap wind- or water-borne soil particles.

The amount of surface and/or canopy cover needed from the field border shall be determined using current approved water and wind erosion prediction technology.

Calculations shall account for the effects of other practices in the management system.

Wind Erosion Reduction. Locate borders to provide a stable area on the windward edge of the field as determined by prevailing wind direction data.

Minimum height of grass or forbs shall be one foot during the critical erosion period.

Water Erosion Reduction. Locate borders to eliminate sloping end rows, headlands, and other areas where concentrated water flows will enter or exit the field.

Orient plant rows as closely as possible to perpendicular to sheet flow direction.

Additional Criteria to Protect Soil and Water Quality

Do not burn the field border if the main goal of the field border is to protect soil or water quality.

Reducing Runoff and Increasing Infiltration.

Locate borders around the perimeter of the field, or as a minimum, install borders to eliminate sloping end rows, headlands and other areas where concentrated water flows will enter or exit the field.

Water Quality – Adsorbed, Dissolved and Suspended Contaminants. As a minimum, locate field borders along the edge(s) of the field where runoff enters or leaves the field. The minimum width for this purpose shall be 30 feet and have a vegetation stem density/retardance of moderate to high (e.g. equivalent to a good stand of wheat).

Design border widths to comply with all applicable State and local regulations regarding manure and chemical application setbacks.

Reducing Soil Compaction from Equipment Parking and Traffic. Border widths will be designed to accommodate equipment turning, parking, loading/unloading equipment, grain harvest operations, etc.

Additional Criteria to Manage Pest Populations

Provide a Harbor for Beneficial organisms (e.g. insects, mites, etc.).

Include the following herbaceous plants that attract beneficial insects:

Assassin bugs – alfalfa, carrot, goldenrod

Big-eyed bugs- alfalfa, carrot, goldenrod

Damsel bugs – alfalfa

Lacewings – carrot

Lady bugs – alfalfa, morning glory, yarrow, goldenrod

Parasitic wasps – yarrow, white clover, goldenrod, carrot, alfalfa

Syrphid flies – buckwheat, candytuft, fennel, ivy, white sweet clover

Tachinid flies – coriander, fennel, ivy, white sweet clover

Yellow jacket wasps – ivy, white sweet clover, carrots

Mowing, harvesting, and pesticide applications will be scheduled to accommodate life cycle requirements of the beneficial insects. Other information can be found online at <http://www.attra.org/attra-pub/farmscape.html>

or

Provide a Habitat to Cause Pests to Congregate. Select plants for the field border that attract pests (e.g. alfalfa strips planted to lure lygus bugs away from a cotton crop).

These include:

Nasturtium – aphids

Radish, nasturtium – flea beetles

Egg plant – potato bugs

Use mechanical, cultural, and/or chemical techniques to reduce pest populations when and where they congregate in the field border.

Mowing, harvesting, pesticide applications and other disturbance activities will be scheduled to accommodate life cycle requirements of the beneficial organisms.

Additional Criteria To Provide Wildlife Food And Cover

Methods that provide wildlife food and cover shall be employed as per the Standard 645 Upland Wildlife Habitat Management.

Mowing, harvest, and weed control activities within the field border will be scheduled after July 15th and prior to October 1 to accommodate reproduction and other requirements of grassland nesting birds. Width of borders for wildlife should be a minimum of 25 feet wide and preferably 50 feet or more. Care should be taken to avoid vegetation over 3 feet tall that will provide escape cover for deer.

Vegetative successional state shall be maintained to accommodate target wildlife species requirements.

Additional Criteria to Increase Carbon Storage

Establish plant species that will produce adequate above- and below-ground biomass for the site (i.e. a positive soil conditioning index).

Maximize the width and length of the herbaceous border to fit the site and increase total biomass production.

Do not burn if the main goal of the field border is carbon storage.

Do not disturb the roots of the established vegetation with tillage.

Additional Criteria to Improve Air Quality

Establish plant species with morphological characteristics that optimize interception and adhesion of airborne particulates. Select plants with persistent roots and residue that stabilize soil aggregates and capture airborne soil particles.

Establish species resistant to damage from equipment traffic.

CONSIDERATIONS

Consider planting field borders around the entire field, not just on the field edges where water enters or leaves the field, for maximizing multiple resource protection.

Establishing a narrow strip of stiff-stemmed upright grass at the crop/field border interface can increase soil particle trapping efficiency of the field border.

Native plants are best suited for wildlife habitat enhancement and provide other ecological benefits where adapted to site conditions and when consistent with producer objectives.

Include native plants that provide diverse pollen and nectar sources to encourage local pollinator populations.

Use field borders as corridors to connect existing or planned habitat blocks.

Prescribed burning, strip disking, or selective herbicide applications are management tools that can be used to maintain suitable habitat for specifically desired wildlife species.

Overseed the field border with legumes for increased plant diversity, soil quality, and wildlife benefits.

Waterbars or berms may be needed to breakup or redirect concentrated water flow within the borders.

In selecting plant species to establish in the field border, among other items, consider the plant's tolerance to:

- Sediment deposition and chemicals planned for application
- Drought in arid areas or where evapotranspiration can potentially exceed precipitation during the field border's active growing period(s).
- Equipment traffic.

Design border widths to match the required field application setback widths for easier management (i.e. land-use and management changes occur in the same location).

Establish plant species that will have the desired visual effects and that will not interfere with field operations or field border maintenance.

Consider the amount of shading that the field border or portions of the field border may experience and select species for those locations accordingly.

The use of native perennial plant species as opposed to annual species provides a longer period of resource protection.

Consider installing other conservation practices on adjacent upland areas to reduce

surface runoff and excessive sedimentation of field borders.

PLANS AND SPECIFICATIONS

Prepare plans and specifications for each field or treatment unit according to the Criteria included in this Standard. Specifications shall describe the requirements for applying this practice to meet the intended purpose. Record practice specifications on the Field Border 386, Conservation Practice Job Sheet. The following components shall be included for recording this specification:

- Field Border widths and lengths based on local design criteria.
- Field Border location(s) within the field(s) or farm boundary.
- Species to be used and the location and planting density of the species used.
- Site preparation requirements.
- Timing of planting and planting method.
- Liming or fertilizer requirements.
- Operation and maintenance requirements.

OPERATION AND MAINTENANCE

Field borders require careful management and maintenance for performance and longevity. The following O&M activities will be planned and applied as needed:

- Repair storm damage.
- Remove sediment from above or within the field border when accumulated sediment either alters the function of the field border or threatens the degradation of the planted species' survival.
- Shut off sprayers and raise tillage equipment to avoid damage to field borders.

- Shape and reseed border areas damaged by animals, chemicals, tillage, or equipment traffic.
- Maintain desired vegetative communities and plant vigor by liming, fertilizing, mowing, disking, or burning and controlling noxious weeds to sustain effectiveness of the border.
- Repair and reseed ephemeral gullies and rills that develop in the border.
- Minimally invasive tillage (e.g. paraplowing) may be performed in rare cases where compaction and vehicle traffic have degraded the field border function. The purpose of the tillage is strictly to decrease bulk density and increase infiltration rates so as to provide a better media for reestablishment of vegetation and field border function.
- Maintenance activities that result in disturbance of vegetation should not be conducted during the nesting season of grass nesting birds.
- Avoid vehicle traffic when soil moisture conditions are saturated.

REFERENCES

K. G. Renard, G. R. Foster, G. A. Weesies, K. D. K. McCool and D. C. Yoder. 1997. Predicting Soil Erosion by Water: A Guide to Conservation Planning with the Revised Universal Soil Loss Equation (RUSLE), Agricultural Handbook Number 703.

Revised Universal Soil Loss Equation Version 2 (RUSLE2) website (checked May 2007): http://fargo.nserl.purdue.edu/rusle2_dataweb/RUSLE2_Index.htm.