

Map Symbol	Map Unit Name	Nontechnical Descriptions
BA	BARBARY ASSOCIATION	This soil is level and very poorly drained. It is a very fluid mineral soil in swamps. This soil is ponded and flooded most of the time. Typically, the soil has a muck surface layer and a gray, very fluid clay underlying material. This soil has low strength. The total subsidence potential is medium. If the soil is drained, it can have a very high shrink-swell potential.
Cc	COMMERCE SILT LOAM	This nearly level, somewhat poorly drained soil is on alluvial plains. It is loamy throughout and has high fertility. Runoff is slow, and water and air move moderately slowly through the soil. A seasonal high water table is about 1.5 to 4 feet below the surface during December through April. The shrink-swell potential is moderate. Slopes range from 0 to 2 percent.
Cm	COMMERCE SILTY CLAY LOAM	This nearly level, somewhat poorly drained soil is on alluvial plains. It is loamy throughout and has high fertility. Runoff is slow, and water and air move moderately slowly through the soil. A seasonal high water table is about 1.5 to 4 feet below the surface during December through April. The shrink-swell potential is moderate. Slopes range from 0 to 2 percent.
FA	FAUSSE ASSOCIATION	These level, very poorly drained soils are in low, depressional areas on the alluvial plain. They formed in alluvium and are clayey throughout their profiles. These soils are ponded or flooded most of the time. Water and air move very slowly through the soils. The soils have high fertility. The shrink-swell potential is very high, but the soils seldom dry enough to shrink and crack. Slopes are less than 1 percent.
Sa	SHARKEY SILTY CLAY LOAM	This level or nearly level, poorly drained soil is on flood plains. The surface layer is loamy and the subsoil is clayey. Cracks form during dry periods, and they seal over during wet periods. Natural fertility is high. Runoff is slow. A seasonal high water table is within 2 feet of the soil surface during December to April. Flooding is rare. The soil dries slowly once wetted. The shrink-swell potential is high or very high in the subsoil. Slopes are less than 1 percent.
Sh	SHARKEY CLAY	This nearly level, poorly drained, soil is on broad flats on the alluvial plain. It is clayey throughout. Natural fertility is medium or high. Runoff is slow or very slow. Water and air move very slowly through the soil. The shrink-swell potential is high or very high. A seasonal high water table is within 2 feet of the soil surface during December through April. Flooding is rare, but it can occur during unusually wet periods. Slopes are less than 1 percent.

Map Symbol	Map Unit Name	Nontechnical Descriptions
Sk	SHARKEY CLAY, FREQUENTLY FLOODED	This level, poorly drained or somewhat poorly drained soil is at low elevations on the alluvial plain. It is flooded frequently for very long periods. This soil is clayey throughout or it has a loamy surface layer and a clayey subsoil. Natural fertility is high. Surface runoff is very slow. Water and air move very slowly through the soil. The seasonal high water table is near the soil surface. This soil has a very high shrink-swell potential. Slopes are less than 1 percent.
Tu	TUNICA CLAY	This level, poorly drained, clayey soil is on the flood plain of the Mississippi River. It has a clay surface layer and subsoil and a silty clay loam underlying material. The surface layer is very sticky when wet and has poor tilth. Cracks form in dry periods and seal over in wet periods. Natural fertility is high. This soil is wet for long periods in winter and spring. Flooding is rare, but it can occur during unusually wet periods. The shrink-swell potential is high in the subsoil.