

FORAGE SUITABILITY GROUP


Loam

FSG No.: G055BY100ND

Major Land Resource Area: 55B - Central Black Glaciated Plains

Physiographic Features

The soils in this group are located on upland positions of till plains and moraines. They are also found on glacial lake and delta plains and glacial stream terraces.



	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	980	1970
Slope (percent):	0	15
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):	0	0
Frequency:	None	None
Duration:	None	None
Runoff Class:	Low	High

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 55B. About 78% of the annual precipitation occurs during the months of April through September. On average there are about 28 days with greater than .1 inches of precipitation during the same time frame. Precipitation is lowest in the north west, and highest in the south in the MLRA. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 25 inches at Forman, ND to 37 inches at Columbia, SD. Snow cover at depths greater than 1 inch range from 32 days at Petersburg, ND to 98 days at Gackle, ND.

Average July temperatures are about 71 degrees F., and average January temperatures are about 7 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -39 at both Petersburg and Oakes in ND, and a high of 114 recorded at Mellette, SD. The MLRA lies in USDA Plant Hardiness Zones 3b and 4a.

At Aberdeen, SD the average annual wind speeds are about 11 MPH. The highest wind speeds occur during March though May, but average monthly wind speeds do not vary significantly throughout the year. It is cloudy about 163 days a year. Average morning relative humidity in June is about 85% and average afternoon humidity is 60%.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>

Freeze-free period (28 deg)(days): (9 years in 10 at least)	115	137
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 28	May 14
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 06	May 23
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Aug 29	Sep 10
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 08	Sep 21

Length of Growing Season (32 deg)(days): (9 years in 10 at least)	92	116
Growing Degree Days (40 deg):	3389	4402
Growing Degree Days (50 deg):	1852	2558
Annual Minimum Temperature:	-35	-25
Mean annual precipitation (inches):	16	21

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.24	0.13	0.30	0.63	1.08	1.72	1.30	0.94	0.76	0.23	0.18	0.24
Precip. More Than	0.60	0.79	2.10	3.58	4.09	5.07	3.66	4.02	3.07	1.92	1.14	0.74
Monthly Average:	0.50	0.43	1.02	1.89	2.41	3.39	2.65	2.27	1.94	1.18	0.57	0.46
Temp. Min.	-8.2	-2.7	11.6	28.1	39.9	50.0	54.0	51.2	40.8	30.3	15.0	-2.0
Temp. Max.	21.8	28.2	41.0	58.2	70.9	80.0	87.3	85.5	74.0	61.5	42.1	26.2
Temp. Avg.	7.4	13.6	26.9	42.8	55.7	65.4	71.0	68.7	57.6	45.8	28.3	12.9

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
ND5764	McVille, ND	1961	1990
ND7027	Petersburg, ND	1961	1990
ND2605	Ellendale, ND	1961	1987
ND3287	Fullerton, ND	1961	1990
ND3117	Forman, ND	1961	1990
ND2949	Fessenden, ND	1961	1990
ND4343	Hurdsfield, ND	1961	1990
SD0020	Aberdeen, SD	1961	1990
SD1873	Columbia, SD	1961	1990
ND2482	Edgeley, ND	1961	1990
ND4937	La Moure, ND	1961	1990
ND3309	Gackle, ND	1961	1990
ND4413	Jamestown, ND	1961	1990
ND8937	Valley City, ND	1961	1990
SD5456	Mellette, SD	1961	1990
ND2605	Oaks, ND	1961	1987

Soil Interpretations

This group consists of well and moderately well drained, moderately coarse to moderately fine textured soils formed mostly in glacial till or alluvium. Available water capacity is high and permeability is moderately slow to moderate.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Modertately slow	To	Moderate
Frost Action Class:	Moderate	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	2	8
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	5

Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.6	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	20

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov/>

Cool Season Grasses

Altai wildrye	F
Canada wildrye	F
Crested wheatgrass	G
Dahurian wildrye	G
Green needlegrass	G
Intermediate wheatgrass	G
Meadow brome	G
Newhy hybrid wheatgrass	G
Pubescent wheatgrass	G
Russian wildrye	G
Slender wheatgrass	G
Smooth brome	G
Tall wheatgrass	G
Western wheatgrass	G

Warm Season Grasses

Big bluestem	G
Blue grama	G
Indiangrass	G
Little bluestem	G
Prairie sandreed	F
Sand bluestem	F
Sideoats grama	G
Switchgrass	G

Legumes

Alfalfa	G
American vetch	G
Birdsfoot trefoil	F
Canada milkvetch	G
Cicer milkvetch	G
Hairy vetch	G
Purple prairieclover	G
Red clover	G
Sainfoin	F
Strawberry clover	F
Sweetclover	G
White clover	G
White prairieclover	G

G - Good adaptation for forage production on this group of soils in this MLRA
F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70% harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25% to 50%.

Forage Crop	Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	9100	3900
Alfalfa/Intermediate wheatgrass	7700	3500
Alfalfa/Smooth brome grass	7700	3500
Big bluestem	6600	2900
Crested wheatgrass	6000	2600
Green needlegrass	4600	2000
Intermediate wheatgrass	6300	2800
Smooth brome grass	6300	2800
Switchgrass	7700	3100
Western wheatgrass	4600	2000

Forage Growth Curves

Growth Curve Number: ND0001

Growth Curve Name: Alfalfa

Growth Curve Description: Alfalfa

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	30	20	15	5	0	0	0

Growth Curve Number: ND0002

Growth Curve Name: Cool season grass

Growth Curve Description: Cool season grass

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	35	10	5	5	0	0	0

Growth Curve Number: ND0003
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. On steeper slopes water erosion is a potential problem during establishment, when renovating stands, and in thin established stands. Livestock trail erosion is a potential problem in established stands.

Management Interpretations

Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

Pasture and hayland can include considerations for wildlife. Delaying grazing on portions of the pasture or rotating pastures will allow nest initiation of grassland nesting birds or species of concern. Nest initiation of most grassland nesting birds occurs from April 15 to June 1. Delaying haying until after July 15 allows for most species to fledge their young. Consider planting species with later maturity to allow for harvesting after nests have fledged. Avoid mowing around the field. Mow back and forth or from the inside to the outside of the field. Consider using flushing bars on swathers and mowers.

FSG Documentation

Similar FSG's:

FSG ID

G055BY109ND
G055BY120ND
G055BY500ND

FSG Narrative

Steep Loam soils are more steeply sloping with greater runoff potential.
Droughty Loam soils are shallower or coarser textured resulting in lower available water capacity and lower production potential.
Overflow soils receive additional moisture due to a favorable landscape position resulting in a higher production potential.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas, Natural Resources Conservation Service (NRCS) National Water and Climate Center, USDA Plant Hardiness Zone maps, National Soil Survey Information System (NASIS) for soil surveys in North Dakota and South Dakota counties in MLRA 55B, North Dakota and South Dakota NRCS Field Office Technical Guide, NRCS National Range and Pasture Handbook, various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:
ND
SD

Forage Suitability Group Approval:

Original Author: Tim Nordquist
Original Date: 9/18/00
Approval by: Jeff Printz
Approval Date: