

| |
|--|
| Table of Contents Section II – County Soil and Site Information Reports |
|--|

| | Issue Date | Date of Last Review | Responsible Staff |
|---|-----------------------|--------------------------------|------------------------------|
| <i>Use and Explanation of Soil Interpretations</i> | | | |
| <i>Explanation of Key Phrases Used in Soil Interpretations</i> | | | |
| Soils Legends | 4/02 | 4/02 | SOI |
| <i>*Acreage and Proportionate Extent of the Soils</i> | | | |
| Soil Descriptions - Nontechnical | 4/02 | 4/02 | SOI |
| <i>Use and Explanation of Nontechnical Descriptions</i> | | | |
| <i>*Nontechnical Soils Description Report</i> | | | |
| Soil Descriptions - Technical | 4/02 | 4/02 | SOI |
| <i>*Map Unit Description Report</i> | | | |
| Cropland Interpretations - Technical | 4/02 | 4/02 | SOI |
| <i>*Land Capability and Yields Per Acre of Crops</i> | | | |
| <i>*Prime Farmland Report</i> | | | |
| <i>*Soil Rating for Plant Growth Index</i> | | | |
| <i>*Soil Properties for Conservation Planning</i> | | | |
| Rangeland, Grazed Forestland, Native Pastureland Interpretations | 4/02 | 4/02 | SOI |
| <i>*Rangeland Productivity Report</i> | | | |
| <i>*Range Site Descriptions</i> | | | |
| Forestland Interpretations | 1/93 | 1/93 | SOI |
| <i>Use and Explanation of Forestland Interpretations</i> | | | |
| <i>*Woodland Management and Productivity</i> | | | |
| Nonagricultural Interpretations | 4/02 | 4/02 | SOI |
| <i>*Building Site Development Report</i> | | | |
| <i>*Construction Materials Report</i> | | | |
| Recreation Interpretations | 4/02 | 4/02 | SOI |
| <i>*Recreational Interpretations</i> | | | |
| Wildlife Interpretations | 4/02 | 4/02 | SOI |
| <i>*Wildlife Interpretations Report</i> | | | |

| | 4/02 | 4/02 | SOI |
|--|---------------|------------------------|----------------------|
| | Issue Date | Date of Last Review | Responsible Staff |
| Pastureland and Hayland Interpretations <i>*Yields Per Acre of Pasture and Hayland</i> | | | |
| Mined Land Interpretations <i>Use and Explanation of Mined Land Interpretations</i> | | | |
| Windbreak Interpretations <i>*Conservation Tree and Shrub Management Report</i> | 4/02 | 4/02 | SOI |
| Engineering Interpretations <i>*Engineering Index Properties</i> <i>*Physical Properties of the Soils</i> <i>*Chemical Properties of the Soils</i> <i>*Water Features</i> <i>*Soil Features</i> <i>*Water Management Report</i> | 4/02 | 4/02 | SOI |
| Waste Disposal Interpretations <i>*Sanitary Facilities Report</i> <i>*Agricultural Waste Management Report</i> | 4/02 | 4/02 | SOI |
| Hydric Soil Interpretations <i>Use and Explanation of Hydric Soil Interpretations</i> <i>*Hydric Soils List</i> | 4/02 | 4/02 | SOI |
| HEL Interpretations <i>Use and Explanation of Highly Erodible Land Interpretations</i> <i>*Highly Erodible Lands Report</i> 1 <i>Highly Erodible Land</i> 2 <i>Potentially Highly Erodible Land</i> 3 <i>Nonhighly Erodible Land</i> | See Report | | SOI |

**County specific computer generated reports.*

ACREAGE AND PROPORTIONATE EXTENT OF THE SOILS

Kearney County, Nebraska: Published

| Map symbol | Soil name | Acres | Percent |
|------------|--|---------|---------|
| Ad | Alda Loam, 0 To 1 Percent Slopes----- | 7,540 | 2.3 |
| Bo | Boel Fine Sandy Loam, 0 To 1 Percent Slopes----- | 2,930 | 0.9 |
| Bu | Butler Silt Loam, 0 To 1 Percent Slopes----- | 12,580 | 3.8 |
| CaC | Coly Silt Loam, 3 To 6 Percent Slopes----- | 8,850 | 2.7 |
| CaD | Coly Silt Loam, 6 To 11 Percent Slopes----- | 1,640 | 0.5 |
| CaF | Coly Silt Loam, 11 To 20 Percent Slopes----- | 1,960 | 0.6 |
| CkB | Coly-Kenesaw Silt Loams, 0 To 3 Percent Slopes----- | 26,616 | 8.0 |
| CoD2 | Coly-Uly Silt Loams, 6 To 11 Percent Slopes, Eroded----- | 3,940 | 1.2 |
| CoF2 | Coly-Uly Silt Loams, 11 To 20 Percent Slopes, Eroded----- | 560 | 0.2 |
| De | Detroit Silt Loam, 0 To 1 Percent Slopes----- | 22,086 | 6.7 |
| Fm | Fillmore Silt Loam, 0 To 1 Percent Slopes----- | 1,100 | 0.3 |
| Gb | Gibbon Loam, 0 To 1 Percent Slopes----- | 1,030 | 0.3 |
| Gc | Gibbon Loam, Saline, 0 To 1 Percent Slopes----- | 990 | 0.3 |
| Go | Gothenburg Loamy Sand, 0 To 2 Percent Slopes----- | 1,590 | 0.5 |
| GP | Gravel Pit----- | 136 | * |
| HeB | Hersh Fine Sandy Loam, 0 To 3 Percent Slopes----- | 2,460 | 0.7 |
| HeC | Hersh Fine Sandy Loam, 3 To 6 Percent Slopes----- | 2,490 | 0.8 |
| HeD | Hersh Fine Sandy Loam, 6 To 11 Percent Slopes----- | 950 | 0.3 |
| Hf | Hobbs Silt Loam, 0 To 2 Percent Slopes----- | 1,640 | 0.5 |
| HgB | Hobbs Silt Loam, Channeled----- | 1,150 | 0.3 |
| Ho | Holdrege Silt Loam, 0 To 1 Percent Slopes----- | 81,280 | 24.5 |
| HoB | Holdrege Silt Loam, 1 To 3 Percent Slopes----- | 22,156 | 6.7 |
| HoC | Holdrege Silt Loam, 3 To 6 Percent Slopes----- | 800 | 0.2 |
| HoC2 | Holdrege Silt Loam, 3 To 6 Percent Slopes, Eroded----- | 4,460 | 1.3 |
| Hp | Holdrege Silt Loam, Overblown, 0 To 1 Percent Slopes----- | 11,030 | 3.3 |
| HpB | Holdrege Silt Loam, Overblown, 1 To 3 Percent Slopes----- | 890 | 0.3 |
| Hr | Hord Silt Loam, 0 To 1 Percent Slopes----- | 3,930 | 1.2 |
| InB | Inavale Loamy Fine Sand, 0 To 3 Percent Slopes----- | 2,060 | 0.6 |
| Ke | Kenesaw Silt Loam, 0 To 1 Percent Slopes----- | 29,416 | 8.9 |
| KgB | Kenesaw-Coly Silt Loams, 1 To 3 Percent Slopes----- | 3,610 | 1.1 |
| Lf | Lex Loam, 0 To 1 Percent Slopes----- | 1,080 | 0.3 |
| Lg | Lex Loam, Saline, 0 To 1 Percent Slopes----- | 360 | 0.1 |
| LoB | Libory Loamy Fine Sand, 0 To 3 Percent Slopes----- | 3,280 | 1.0 |
| M-W | Miscellaneous Water (sewage Lagoons)----- | 26 | * |
| Ma | Massie Silty Clay Loam, 0 To 1 Percent Slopes----- | 980 | 0.3 |
| Pm | Platte Loam, 0 To 1 Percent Slopes----- | 1,970 | 0.6 |
| Ru | Rusco Silt Loam, 0 To 1 Percent Slopes----- | 1,640 | 0.5 |
| Sc | Scott Silt Loam, 0 To 1 Percent Slopes----- | 2,070 | 0.6 |
| SmB | Simeon Sandy Loam, 0 To 3 Percent Slopes----- | 9,180 | 2.8 |
| To | Tryon Loamy Fine Sand, 0 To 1 Percent Slopes----- | 990 | 0.3 |
| UcF | Uly-Coly Silt Loams, 11 To 20 Percent Slopes----- | 2,290 | 0.7 |
| UsF | Ustorthents, Steep----- | 200 | * |
| VaB | Valentine Loamy Fine Sand, 0 To 3 Percent Slopes----- | 4,690 | 1.4 |
| VaD | Valentine Loamy Fine Sand, 3 To 9 Percent Slopes----- | 9,460 | 2.9 |
| VaF | Valentine Loamy Fine Sand, Rolling----- | 18,540 | 5.6 |
| VbD | Valentine-Els Loamy Fine Sands, 0 To 9 Percent Slopes----- | 6,840 | 2.1 |
| W | Water----- | 805 | 0.2 |
| Wa | Wann Fine Sandy Loam, 0 To 1 Percent Slopes----- | 5,640 | 1.7 |
| | Total----- | 331,911 | 100.0 |

* Less than 0.1 percent.

NONTECHNICAL SOIL DESCRIPTIONS
Kearney County, Nebraska

Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units, shown in the NonTechnical Descriptions report. These descriptions are written in terminology that Non-technical users of soil survey information can understand. Nontechnical soil descriptions are a powerful tool for creating reports. These high quality, easy to read reports can be generated by conservation planners and other NRCS employees for distribution to land users. Soil map unit descriptions and National Soil Information System records are the basis for these descriptions.

Ad Alda Loam, 0 To 1 Percent Slopes

Alda soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Bo Boel Fine Sandy Loam, 0 To 1 Percent Slopes

Boel soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium and/or sandy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Bu Butler Silt Loam, 0 To 1 Percent Slopes

Butler soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level playa on upland. The runoff class is negligible. The parent material consists of loess. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 12 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clayey - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

CaC Coly Silt Loam, 3 To 6 Percent Slopes

Coly soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is low. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

CaD Coly Silt Loam, 6 To 11 Percent Slopes

Coly soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

CaF Coly Silt Loam, 11 To 20 Percent Slopes

Coly soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

CkB Coly-Kenesaw Silt Loams, 0 To 3 Percent Slopes

Coly soil makes up 60 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

Kenesaw soil makes up 40 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping hillslope on upland. The runoff class is low. The parent material consists of calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

CoD2 Coly-Uly Silt Loams, 6 To 11 Percent Slopes, Eroded

Coly soil makes up 60 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Uly soil makes up 40 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

CoF2 Coly-Uly Silt Loams, 11 To 20 Percent Slopes, Eroded

Coly soil makes up 60 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Uly soil makes up 40 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

De Detroit Silt Loam, 0 To 1 Percent Slopes

Detroit soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flat on upland. The runoff class is medium. The parent material consists of silty alluvium over loess. This soil is moderately well drained. The slowest permeability is slow. It has a high available water capacity and a high shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 1.

Fm Fillmore Silt Loam, 0 To 1 Percent Slopes

Fillmore soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level playa on upland. The runoff class is negligible. The parent material consists of loess. This soil is somewhat poorly drained. The slowest permeability is very slow. It has a high available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 12 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 3w.

Gb Gibbon Loam, 0 To 1 Percent Slopes

Gibbon soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

Gc Gibbon Loam, Saline, 0 To 1 Percent Slopes

Gibbon soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of stratified calcareous silty alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 15 percent calcium carbonate. This soil contains a moderately saline horizon, it has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3s. It is in the nonirrigated land capability classification 4s.

Go Gothenburg Loamy Sand, 0 To 2 Percent Slopes

Gothenburg soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of sandy and gravelly alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a very low available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It is in the nonirrigated land capability classification 7s.

HeB Hersh Fine Sandy Loam, 0 To 3 Percent Slopes

Hersh soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping hillslope on upland. The runoff class is very low. The parent material consists of coarse-loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 3e.

HeC Hersh Fine Sandy Loam, 3 To 6 Percent Slopes

Hersh soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is very low. The parent material consists of coarse-loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

HeD Hersh Fine Sandy Loam, 6 To 11 Percent Slopes

Hersh soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hillslope on upland. The runoff class is low. The parent material consists of coarse-loamy eolian deposits. This soil is well drained. The slowest permeability is moderately rapid. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

Hf Hobbs Silt Loam, 0 To 2 Percent Slopes

Hobbs soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on drainageway on upland. The runoff class is negligible. The parent material consists of stratified silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty Overflow - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

HgB Hobbs Silt Loam, Channeled

Hobbs soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on drainageway on upland. The runoff class is low. The parent material consists of stratified silty alluvium. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is frequently flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty Overflow - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6w.

Ho Holdrege Silt Loam, 0 To 1 Percent Slopes

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level broad interstream divide on upland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1 It is in the nonirrigated land capability classification 2c.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

HoB Holdrege Silt Loam, 1 To 3 Percent Slopes

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

HoC Holdrege Silt Loam, 3 To 6 Percent Slopes

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

HoC2 Holdrege Silt Loam, 3 To 6 Percent Slopes, Eroded

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping hillslope on upland. The runoff class is medium. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Hp Holdrege Silt Loam, Overblown, 0 To 1 Percent Slopes

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level summit interfluvium on upland. The runoff class is negligible. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

HpB Holdrege Silt Loam, Overblown, 1 To 3 Percent Slopes

Holdrege soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of loess. This soil is well drained. The slowest permeability is moderately slow. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Hr Hord Silt Loam, 0 To 1 Percent Slopes

Hord soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level broad interstream divide on upland. The runoff class is negligible. The parent material consists of alluvium and/or loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

InB Inavale Loamy Fine Sand, 0 To 3 Percent Slopes

Inavale soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping flood plain on river valley. The runoff class is negligible. The parent material consists of sandy alluvium. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 4e.

Ke Kenesaw Silt Loam, 0 To 1 Percent Slopes

Kenesaw soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level summit interfluvium on upland. The runoff class is negligible. The parent material consists of calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 1. It is in the nonirrigated land capability classification 2c.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

KgB Kenesaw-Coly Silt Loams, 1 To 3 Percent Slopes

Kenesaw soil makes up 55 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Coly soil makes up 45 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a gently sloping hillslope on upland. The runoff class is low. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2e. It is in the nonirrigated land capability classification 2e.

Lf Lex Loam, 0 To 1 Percent Slopes

Lex soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 24 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a very slightly saline horizon, it has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3w. It is in the nonirrigated land capability classification 3w.

Lg Lex Loam, Saline, 0 To 1 Percent Slopes

Lex soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is low. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately slow. It has a moderate available water capacity and a moderate shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 27 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil contains a moderately saline horizon, it has a horizon that is moderately sodic. This soil is in the Saline Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3s. It is in the nonirrigated land capability classification 4s.

LoB Libory Loamy Fine Sand, 0 To 3 Percent Slopes

Libory soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping hillslope on upland, stream terrace on river valley. The runoff class is very low. The parent material consists of eolian sands over loess. This soil is moderately well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Sandy Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 3e. It is in the nonirrigated land capability classification 3e.

Ma Massie Silty Clay Loam, 0 To 1 Percent Slopes

Massie soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level playa on upland. The runoff class is negligible. The parent material consists of alluvium over loess. This soil is very poorly drained. The slowest permeability is very slow. It has a high available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. It is in the nonirrigated land capability classification 8w.

Pm Platte Loam, 0 To 1 Percent Slopes

Platte soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of loamy alluvium over sandy and gravelly alluvium. This soil is somewhat poorly drained. The slowest permeability is moderate. It has a low available water capacity and a low shrink swell potential. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 24 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

Ru Rusco Silt Loam, 0 To 1 Percent Slopes

Rusco soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level depression on upland. The runoff class is negligible. The parent material consists of loamy eolian deposits over silty alluvium. This soil is moderately well drained. The slowest permeability is moderately slow. It has a high available water capacity and a moderate shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Silty Lowland - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

Sc Scott Silt Loam, 0 To 1 Percent Slopes

Scott soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level playa on upland. The runoff class is negligible. The parent material consists of loess. This soil is poorly drained. The slowest permeability is very slow. It has a high available water capacity and a very high shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Clayey Overflow - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 4w.

SmB Simeon Sandy Loam, 0 To 3 Percent Slopes

Simeon soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping stream terrace on river valley. The runoff class is very low. The parent material consists of sandy and gravelly alluvium. This soil is excessively drained. The slowest permeability is moderately rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Shallow To Gravel - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4s. It is in the nonirrigated land capability classification 6s.

To Tryon Loamy Fine Sand, 0 To 1 Percent Slopes

Tryon soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level swale on flood plain on river valley. The runoff class is negligible. The parent material consists of sandy eolian deposits over sandy alluvium. This soil is poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 9 inches. The soil contains a maximum amount of 5 percent calcium carbonate. This soil is in the Wet Subirrigated - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 5w.

UcF Uly-Coly Silt Loams, 11 To 20 Percent Slopes

Uly soil makes up 60 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. This soil is in the Silty - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

Coly soil makes up 40 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on upland. The runoff class is medium. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a very high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 10 percent calcium carbonate. This soil is in the Limy Upland - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

UsF Ustorthents, Steep

Ustorthents soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately steep to steep spoil bank. The runoff class is high. The parent material consists of fine-silty calcareous loess. This soil is well drained. The slowest permeability is moderate. It has a high available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. The soil contains a maximum amount of 15 percent calcium carbonate. It is in the nonirrigated land capability classification 7s.

VaB Valentine Loamy Fine Sand, 0 To 3 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping hummock on sand sheet on upland. The runoff class is negligible. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sandy - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 4e.

VaD Valentine Loamy Fine Sand, 3 To 9 Percent Slopes

Valentine soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hummock on sand sheet on upland. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

NONTECHNICAL SOIL DESCRIPTIONS--Continued
Kearney County, Nebraska

VaF Valentine Loamy Fine Sand, Rolling

Valentine soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a strongly sloping to moderately steep hillslope on sand sheet on upland. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. It is in the nonirrigated land capability classification 6e.

VbD Valentine-Els Loamy Fine Sands, 0 To 9 Percent Slopes

Valentine soil makes up 65 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a moderately sloping to strongly sloping hummock on sand sheet on upland. The runoff class is very low. The parent material consists of eolian sands. This soil is excessively drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The seasonal high water table is at a depth of more than 6 feet. This soil is in the Sands - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4e. It is in the nonirrigated land capability classification 6e.

Els soil makes up 35 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level to gently sloping swale on sand sheet on upland. The runoff class is negligible. The parent material consists of sandy alluvium and/or sandy eolian deposits. This soil is somewhat poorly drained. The slowest permeability is rapid. It has a low available water capacity and a low shrink swell potential. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 4w. It is in the nonirrigated land capability classification 4w.

Wa Wann Fine Sandy Loam, 0 To 1 Percent Slopes

Wann soil makes up 100 percent of the map unit. This map unit is in the Central Loess Plains Major Land Resource Area. This soil occurs on a nearly level flood plain on river valley. The runoff class is negligible. The parent material consists of calcareous loamy alluvium. This soil is somewhat poorly drained. The slowest permeability is moderately rapid. It has a moderate available water capacity and a low shrink swell potential. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 30 inches. The soil contains a maximum amount of 5 percent calcium carbonate. It has a horizon that is slightly sodic. This soil is in the Subirrigated - Veg. Zone 3 range site. This soil is in the irrigated land capability class 2w. It is in the nonirrigated land capability classification 2w.

Ad—Alda loam, 0 to 1 percent slopes

Map Unit Composition

Alda: 100 percent

Component Descriptions

Alda

MLRA: 75 - Central Loess Plains*Landform:* Flood plain on river valley*Parent material:* Loamy alluvium over sandy and gravelly alluvium*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* Moderate (About 6.3 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Rare*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Negligible*Ecological site:* Subirrigated - Veg. Zone 3*Land capability (irrigated):* 3w*Land capability (nonirrigated):* 3w*Typical Profile:*

H1—0 to 12 inches; loam

H2—12 to 30 inches; fine sandy loam

H3—30 to 60 inches; coarse sand

Minor Components**Wt At 0-1 Foot****Bo—Boel fine sandy loam, 0 to 1 percent slopes**

Map Unit Composition

Boel: 100 percent

Component Descriptions

Boel

MLRA: 75 - Central Loess Plains*Landform:* Flood plain on river valley*Parent material:* Loamy alluvium and/or sandy alluvium*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Moderately rapid (About 2.00 in/hr)*Available water capacity:* Moderate (About 6.2 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Rare*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Negligible*Ecological site:* Subirrigated - Veg. Zone 3*Land capability (irrigated):* 3w*Land capability (nonirrigated):* 3w*Typical Profile:*

H1—0 to 17 inches; fine sandy loam

H2—17 to 54 inches; fine sand

H3—54 to 60 inches; loamy fine sand

Minor Components**Wt At 0-1 Foot****Bu—Butler silt loam, 0 to 1 percent slopes**

Map Unit Composition

Butler: 100 percent

Component Descriptions

Butler

MLRA: 75 - Central Loess Plains*Landform:* Playa on upland*Parent material:* Loess*Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Very slow (About 0.01 in/hr)*Available water capacity:* High (About 10.1 inches)*Shrink-swell potential:* Very high (About 10.0 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 6 to 18 inches*Runoff class:* Negligible*Ecological site:* Clayey - Veg. Zone 3*Land capability (irrigated):* 2w*Land capability (nonirrigated):* 2w*Typical Profile:*

H1—0 to 12 inches; silt loam

H2—12 to 35 inches; silty clay

H3—35 to 41 inches; silty clay loam
H4—41 to 60 inches; silt loam

Minor Components

Fillmore

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Clayey Overflow - Veg. Zone
3

Scott

Slope: 0 to 1 percent
Drainage class: Poorly drained
Ecological site: Clayey Overflow - Veg. Zone
3

CaC—Coly silt loam, 3 to 6 percent slopes

Map Unit Composition

Coly: 100 percent

Component Descriptions

Coly
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Limy Upland - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 5 inches; silt loam
H2—5 to 60 inches; silt loam

Minor Components

Wt At 0-1 Foot

CaD—Coly silt loam, 6 to 11 percent slopes

Map Unit Composition

Coly: 100 percent

Component Descriptions

Coly
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 6 to 11 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Limy Upland - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 5 inches; silt loam
H2—5 to 60 inches; silt loam

CaF—Coly silt loam, 11 to 20 percent slopes

Map Unit Composition

Coly: 100 percent

Component Descriptions

Coly
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 11 to 20 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Limy Upland - Veg. Zone 3

Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 4 inches; silt loam

H2—4 to 60 inches; silt loam

CkB—Coly-Kenesaw silt loams, 0 to 3 percent slopes

Map Unit Composition

Coly: 60 percent

Kenesaw: 40 percent

Component Descriptions

Coly

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Fine-silty calcareous loess

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Limy Upland - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 6 inches; silt loam

H2—6 to 60 inches; silt loam

Kenesaw

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Calcareous loess

Slope: 0 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Very high (About 12.0 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 10 inches; silt loam

H2—10 to 23 inches; silt loam

H3—23 to 60 inches; silt loam

Minor Components

Fillmore

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

CoD2—Coly-Uly silt loams, 6 to 11 percent slopes, Eroded

Map Unit Composition

Coly: 60 percent

Uly: 40 percent

Component Descriptions

Coly

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Fine-silty calcareous loess

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium

Ecological site: Limy Upland - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 5 inches; silt loam

H2—5 to 60 inches; silt loam

Uly

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 6 to 11 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.3 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 4e
Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 7 inches; silt loam
H2—7 to 28 inches; silt loam
H3—28 to 60 inches; silt loam

CoF2—Coly-Uly silt loams, 11 to 20 percent slopes, Eroded

Map Unit Composition

Coly: 60 percent
Uly: 40 percent

Component Descriptions

Coly
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 11 to 20 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Limy Upland - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 6 inches; silt loam
H2—6 to 60 inches; silt loam

Uly

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 11 to 20 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.3 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 7 inches; silt loam
H2—7 to 28 inches; silt loam
H3—28 to 60 inches; silt loam

De—Detroit silt loam, 0 to 1 percent slopes

Map Unit Composition

Detroit: 100 percent

Component Descriptions

Detroit
MLRA: 75 - Central Loess Plains
Landform: Flat on upland
Parent material: Silty alluvium over loess
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Slow (About 0.06 in/hr)
Available water capacity: High (About 11.2 inches)
Shrink-swell potential: High (About 7.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty Lowland - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 1

Typical Profile:

H1—0 to 16 inches; silt loam
H2—16 to 38 inches; silty clay loam

H3—38 to 60 inches; silty clay loam

Minor Components

Fillmore

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Ecological site: Clayey Overflow - Veg. Zone

3

Fm—Fillmore silt loam, 0 to 1 percent slopes

Map Unit Composition

Fillmore: 100 percent

Component Descriptions

Fillmore

MLRA: 75 - Central Loess Plains

Landform: Playa on upland

Parent material: Loess

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 9.9 inches)

Shrink-swell potential: Very high (About 10.0 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 0 to 24 inches

Runoff class: Negligible

Ecological site: Clayey Overflow - Veg. Zone 3

Land capability (irrigated): 4w

Land capability (nonirrigated): 3w

Typical Profile:

H1—0 to 12 inches; silt loam

H2—12 to 35 inches; silty clay

H3—35 to 44 inches; silty clay loam

H4—44 to 60 inches; silt loam

Minor Components

Scott

Slope: 0 to 1 percent

Drainage class: Poorly drained

Ecological site: Clayey Overflow - Veg. Zone

3

Gb—Gibbon loam, 0 to 1 percent slopes

Map Unit Composition

Gibbon: 100 percent

Component Descriptions

Gibbon

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Stratified calcareous silty alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderate (About 0.57 in/hr)

Available water capacity: High (About 11.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: About 18 to 36 inches

Runoff class: Negligible

Ecological site: Subirrigated - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 14 inches; loam

H2—14 to 37 inches; silt loam

H3—37 to 60 inches; stratified very fine sandy loam to silt loam

Minor Components

Wt At 0-1 Foot

Gc—Gibbon loam, Saline, 0 to 1 percent slopes

Map Unit Composition

Gibbon: 100 percent

Component Descriptions

Gibbon

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Stratified calcareous silty alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Moderate (About 3.0 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 18 to 36 inches
Runoff class: Negligible
Ecological site: Subirrigated - Veg. Zone 3
Land capability (irrigated): 3s
Land capability (nonirrigated): 4s

Typical Profile:

H1—0 to 10 inches; loam
 H2—10 to 40 inches; silt loam
 H3—40 to 60 inches; silty clay loam

Minor Components**Wt At 0-1 Foot****Go—Gothenburg loamy sand, 0 to 2 percent slopes**

Map Unit Composition

Gothenburg: 100 percent

Component Descriptions

Gothenburg
MLRA: 75 - Central Loess Plains
Landform: Flood plain on river valley
Parent material: Sandy and gravelly alluvium
Slope: 0 to 2 percent
Drainage class: Poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Very low (About 2.2 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Frequent
Depth to seasonal water saturation: About 0 to 18 inches
Runoff class: Negligible
Land capability (nonirrigated): 7s

Typical Profile:

H1—0 to 3 inches; loamy sand
 H2—3 to 8 inches; sand
 H3—8 to 60 inches; stratified coarse sand to gravelly coarse sand

GP—Gravel Pit

Map Unit Composition

Pits: 100 percent

Component Descriptions

Pits
MLRA: 75 - Central Loess Plains
Slope: 0 to 30 percent
Drainage class: Excessively drained
Slowest permeability: Rapid (About 6.00 in/hr)
Available water capacity: Low (About 3.5 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Land capability (nonirrigated): 8s

HeB—Hersh fine sandy loam, 0 to 3 percent slopes

Map Unit Composition

Hersh: 100 percent

Component Descriptions

Hersh
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Coarse-loamy eolian deposits
Slope: 0 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: High (About 9.6 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Sandy - Veg. Zone 3
Land capability (irrigated): 2e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; fine sandy loam
 H2—7 to 11 inches; fine sandy loam

H3—11 to 60 inches; fine sandy loam

Minor Components

Fillmore

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

HeC—Hersh fine sandy loam, 3 to 6 percent slopes

Map Unit Composition

Hersh: 100 percent

Component Descriptions

Hersh

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Coarse-loamy eolian deposits

Slope: 3 to 6 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: High (About 9.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Very low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 7 inches; fine sandy loam

H2—7 to 11 inches; fine sandy loam

H3—11 to 60 inches; fine sandy loam

HeD—Hersh fine sandy loam, 6 to 11 percent slopes

Map Unit Composition

Hersh: 100 percent

Component Descriptions

Hersh

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Coarse-loamy eolian deposits

Slope: 6 to 11 percent

Drainage class: Well drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: High (About 9.6 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Sandy - Veg. Zone 3

Land capability (irrigated): 4e

Land capability (nonirrigated): 4e

Typical Profile:

H1—0 to 7 inches; fine sandy loam

H2—7 to 11 inches; fine sandy loam

H3—11 to 60 inches; fine sandy loam

Hf—Hobbs silt loam, 0 to 2 percent slopes

Map Unit Composition

Hobbs: 100 percent

Component Descriptions

Hobbs

MLRA: 75 - Central Loess Plains

Landform: Flood plain on drainageway on upland

Parent material: Stratified silty alluvium

Slope: 0 to 2 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Occasional

Depth to seasonal water saturation: More than 6 feet

Runoff class: Negligible

Ecological site: Silty Overflow - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 6 inches; silt loam
 H2—6 to 34 inches; stratified silt loam
 H3—34 to 60 inches; silt loam

Minor Components**Wt At 0-1 Foot****HgB—Hobbs silt loam, Channeled**

Map Unit Composition

Hobbs: 100 percent

Component Descriptions

Hobbs

MLRA: 75 - Central Loess Plains

Landform: Flood plain on drainageway on upland

Parent material: Stratified silty alluvium

Slope: 0 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.8 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Frequent

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty Overflow - Veg. Zone 3

Land capability (nonirrigated): 6w

Typical Profile:

H1—0 to 9 inches; silt loam

H2—9 to 34 inches; stratified silt loam

H3—34 to 60 inches; silt loam

Ho—Holdrege silt loam, 0 to 1 percent slopes

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege

MLRA: 75 - Central Loess Plains

Landform: Broad interstream divide on upland

Parent material: Loess

Slope: 0 to 1 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Very high (About 12.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Silty - Veg. Zone 3

Land capability (irrigated): 1

Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 13 inches; silt loam

H2—13 to 25 inches; silty clay loam

H3—25 to 31 inches; silt loam

H4—31 to 60 inches; silt loam

Minor Components**Fillmore**

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Ecological site: Clayey Overflow - Veg. Zone 3

HoB—Holdrege silt loam, 1 to 3 percent slopes

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Loess

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Very high (About 12.3 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 2e
Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 25 inches; silty clay loam
 H3—25 to 31 inches; silt loam
 H4—31 to 60 inches; silt loam

Minor Components**Fillmore**

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

HoC—Holdrege silt loam, 3 to 6 percent slopes

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Loess
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Very high (About 12.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 25 inches; silty clay loam
 H3—25 to 31 inches; silt loam
 H4—31 to 60 inches; silt loam

HoC2—Holdrege silt loam, 3 to 6 percent slopes, Eroded

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Loess
Slope: 3 to 6 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Very high (About 12.3 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 3e
Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 25 inches; silty clay loam
 H3—25 to 31 inches; silt loam
 H4—31 to 60 inches; silt loam

Hp—Holdrege silt loam, Overblown, 0 to 1 percent slopes

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege
MLRA: 75 - Central Loess Plains
Landform: Interfluvium on upland
Hillslope position: Summit
Parent material: Loess
Slope: 0 to 1 percent
Drainage class: Well drained

Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Very high (About 12.6 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 27 inches; silt loam
 H2—27 to 43 inches; silty clay loam
 H3—43 to 47 inches; silt loam
 H4—47 to 60 inches; silt loam

Minor Components**Fillmore**

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

HpB—Holdrege silt loam, Overblown, 1 to 3 percent slopes

Map Unit Composition

Holdrege: 100 percent

Component Descriptions

Holdrege

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Loess
Slope: 1 to 3 percent
Drainage class: Well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: Very high (About 12.6 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Low
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

H1—0 to 27 inches; silt loam
 H2—27 to 43 inches; silty clay loam
 H3—43 to 47 inches; silt loam
 H4—47 to 60 inches; silt loam

Minor Components**Fillmore**

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

Hr—Hord silt loam, 0 to 1 percent slopes

Map Unit Composition

Hord: 100 percent

Component Descriptions

Hord

MLRA: 75 - Central Loess Plains
Landform: Broad interstream divide on upland
Parent material: Alluvium and/or loess
Slope: 0 to 1 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: Very high (About 12.1 inches)
Shrink-swell potential: Moderate (About 3.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Negligible
Ecological site: Silty - Veg. Zone 3
Land capability (irrigated): 1
Land capability (nonirrigated): 2c

Typical Profile:

H1—0 to 13 inches; silt loam
 H2—13 to 43 inches; silt loam
 H3—43 to 60 inches; silt loam

Minor Components**Fillmore**

Slope: 0 to 1 percent
Drainage class: Somewhat poorly drained
Ecological site: Clayey Overflow - Veg. Zone 3

InB—Inavale loamy fine sand, 0 to 3 percent slopes

Map Unit Composition

Inavale: 100 percent

Component Descriptions

Inavale

MLRA: 75 - Central Loess Plains*Landform:* Flood plain on river valley*Parent material:* Sandy alluvium*Slope:* 0 to 3 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 4.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* Rare*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy Lowland - Veg. Zone 3*Land capability (irrigated):* 3e*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 5 inches; loamy fine sand

H2—5 to 10 inches; loamy fine sand

H3—10 to 60 inches; fine sand

Minor Components**Tryon***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**Ke—Kenesaw silt loam, 0 to 1 percent slopes**

Map Unit Composition

Kenesaw: 100 percent

Component Descriptions

Kenesaw

MLRA: 75 - Central Loess Plains*Landform:* Interfluvium on upland*Hillslope position:* Summit*Parent material:* Calcareous loess*Slope:* 0 to 1 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 12.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Silty - Veg. Zone 3*Land capability (irrigated):* 1*Land capability (nonirrigated):* 2c*Typical Profile:*

H1—0 to 8 inches; silt loam

H2—8 to 22 inches; silt loam

H3—22 to 60 inches; silt loam

Minor Components**Fillmore***Slope:* 0 to 1 percent*Drainage class:* Somewhat poorly drained*Ecological site:* Clayey Overflow - Veg. Zone 3**KgB—Kenesaw-Coly silt loams, 1 to 3 percent slopes**

Map Unit Composition

Kenesaw: 55 percent

Coly: 45 percent

Component Descriptions

Kenesaw

MLRA: 75 - Central Loess Plains*Landform:* Hillslope on upland*Parent material:* Calcareous loess*Slope:* 1 to 3 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 12.0 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Low*Ecological site:* Silty - Veg. Zone 3*Land capability (irrigated):* 2e

Land capability (nonirrigated): 2e

Typical Profile:

- H1—0 to 7 inches; silt loam
- H2—7 to 18 inches; silt loam
- H3—18 to 60 inches; silt loam

Coly

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland

Parent material: Fine-silty calcareous loess

Slope: 1 to 3 percent

Drainage class: Well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 11.9 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: None

Depth to seasonal water saturation: More than 6 feet

Runoff class: Low

Ecological site: Limy Upland - Veg. Zone 3

Land capability (irrigated): 2e

Land capability (nonirrigated): 2e

Typical Profile:

- H1—0 to 5 inches; silt loam
- H2—5 to 60 inches; silt loam

Lf—Lex loam, 0 to 1 percent slopes

Map Unit Composition

Lex: 100 percent

Component Descriptions

Lex

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Moderate (About 6.2 inches)

Shrink-swell potential: Low (About 2.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: About 12 to 36 inches

Runoff class: Low

Ecological site: Subirrigated - Veg. Zone 3

Land capability (irrigated): 3w

Land capability (nonirrigated): 3w

Typical Profile:

- H1—0 to 11 inches; loam
- H2—11 to 24 inches; stratified sandy loam to silty clay loam
- H3—24 to 60 inches; coarse sand

Minor Components

Wt At 0-1 Foot

Lg—Lex loam, Saline, 0 to 1 percent slopes

Map Unit Composition

Lex: 100 percent

Component Descriptions

Lex

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately slow (About 0.20 in/hr)

Available water capacity: Moderate (About 7.0 inches)

Shrink-swell potential: Moderate (About 4.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: About 18 to 36 inches

Runoff class: Low

Ecological site: Saline Subirrigated - Veg. Zone 3

Land capability (irrigated): 3s

Land capability (nonirrigated): 4s

Typical Profile:

- H1—0 to 12 inches; loam
- H2—12 to 31 inches; silt loam
- H3—31 to 60 inches; coarse sand

Minor Components

Wt At 0-1 Foot

LoB—Libory loamy fine sand, 0 to 3 percent slopes

Map Unit Composition

Libory: 100 percent

Component Descriptions

Libory

MLRA: 75 - Central Loess Plains

Landform: Hillslope on upland, stream terrace on river valley

Parent material: Eolian sands over loess

Slope: 0 to 3 percent

Drainage class: Moderately well drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: High (About 9.3 inches)

Shrink-swell potential: Low (About 2.0 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 18 to 36 inches

Runoff class: Very low

Ecological site: Sandy Lowland - Veg. Zone 3

Land capability (irrigated): 3e

Land capability (nonirrigated): 3e

Typical Profile:

H1—0 to 16 inches; loamy fine sand
H2—16 to 24 inches; loamy fine sand
H3—24 to 60 inches; silt loam

**Minor Components
Wt At 0-1 Foot****M-W—Miscellaneous Water
(sewage Lagoons)**

Map Unit Composition

Miscellaneous Water: 100 percent

Component Descriptions

Miscellaneous Water

MLRA: 75 - Central Loess Plains

Depth to seasonal water saturation: More than 6 feet

Ma—Massie silty clay loam, 0 to 1 percent slopes

Map Unit Composition

Massie: 100 percent

Component Descriptions

Massie

MLRA: 75 - Central Loess Plains

Landform: Playa on upland

Parent material: Alluvium over loess

Slope: 0 to 1 percent

Drainage class: Very poorly drained

Slowest permeability: Very slow (About 0.00 in/hr)

Available water capacity: High (About 9.6 inches)

Shrink-swell potential: Very high (About 9.0 LEP)

Flooding hazard: None

Depth to seasonal water saturation: About 0 to 12 inches

Runoff class: Negligible

Land capability (nonirrigated): 8w

Typical Profile:

H1—0 to 10 inches; silty clay loam
H2—10 to 60 inches; silty clay

Pm—Platte loam, 0 to 1 percent slopes

Map Unit Composition

Platte: 100 percent

Component Descriptions

Platte

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Loamy alluvium over sandy and gravelly alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderate (About 0.60 in/hr)

Available water capacity: Low (About 3.7 inches)

Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Occasional
Depth to seasonal water saturation: About 12 to 36 inches
Runoff class: Negligible
Ecological site: Subirrigated - Veg. Zone 3
Land capability (irrigated): 4w
Land capability (nonirrigated): 4w

Typical Profile:

H1—0 to 6 inches; loam
 H2—6 to 12 inches; loam
 H3—12 to 60 inches; gravelly coarse sand

Minor Components

Gothenburg

Slope: 0 to 2 percent
Drainage class: Poorly drained

Ru—Rusco silt loam, 0 to 1 percent slopes

Map Unit Composition

Rusco: 100 percent

Component Descriptions

Rusco
MLRA: 75 - Central Loess Plains
Landform: Depression on upland
Parent material: Loamy eolian deposits over silty alluvium
Slope: 0 to 1 percent
Drainage class: Moderately well drained
Slowest permeability: Moderately slow (About 0.20 in/hr)
Available water capacity: High (About 11.7 inches)
Shrink-swell potential: Moderate (About 4.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 0 to 0 inches
Runoff class: Negligible
Ecological site: Silty Lowland - Veg. Zone 3
Land capability (irrigated): 2w
Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 5 inches; silt loam
 H2—5 to 30 inches; silty clay loam
 H3—30 to 60 inches; silt loam

Minor Components
Perched Wt

Sc—Scott silt loam, 0 to 1 percent slopes

Map Unit Composition

Scott: 100 percent

Component Descriptions

Scott
MLRA: 75 - Central Loess Plains
Landform: Playa on upland
Parent material: Loess
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Very slow (About 0.00 in/hr)
Available water capacity: High (About 9.2 inches)
Shrink-swell potential: Very high (About 9.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: About 0 to 12 inches
Runoff class: Negligible
Ecological site: Clayey Overflow - Veg. Zone 3
Land capability (nonirrigated): 4w

Typical Profile:

H1—0 to 6 inches; silt loam
 H2—6 to 37 inches; clay
 H3—37 to 48 inches; silty clay loam
 H4—48 to 60 inches; silt loam

SmB—Simeon sandy loam, 0 to 3 percent slopes

Map Unit Composition

Simeon: 100 percent

Component Descriptions

Simeon
MLRA: 75 - Central Loess Plains
Landform: Stream terrace on river valley
Parent material: Sandy and gravelly alluvium

Slope: 0 to 3 percent
Drainage class: Excessively drained
Slowest permeability: Moderately rapid (About 2.00 in/hr)
Available water capacity: Low (About 5.1 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Very low
Ecological site: Shallow To Gravel - Veg. Zone 3
Land capability (irrigated): 4s
Land capability (nonirrigated): 6s

Typical Profile:

H1—0 to 6 inches; sandy loam
 H2—6 to 60 inches; coarse sand

To—Tryon loamy fine sand, 0 to 1 percent slopes

Map Unit Composition

Tryon: 100 percent

Component Descriptions

Tryon
MLRA: 75 - Central Loess Plains
Landform: Swale on flood plain on river valley
Parent material: Sandy eolian deposits over sandy alluvium
Slope: 0 to 1 percent
Drainage class: Poorly drained
Slowest permeability: Rapid (About 5.95 in/hr)
Available water capacity: Low (About 4.3 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: Rare
Depth to seasonal water saturation: About 0 to 18 inches
Runoff class: Negligible
Ecological site: Wet Subirrigated - Veg. Zone 3
Land capability (nonirrigated): 5w

Typical Profile:

H1—0 to 4 inches; loamy fine sand
 H2—4 to 60 inches; fine sand

UcF—Uly-Coly silt loams, 11 to 20 percent slopes

Map Unit Composition

Uly: 60 percent
 Coly: 40 percent

Component Descriptions

Uly
MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 11 to 20 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.4 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Silty - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 9 inches; silt loam
 H2—9 to 27 inches; silt loam
 H3—27 to 60 inches; silt loam

Coly

MLRA: 75 - Central Loess Plains
Landform: Hillslope on upland
Parent material: Fine-silty calcareous loess
Slope: 11 to 20 percent
Drainage class: Well drained
Slowest permeability: Moderate (About 0.60 in/hr)
Available water capacity: High (About 11.9 inches)
Shrink-swell potential: Low (About 1.5 LEP)
Flooding hazard: None
Depth to seasonal water saturation: More than 6 feet
Runoff class: Medium
Ecological site: Limy Upland - Veg. Zone 3
Land capability (nonirrigated): 6e

Typical Profile:

H1—0 to 4 inches; silt loam
 H2—4 to 60 inches; silt loam

UsF—Ustorthents, Steep

Map Unit Composition

Ustorhents: 100 percent

Component Descriptions

Ustorhents

MLRA: 75 - Central Loess Plains*Landform:* Spoil bank*Parent material:* Fine-silty calcareous loess*Slope:* 17 to 30 percent*Drainage class:* Well drained*Slowest permeability:* Moderate (About 0.60 in/hr)*Available water capacity:* High (About 11.8 inches)*Shrink-swell potential:* Low (About 2.9 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* High*Land capability (nonirrigated):* 7s*Typical Profile:*

H1—0 to 60 inches; silt loam

VaB—Valentine loamy fine sand, 0 to 3 percent slopes

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

MLRA: 75 - Central Loess Plains*Landform:* Hummock on sand sheet on upland*Parent material:* Eolian sands*Slope:* 0 to 3 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Negligible*Ecological site:* Sandy - Veg. Zone 3*Land capability (irrigated):* 4e*Land capability (nonirrigated):* 4e*Typical Profile:*

H1—0 to 5 inches; loamy fine sand

H2—5 to 8 inches; loamy fine sand

H3—8 to 60 inches; fine sand

Minor Components**Tryon***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**VaD—Valentine loamy fine sand, 3 to 9 percent slopes**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

MLRA: 75 - Central Loess Plains*Landform:* Hummock on sand sheet on upland*Parent material:* Eolian sands*Slope:* 3 to 9 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands - Veg. Zone 3*Land capability (irrigated):* 4e*Land capability (nonirrigated):* 6e*Typical Profile:*

H1—0 to 5 inches; loamy fine sand

H2—5 to 8 inches; loamy fine sand

H3—8 to 60 inches; fine sand

Minor Components**Tryon***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**VaF—Valentine loamy fine sand, Rolling**

Map Unit Composition

Valentine: 100 percent

Component Descriptions

Valentine

MLRA: 75 - Central Loess Plains*Landform:* Hillslope on sand sheet on upland*Parent material:* Eolian sands*Slope:* 9 to 17 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands - Veg. Zone 3*Land capability (nonirrigated):* 6e*Typical Profile:*

H1—0 to 5 inches; loamy fine sand

H2—5 to 8 inches; loamy fine sand

H3—8 to 60 inches; fine sand

Minor Components**Tryon***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**VbD—Valentine-Els loamy fine sands, 0 to 9 percent slopes**

Map Unit Composition

Valentine: 65 percent

Els: 35 percent

Component Descriptions

Valentine

MLRA: 75 - Central Loess Plains*Landform:* Hummock on sand sheet on upland*Parent material:* Eolian sands*Slope:* 3 to 9 percent*Drainage class:* Excessively drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* More than 6 feet*Runoff class:* Very low*Ecological site:* Sands - Veg. Zone 3*Land capability (irrigated):* 4e*Land capability (nonirrigated):* 6e*Typical Profile:*

H1—0 to 5 inches; loamy fine sand

H2—5 to 8 inches; loamy fine sand

H3—8 to 60 inches; fine sand

Els

MLRA: 75 - Central Loess Plains*Landform:* Swale on sand sheet on upland*Parent material:* Sandy alluvium and/or sandy eolian deposits*Slope:* 0 to 3 percent*Drainage class:* Somewhat poorly drained*Slowest permeability:* Rapid (About 5.95 in/hr)*Available water capacity:* Low (About 3.9 inches)*Shrink-swell potential:* Low (About 1.5 LEP)*Flooding hazard:* None*Depth to seasonal water saturation:* About 18 to 36 inches*Runoff class:* Negligible*Ecological site:* Subirrigated - Veg. Zone 3*Land capability (irrigated):* 4w*Land capability (nonirrigated):* 4w*Typical Profile:*

H1—0 to 6 inches; loamy fine sand

H2—6 to 10 inches; loamy sand

H3—10 to 60 inches; fine sand

Minor Components**Tryon***Slope:* 0 to 1 percent*Drainage class:* Poorly drained*Ecological site:* Wet Subirrigated - Veg. Zone 3**W—Water**

Map Unit Composition

Water: 100 percent

Component Descriptions

Water

MLRA: 75 - Central Loess Plains*Depth to seasonal water saturation:* More than 6 feet

General Considerations: Water includes streams, lakes, ponds, and estuaries. These areas are covered with water in most years, at least during the period that is warm enough for plants to grow. Many areas are covered throughout the year.

Wa—Wann fine sandy loam, 0 to 1 percent slopes

Map Unit Composition

Wann: 100 percent

Component Descriptions

Wann

MLRA: 75 - Central Loess Plains

Landform: Flood plain on river valley

Parent material: Calcareous loamy alluvium

Slope: 0 to 1 percent

Drainage class: Somewhat poorly drained

Slowest permeability: Moderately rapid (About 2.00 in/hr)

Available water capacity: Moderate (About 8.2 inches)

Shrink-swell potential: Low (About 1.5 LEP)

Flooding hazard: Rare

Depth to seasonal water saturation: About 18 to 42 inches

Runoff class: Negligible

Ecological site: Subirrigated - Veg. Zone 3

Land capability (irrigated): 2w

Land capability (nonirrigated): 2w

Typical Profile:

H1—0 to 14 inches; fine sandy loam

H2—14 to 48 inches; fine sandy loam

H3—48 to 60 inches; stratified sandy clay loam to fine sand

Minor Components

Wt At 0-1 Foot

Land capability classification shows, in a general way, the suitability of soils for most kinds of field crops. Crops that require special management are excluded. The soils are grouped according to their limitations for field crops, the risk of damage if they are used for crops, and the way they respond to management. The criteria used in grouping the soils do not include major and generally expensive land-forming that would change slope, depth, or other characteristics of the soils, nor do they include possible but unlikely major reclamation projects. Capability classification is not a substitute for interpretations designed to show suitability and limitations of groups of soils for rangeland, for forestland, or for engineering purposes. In the capability system, soils are generally grouped at three levels: capability class, subclass, and unit.

Capability classes, the broadest groups, are designated by the numbers 1 through 8. The numbers indicate progressively greater limitations and narrower choices for practical use. The classes are defined as follows:

(Class 1) soils have slight limitations that restrict their use.

(Class 2) soils have moderate limitations that restrict the choice of plants or that require moderate conservation practices.

(Class 3) soils have severe limitations that restrict the choice of plants or that require special conservation practices, or both.

(Class 4) soils have very severe limitations that restrict the choice of plants or that require very careful management, or both.

(Class 5) soils are subject to little or no erosion but have other limitations, impractical to remove, that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 6) soils have severe limitations that make them generally unsuitable for cultivation and that restrict their use mainly to pasture, rangeland, forestland, or wildlife habitat.

(Class 7) soils have very severe limitations that make them unsuitable for cultivation and that restrict their use mainly to grazing, forestland, or wildlife habitat.

(Class 8) soils and miscellaneous areas have limitations that preclude commercial plant production and that restrict their use to recreational purposes, wildlife habitat, watershed, or esthetic purposes.

Capability subclasses are soil groups within one class. They are designated by adding a small letter, e, w, s, or c, to the class numeral, for example, 2e. The letter e shows that the main hazard is the risk of erosion unless close-growing plant cover is maintained; w shows that water in or on the soil interferes with plant growth or cultivation (in some soils the wetness can be partly corrected by artificial drainage); s shows that the soil is limited mainly because it is shallow, droughty, or stony; and c, used in only some parts of the United States, shows that the chief limitation is climate that is very cold or very dry.

In class 1 there are no subclasses because the soils of this class have few limitations. Class 5 contains only the subclasses indicated by w, s, or c because the soils in class 5 are subject to little or no erosion. They have other limitations that restrict their use to pasture, rangeland, forestland, wildlife habitat, or recreation.

Capability units are soil groups within a subclass. The soils in a capability unit are enough alike to be suited to the same crops and pasture plants, to require similar management, and to have similar productivity. Capability units are generally designated by adding an Arabic numeral to the subclass symbol, for example, 2e-4 and 3e-6. These units are not given in all soil surveys.

The capability classification of map units in this survey area is given in the section "Detailed Soil Map Units" and in the Land Capability and Component Yields table.

Crop Yield Estimates

The average yields per acre that can be expected of the principal crops under a high level of management are shown in "Land Capability and Component Yields" table. In any given year, yields may be higher or lower than those indicated in the table because of variations in rainfall and other climatic factors. The land capability classification of map units in the survey area also is shown in the table.

The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations also are considered.

The management needed to obtain the indicated yields of the various crops depends on the kind of soil and the crop. Management can include drainage, erosion control, and protection from flooding; the proper planting and seeding rates; suitable high-yielding crop varieties; appropriate and timely tillage; control of weeds, plant diseases, and harmful insects; favorable soil reaction and optimum levels of nitrogen, phosphorus, potassium, and trace elements for each crop; effective use of crop residue, animal waste manure, and green manure crops; and harvesting that ensures the smallest possible loss.

For yields of irrigated crops, it is assumed that the irrigation system is adapted to the soils and to the crops grown, that good-quality irrigation water is uniformly applied as needed, and that tillage is kept to a minimum.

The estimated yields reflect the productive capacity of each soil for each of the principal crops. Yields are likely to increase as new production technology is developed. The productivity of a given soil compared with that of other soils, however, is not likely to change.

Crops other than those shown in this table, are grown in the survey area, but estimated yields are not listed because the acreage of such crops is small. The local office of the Natural Resources Conservation Service (NRCS) or the Cooperative Extension Service (CES) can provide information about the management and productivity of the soils for those crops.

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued
Kearney County, Nebraska

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

| Map symbol and soil name | Land Capability | | Corn | | Grain sorghum | | Soybeans | | Winter wheat | |
|--------------------------|-----------------|-----|-------|--------|---------------|--------|----------|-----|--------------|-----|
| | N | I | N | I | N | I | N | I | N | I |
| | | | Bu | | Bu | | | | Bu | |
| Ad: ALDA----- | 3w | 3w | 78.00 | 150.00 | 85.00 | 110.00 | --- | --- | 39.00 | --- |
| Bo: BOEL----- | 3w | 3w | 46.00 | 132.00 | 52.00 | 90.00 | --- | --- | 30.00 | --- |
| Bu: BUTLER----- | 2w | 2w | 59.00 | 162.00 | 81.00 | 110.00 | --- | --- | 44.00 | --- |
| CaC: COLY----- | 3e | 3e | 35.00 | 132.00 | 39.00 | 95.00 | --- | --- | 35.00 | --- |
| CaD: COLY----- | 4e | 4e | 33.00 | 108.00 | 35.00 | 90.00 | --- | --- | 33.00 | --- |
| CaF: COLY----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| CkB: COLY----- | 2e | 2e | 51.00 | 156.00 | 65.00 | 115.00 | --- | --- | 49.00 | --- |
| KENESAW----- | 2e | 2e | 51.00 | 156.00 | 65.00 | 115.00 | --- | --- | 49.00 | --- |
| CoD2: COLY----- | 4e | 4e | 23.00 | --- | 33.00 | 80.00 | --- | --- | 31.00 | --- |
| ULY----- | 4e | 4e | 23.00 | --- | 33.00 | 80.00 | --- | --- | 31.00 | --- |
| CoF2: COLY----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| ULY----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| De: DETROIT----- | 1 | 1 | 68.00 | 180.00 | 85.00 | 125.00 | --- | --- | 59.00 | --- |
| Fm: FILLMORE----- | 3w | 4w | 52.00 | 120.00 | 65.00 | 110.00 | --- | --- | 29.00 | --- |
| Gb: GIBBON----- | 2w | 2w | 91.00 | 162.00 | 91.00 | 115.00 | --- | --- | 42.00 | --- |
| Gc: GIBBON----- | 4s | 3s | 46.00 | 108.00 | 52.00 | 90.00 | --- | --- | 26.00 | --- |
| Go: GOTHENBURG----- | 7s | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| GP: PITS----- | 8s | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| HeB: HERSH----- | 3e | 2e | 46.00 | 144.00 | 48.00 | 95.00 | --- | --- | 33.00 | --- |
| HeC: HERSH----- | 3e | 3e | 40.00 | 144.00 | 43.00 | 85.00 | --- | --- | 30.00 | --- |
| HeD: HERSH----- | 4e | 4e | 34.00 | 108.00 | 26.00 | 75.00 | --- | --- | 23.00 | --- |
| Hf: HOBBS----- | 2w | 2w | 88.00 | 168.00 | 94.00 | 120.00 | --- | --- | 52.00 | --- |
| HgB: HOBBS----- | 6w | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Ho: HOLDREGE----- | 2c | 1 | 65.00 | 180.00 | 72.00 | 125.00 | --- | --- | 59.00 | --- |
| HoB: HOLDREGE----- | 2e | 2e | 55.00 | 168.00 | 65.00 | 120.00 | --- | --- | 55.00 | --- |
| HoC: HOLDREGE----- | 3e | 3e | 42.00 | 144.00 | 59.00 | 100.00 | --- | --- | 46.00 | --- |
| HoC2: HOLDREGE----- | 3e | 3e | 35.00 | 132.00 | 46.00 | 95.00 | --- | --- | 36.00 | --- |
| Hp: HOLDREGE----- | 2c | 1 | 72.00 | 180.00 | 72.00 | 125.00 | --- | --- | 59.00 | --- |

LAND CAPABILITY AND YIELDS PER ACRE OF CROPS--Continued
Kearney County, Nebraska

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)

| Map symbol and soil name | Land Capability | | Corn | | Grain sorghum | | Soybeans | | Winter wheat | |
|-------------------------------------|-----------------|-----|-------|--------|---------------|--------|----------|-----|--------------|-----|
| | N | I | N | I | N | I | N | I | N | I |
| | | | Bu | | Bu | | | | Bu | |
| HpB: HOLDREGE----- | 2e | 2e | 55.00 | 168.00 | 65.00 | 120.00 | --- | --- | 55.00 | --- |
| Hr: HORD----- | 2c | 1 | 72.00 | 180.00 | 85.00 | 125.00 | --- | --- | 56.00 | --- |
| InB: INAVALE----- | 4e | 3e | 33.00 | 120.00 | 39.00 | 80.00 | --- | --- | 25.00 | --- |
| Ke: KENESAW----- | 2c | 1 | 65.00 | 162.00 | 72.00 | 120.00 | --- | --- | 59.00 | --- |
| KgB: KENESAW----- | 2e | 2e | 59.00 | 168.00 | 65.00 | 120.00 | --- | --- | 48.00 | --- |
| COLY----- | 2e | 2e | 59.00 | 168.00 | 65.00 | 120.00 | --- | --- | 48.00 | --- |
| Lf: LEX----- | 3w | 3w | 91.00 | 156.00 | 100.00 | 115.00 | --- | --- | 46.00 | --- |
| Lg: LEX----- | 4s | 3s | 39.00 | 108.00 | 52.00 | 85.00 | --- | --- | 26.00 | --- |
| LoB: LIBORY----- | 3e | 3e | 59.00 | 144.00 | 59.00 | 115.00 | --- | --- | 33.00 | --- |
| M-w: MISCELLANEOUS WATER----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Ma: MASSIE----- | 8w | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Pm: PLATTE----- | 4w | 4w | 52.00 | 102.00 | 59.00 | 85.00 | --- | --- | --- | --- |
| Ru: RUSCO----- | 2w | 2w | 59.00 | 162.00 | 78.00 | 110.00 | --- | --- | 39.00 | --- |
| Sc: SCOTT----- | 4w | --- | --- | --- | 39.00 | --- | --- | --- | 20.00 | --- |
| SmB: SIMEON----- | 6s | 4s | --- | 108.00 | --- | 90.00 | --- | --- | --- | --- |
| To: TRYON----- | 5w | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| UcF: ULY----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| COLY----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| UsF: USTORTHENTS----- | 7s | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| VaB: VALENTINE----- | 4e | 4e | 33.00 | 132.00 | 39.00 | 95.00 | --- | --- | 25.00 | --- |
| VaD: VALENTINE----- | 6e | 4e | --- | 120.00 | --- | --- | --- | --- | --- | --- |
| VaF: VALENTINE----- | 6e | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| VbD: VALENTINE----- | 6e | 4e | --- | 132.00 | --- | --- | --- | --- | --- | --- |
| ELS----- | 4w | 4w | --- | 132.00 | --- | --- | --- | --- | --- | --- |
| W: WATER----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Wa: WANN----- | 2w | 2w | 91.00 | 168.00 | 91.00 | 110.00 | --- | --- | 39.00 | --- |

Prime farmland is one of several kinds of important farmland defined by the U.S. Department of Agriculture. It is of major importance in meeting the Nation's short- and long-range needs for food and fiber. Because the supply of high-quality farmland is limited, the U.S. Department of Agriculture recognizes that responsible levels of government, as well as individuals, should encourage and facilitate the wise use of our Nation's prime farmland.

Prime farmland, as defined by the U.S. Department of Agriculture, is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops and is available for these uses. It could be cultivated land, pastureland, forestland, or other land, but it is not urban or built-up land or water areas. The soil qualities, growing season, and moisture supply are those needed for the soil to economically produce sustained high yields of crops when proper management, including water management, and acceptable farming methods are applied. In general, prime farmland has an adequate and dependable supply of moisture from precipitation or irrigation, a favorable temperature and growing season, acceptable acidity or alkalinity, an acceptable salt and sodium content, and few or no rocks. It is permeable to water and air. It is not excessively erodible or saturated with water for long periods, and it either is not frequently flooded during the growing season or is protected from flooding. Slope ranges mainly from 0 to 6 percent. More detailed information about the criteria for prime farmland is available at the local office of the Natural Resources Conservation Service.

A recent trend in land use in some parts of the survey area has been the loss of some prime farmland to industrial and urban uses. The loss of prime farmland to other uses puts pressure on marginal lands, which generally are more erodible, droughty, and less productive and cannot be easily cultivated.

The map units in the survey area that are considered prime farmland are listed in the following table. This list does not constitute a recommendation for a particular land use. On some soils included in the list, measures that overcome a hazard or limitation, such as flooding, wetness, and droughtiness, are needed. Onsite evaluation is needed to determine whether or not the hazard or limitation has been overcome by corrective measures. The extent of each listed map unit is shown in the "Acres and Proportionate Extent of Soils" table. The location is shown on the detailed soil maps. The soil qualities that affect use and management are described in other tables in this document."

| Map symbol | Mapunit name | Farmland Classification |
|------------|--|------------------------------|
| CkB | Coly-kenesaw silt loams, 0 to 3 percent slopes | All areas are prime farmland |
| De | Detroit silt loam, 0 to 1 percent slopes | All areas are prime farmland |
| HeB | Hersh fine sandy loam, 0 to 3 percent slopes | All areas are prime farmland |
| HeC | Hersh fine sandy loam, 3 to 6 percent slopes | All areas are prime farmland |
| Hf | Hobbs silt loam, 0 to 2 percent slopes | All areas are prime farmland |
| Ho | Holdrege silt loam, 0 to 1 percent slopes | All areas are prime farmland |
| HoB | Holdrege silt loam, 1 to 3 percent slopes | All areas are prime farmland |
| HoC | Holdrege silt loam, 3 to 6 percent slopes | All areas are prime farmland |
| HoC2 | Holdrege silt loam, 3 to 6 percent slopes, eroded | All areas are prime farmland |
| Hp | Holdrege silt loam, overblown, 0 to 1 percent slopes | All areas are prime farmland |
| HpB | Holdrege silt loam, overblown, 1 to 3 percent slopes | All areas are prime farmland |
| Hr | Hord silt loam, 0 to 1 percent slopes | All areas are prime farmland |
| Ke | Kenesaw silt loam, 0 to 1 percent slopes | All areas are prime farmland |
| KgB | Kenesaw-coly silt loams, 1 to 3 percent slopes | All areas are prime farmland |
| Ru | Rusco silt loam, 0 to 1 percent slopes | All areas are prime farmland |
| Ad | Alda loam, 0 to 1 percent slopes | Prime farmland if drained |
| Bu | Butler silt loam, 0 to 1 percent slopes | Prime farmland if drained |
| Gb | Gibbon loam, 0 to 1 percent slopes | Prime farmland if drained |
| Lf | Lex loam, 0 to 1 percent slopes | Prime farmland if drained |
| Wa | Wann fine sandy loam, 0 to 1 percent slopes | Prime farmland if drained |

The "Soil Rating for Plant Growth, modified 1998" (SRPG) is a relative rating of the capacity of a soil to produce a specific plant under a defined management system. The index is determined from yield data on a few benchmark soils and is used to calculate yields, the net returns from crops, land assessment values, and taxes and to perform risk analysis when land management decisions are made. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

| Map symbol | Soil name | Crop Index |
|------------|--|------------|
| Ad | Alda Loam, 0 To 1 Percent Slopes----- | 35 |
| Bo | Boel Fine Sandy Loam, 0 To 1 Percent Slopes----- | 30 |
| Bu | Butler Silt Loam, 0 To 1 Percent Slopes----- | 65 |
| CaC | Coly Silt Loam, 3 To 6 Percent Slopes----- | 53 |
| CaD | Coly Silt Loam, 6 To 11 Percent Slopes----- | 49 |
| CaF | Coly Silt Loam, 11 To 20 Percent Slopes----- | 41 |
| CkB | Coly-Kenesaw Silt Loams, 0 To 3 Percent Slopes----- | 61 |
| CoD2 | Coly-Uly Silt Loams, 6 To 11 Percent Slopes, Eroded----- | 54 |
| CoF2 | Coly-Uly Silt Loams, 11 To 20 Percent Slopes, Eroded----- | 45 |
| De | Detroit Silt Loam, 0 To 1 Percent Slopes----- | 69 |
| Fm | Fillmore Silt Loam, 0 To 1 Percent Slopes----- | 46 |
| GP | Gravel Pit----- | 18 |
| Gb | Gibbon Loam, 0 To 1 Percent Slopes----- | 51 |
| Gc | Gibbon Loam, Saline, 0 To 1 Percent Slopes----- | 47 |
| Go | Gothenburg Loamy Sand, 0 To 2 Percent Slopes----- | 14 |
| HeB | Hersh Fine Sandy Loam, 0 To 3 Percent Slopes----- | 61 |
| HeC | Hersh Fine Sandy Loam, 3 To 6 Percent Slopes----- | 59 |
| HeD | Hersh Fine Sandy Loam, 6 To 11 Percent Slopes----- | 55 |
| Hf | Hobbs Silt Loam, 0 To 2 Percent Slopes----- | 66 |
| HgB | Hobbs Silt Loam, Channeled----- | 52 |
| Ho | Holdrege Silt Loam, 0 To 1 Percent Slopes----- | 74 |
| HoB | Holdrege Silt Loam, 1 To 3 Percent Slopes----- | 73 |
| HoC | Holdrege Silt Loam, 3 To 6 Percent Slopes----- | 70 |
| HoC2 | Holdrege Silt Loam, 3 To 6 Percent Slopes, Eroded----- | 70 |
| Hp | Holdrege Silt Loam, Overblown, 0 To 1 Percent Slopes----- | 74 |
| HpB | Holdrege Silt Loam, Overblown, 1 To 3 Percent Slopes----- | 73 |
| Hr | Hord Silt Loam, 0 To 1 Percent Slopes----- | 75 |
| InB | Inavale Loamy Fine Sand, 0 To 3 Percent Slopes----- | 33 |
| Ke | Kenesaw Silt Loam, 0 To 1 Percent Slopes----- | 71 |
| KgB | Kenesaw-Coly Silt Loams, 1 To 3 Percent Slopes----- | 63 |
| Lf | Lex Loam, 0 To 1 Percent Slopes----- | 37 |
| Lg | Lex Loam, Saline, 0 To 1 Percent Slopes----- | 33 |
| LoB | Libory Loamy Fine Sand, 0 To 3 Percent Slopes----- | 55 |
| M-W | Miscellaneous Water (sewage Lagoons)----- | 0 |
| Ma | Massie Silty Clay Loam, 0 To 1 Percent Slopes----- | 6 |
| Pm | Platte Loam, 0 To 1 Percent Slopes----- | 30 |
| Ru | Rusco Silt Loam, 0 To 1 Percent Slopes----- | 31 |
| Sc | Scott Silt Loam, 0 To 1 Percent Slopes----- | 19 |
| SmB | Simeon Sandy Loam, 0 To 3 Percent Slopes----- | 30 |
| To | Tryon Loamy Fine Sand, 0 To 1 Percent Slopes----- | 26 |
| UcF | Uly-Coly Silt Loams, 11 To 20 Percent Slopes----- | 47 |
| UsF | Ustorthents, Steep----- | 23 |
| VaB | Valentine Loamy Fine Sand, 0 To 3 Percent Slopes----- | 27 |
| VaD | Valentine Loamy Fine Sand, 3 To 9 Percent Slopes----- | 25 |
| VaF | Valentine Loamy Fine Sand, Rolling----- | 22 |
| VbD | Valentine-Els Loamy Fine Sands, 0 To 9 Percent Slopes----- | 25 |
| W | Water----- | 0 |
| Wa | Wann Fine Sandy Loam, 0 To 1 Percent Slopes----- | 43 |

Kearney County, Nebraska: Published
 Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

| Map symbol and soil name | Percent | Irr Cap Class | Nonirr Cap Class | Prime Farmland | Hydro-logic Group | Range site name | Windbreak suitability group | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|---------|---------------|------------------|------------------------------|-------------------|-------------------------------|-----------------------------|-----------------|-----|---|------------------------|------------------------|
| | | | | | | | | K | Kf | T | | |
| Ad:ALDA----- | 100 | 3w- | 3w | Prime farmland if drained | C | Subirrigated - Veg. Zone 3 | | .28 | .28 | 4 | 5 | 56 |
| Bo:BOEL----- | 100 | 3w- | 3w | Not prime farmland | A | Subirrigated - Veg. Zone 3 | | .20 | .20 | 3 | 3 | 86 |
| Bu:BUTLER----- | 100 | 2w- | 2w | Prime farmland if drained | D | Clayey - Veg. Zone 3 | | .37 | .37 | 3 | 6 | 48 |
| CaC:COLY----- | 100 | 3e- | 3e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CaD:COLY----- | 100 | 4e- | 4e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CaF:COLY----- | 100 | N/A | 6e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CkB:COLY----- | 60 | 2e- | 2e | All areas are prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CkB:KENESAW---- | 40 | 2e- | 2e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| CoD2:COLY----- | 60 | 4e- | 4e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CoD2:ULY----- | 40 | 4e- | 4e | Not prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| CoF2:COLY----- | 60 | N/A | 6e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| CoF2:ULY----- | 40 | N/A | 6e | Not prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| De:DETROIT----- | 100 | 1- | 1 | All areas are prime farmland | C | Silty Lowland - Veg. Zone 3 | | .37 | .37 | 5 | 6 | 48 |
| Fm:FILLMORE---- | 100 | 4w- | 3w | Not prime farmland | D | Clayey Overflow - Veg. Zone 3 | | .37 | .37 | 3 | 6 | 48 |
| GP:PITS----- | 100 | N/A | 8s | Not prime farmland | A | Unspecified | | .10 | .17 | 2 | 8 | 0 |
| Gb:GIBBON----- | 100 | 2w- | 2w | Prime farmland if drained | B | Subirrigated - Veg. Zone 3 | | .28 | .28 | 5 | 4L | 86 |
| Gc:GIBBON----- | 100 | 3s- | 4s | Not prime farmland | B | Subirrigated - Veg. Zone 3 | | .28 | .28 | 5 | 4L | 86 |
| Go:GOTHENBURG--- | 100 | N/A | 7s | Not prime farmland | D | Unspecified | | .17 | .17 | 5 | 8 | 0 |

Kearney County, Nebraska: Published
 Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

| Map symbol and soil name | Percent | Irr Cap Class | Nonirr Cap Class | Prime Farmland | Hydro-logic Group | Range site name | Windbreak suitability group | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|---------|---------------|------------------|------------------------------|-------------------|------------------------------|-----------------------------|-----------------|-----|---|------------------------|------------------------|
| | | | | | | | | K | Kf | T | | |
| HeB:HERSH----- | 100 | 2e- | 3e | All areas are prime farmland | B | Sandy - Veg. Zone 3 | | .24 | .24 | 5 | 3 | 86 |
| HeC:HERSH----- | 100 | 3e- | 3e | All areas are prime farmland | B | Sandy - Veg. Zone 3 | | .24 | .24 | 5 | 3 | 86 |
| HeD:HERSH----- | 100 | 4e- | 4e | Not prime farmland | B | Sandy - Veg. Zone 3 | | .24 | .24 | 5 | 3 | 86 |
| Hf:HOBBS----- | 100 | 2w- | 2w | All areas are prime farmland | B | Silty Overflow - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| HgB:HOBBS----- | 100 | N/A | 6w | Not prime farmland | B | Silty Overflow - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| Ho:HOLDREGE---- | 100 | 1- | 2c | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| HoB:HOLDREGE---- | 100 | 2e- | 2e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| HoC:HOLDREGE---- | 100 | 3e- | 3e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| HoC2:HOLDREGE--- | 100 | 3e- | 3e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| Hp:HOLDREGE---- | 100 | 1- | 2c | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| HpB:HOLDREGE---- | 100 | 2e- | 2e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| Hr:HORD----- | 100 | 1- | 2c | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| InB:INAVALE---- | 100 | 3e- | 4e | Not prime farmland | A | Sandy Lowland - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| Ke:KENESAW----- | 100 | 1- | 2c | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| KgB:KENESAW----- | 55 | 2e- | 2e | All areas are prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| KgB:COLY----- | 45 | 2e- | 2e | All areas are prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| Lf:LEX----- | 100 | 3w- | 3w | Prime farmland if drained | B | Subirrigated - Veg. Zone 3 | | .28 | .28 | 4 | 4L | 86 |

Kearney County, Nebraska: Published
 Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

| Map symbol and soil name | Percent | Irr Cap Class | Nonirr Cap Class | Prime Farmland | Hydro-logic Group | Range site name | Windbreak suitability group | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|-------------------------------------|---------|---------------|------------------|------------------------------|-------------------|-----------------------------------|-----------------------------|-----------------|-----|---|------------------------|------------------------|
| | | | | | | | | K | Kf | T | | |
| Lg:LEX----- | 100 | 3s- | 4s | Not prime farmland | B | Saline Subirrigated - Veg. Zone 3 | | .28 | .28 | 4 | 4L | 86 |
| LoB:LIBORY----- | 100 | 3e- | 3e | Not prime farmland | A | Sandy Lowland - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| M- W:MISCELLANEOUS WATER----- | 100 | N/A | N/A | Not prime farmland | | Unspecified | | --- | --- | - | --- | --- |
| Ma:MASSIE----- | 100 | N/A | 8w | Not prime farmland | D | Unspecified | | .37 | .37 | 3 | 7 | 38 |
| Pm:PLATTE----- | 100 | 4w- | 4w | Not prime farmland | B | Subirrigated - Veg. Zone 3 | | .28 | .28 | 3 | 4L | 86 |
| Ru:RUSCO----- | 100 | 2w- | 2w | All areas are prime farmland | C | Silty Lowland - Veg. Zone 3 | | .32 | .32 | 5 | 5 | 56 |
| Sc:SCOTT----- | 100 | N/A | 4w | Not prime farmland | D | Clayey Overflow - Veg. Zone 3 | | .37 | .37 | 3 | 6 | 48 |
| SmB:SIMEON----- | 100 | 4s- | 6s | Not prime farmland | A | Shallow To Gravel - Veg. Zone 3 | | .24 | .24 | 5 | 3 | 86 |
| To:TRYON----- | 100 | N/A | 5w | Not prime farmland | D | Wet Subirrigated - Veg. Zone 3 | | .17 | .17 | 5 | 8 | 0 |
| UcF:ULY----- | 60 | N/A | 6e | Not prime farmland | B | Silty - Veg. Zone 3 | | .32 | .32 | 5 | 6 | 48 |
| UcF:COLY----- | 40 | N/A | 6e | Not prime farmland | B | Limy Upland - Veg. Zone 3 | | .43 | .43 | 5 | 4L | 86 |
| UsF:USTORTHENTS- | 100 | N/A | 7s | Not prime farmland | B | Unspecified | | .43 | .43 | 5 | 4L | 86 |
| VaB:VALENTINE--- | 100 | 4e- | 4e | Not prime farmland | A | Sandy - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| VaD:VALENTINE--- | 100 | 4e- | 6e | Not prime farmland | A | Sands - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| VaF:VALENTINE--- | 100 | N/A | 6e | Not prime farmland | A | Sands - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| VbD:VALENTINE--- | 65 | 4e- | 6e | Not prime farmland | A | Sands - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| VbD:ELS----- | 35 | 4w- | 4w | Not prime farmland | A | Subirrigated - Veg. Zone 3 | | .17 | .17 | 5 | 2 | 134 |
| W:WATER----- | 100 | N/A | N/A | Not prime farmland | | Unspecified | | --- | --- | - | --- | 0 |
| Wa:WANN----- | 100 | 2w- | 2w | Prime farmland if drained | B | Subirrigated - Veg. Zone 3 | | .20 | .20 | 5 | 3 | 86 |

Field Office Thunderbook: Soils Properties for Conservation Planning

(Entries under "Erosion factors--T" apply to the entire profile. Entries under "K", "Kf", "Wind Erodibility Group" and "Wind Erodibility Index" apply only to the surface layer)

| Map symbol and soil name | Percent | Irr Cap Class | Nonirr Cap Class | Prime Farmland | Hydrologic Group | Range site name | Windbreak suitability group | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|---------|---------------|------------------|----------------|------------------|-----------------|-----------------------------|-----------------|----|---|------------------------|------------------------|
| | | | | | | | | K | Kf | T | | |
| | | | | | | | | | | | | |

RANGELAND PRODUCTIVITY
Kearney County, Nebraska

Use and Explanation of Rangeland, Grazed Forest Land, Native Pastureland Interpretations

Information in this subsection can be used to plan the use and management of soils for rangeland, grazed forest land, and native pasture. Different kinds of soils vary in their capacity to produce native grasses and other plants suitable for grazing. Information in this subsection provides groupings of similar soils and estimates of potential forage production, which can be used to determine livestock stocking rates.

Rangeland. Range is land on which the native vegetation (climax or natural potential plant community) is predominantly grasses, grasslike plants, forbs, and shrubs suitable for grazing and browsing. Range includes natural grasslands, savannas, many wetlands, some deserts, tundra, and certain shrub and forb communities. Rangeland receives no regular or frequent cultural treatment. The composition and production of the plant community are determined by soil, climate, topography, overstory canopy, and grazing management.

Grazed Forest Land. Includes land on which the understory includes, as an integral part of the forest plant community, plants that can be grazed without significantly impairing other forest values.

Native Pasture. Includes land on which the native vegetation (climax or natural potential plant community) is forest but which is used and managed primarily for production of native plants for forage. Native pasture includes cut-over forest land and forest land cleared and now managed for native or naturalized forage plants.

Rangeland

In areas that have similar climate and topography, differences in the kind and amount of vegetation produced on rangeland are closely related to the kind of soil. Effective management based on the relationship between the soils and vegetation and water.

The Rangeland, Grazed Forest land, Native Pastureland Interpretations shows, for each soil that supports rangeland vegetation, the ecological site and the potential annual production of vegetation in favorable, normal, unfavorable years. An explanation of the column headings in this table follows.

An ecological site is the product of all the environmental factors responsible for its development. It has characteristic soils that have developed over time throughout the soil development process; a characteristic hydrology, particularly infiltration and runoff, that has developed over time; and a characteristic plant community (kind and amount of vegetation). The hydrology of a site is influenced by development of the soil and plant community. The vegetation, soils, and hydrology are all interrelated. Each is influenced by the others and influences the development of the others. The plant community on an ecological site is typified by an association of species that differs from that of other ecological sites in the kind and/or proportion of species or in total production. Descriptions of ecological sites are provided in the Field Office Technical Guide, which is available in local offices of the Natural Resources Conservation Service.

Total dry-weight production is the amount of vegetation that can be expected to grow annually on well managed rangeland that is supporting the potential natural plant community. It includes all vegetation, whether or not it is palatable to grazing animals. It includes the current year's growth of leaves, twigs, and fruits of woody plants. It does not include the increase in stem diameter of trees and shrubs. It is expressed in pounds per acre of air-dry vegetation for favorable, average, and unfavorable years. In a favorable year, the amount and distribution of precipitation and the temperatures make growing conditions substantially better than average. In a normal year, growing conditions are about average. In an unfavorable year, growing conditions are well below average, generally because of low available soil moisture. Yields are adjusted to a common percent of air-dry moisture content.

Range management requires a knowledge of the kinds of soil and of the potential natural plant community. It also requires an evaluation of the present range similarity index and rangeland trend. Range similarity index is determined by comparing the present plant community with the potential natural plant community on a particular rangeland ecological site. The more closely the existing community resembles the potential community, the higher the range similarity index. Rangeland trend is defined as the direction of change in an existing plant community relative to the potential natural plant community. Further information about the range similarity index and rangeland trend is available in chapter 4 of the National Range and Pasture Handbook, which is available in local offices of the Natural Resources Conservation Service. The objective in range management is to control grazing so that the plants growing on a site are about the same in kind and amount as the potential natural plant community for that site. Such management generally results in the optimum production of vegetation, control of undesirable brush species, conservation of water, and control of erosion. Sometimes, however, an area with a range similarity index somewhat below the potential meets grazing needs, provides wildlife habitat, and protects soil and water resources.

RANGELAND PRODUCTIVITY--Continued
Kearney County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

| Map symbol and soil name | Ecological site | Total dry-weight production | | |
|-----------------------------|--------------------------------------|-----------------------------|-----------------|---------------------|
| | | Favorable year | Average year | Unfavorable year |
| | | Lb/acre | Lb/acre | Lb/acre |
| Ad: | | | | |
| Alda----- | Subirrigated - Veg. Zone 3 | 5,200 | 4,900 | 4,600 |
| Bo: | | | | |
| Boel----- | Subirrigated - Veg. Zone 3 | 5,200 | 4,900 | 4,600 |
| Bu: | | | | |
| Butler----- | Clayey - Veg. Zone 3 | 3,800 | 3,400 | 3,000 |
| CaC: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| CaD: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| CaF: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| CkB: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| Kenesaw----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| CoD2: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| Uly----- | Silty - Veg. Zone 3 | 3,700 | 3,200 | 2,700 |
| CoF2: | | | | |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| Uly----- | Silty - Veg. Zone 3 | 3,700 | 3,200 | 2,700 |
| De: | | | | |
| Detroit----- | Silty Lowland - Veg. Zone 3 | 6,000 | 4,500 | 3,000 |
| Fm: | | | | |
| Fillmore----- | Clayey Overflow - Veg. Zone 3 | 3,200 | 2,700 | 2,200 |
| Gb: | | | | |
| Gibbon----- | Subirrigated - Veg. Zone 3 | 5,500 | 5,300 | 5,000 |
| Gc: | | | | |
| Gibbon----- | Subirrigated - Veg. Zone 3 | 4,000 | 3,800 | 3,500 |
| Go: | | | | |
| Gothenburg----- | --- | --- | --- | --- |
| GP: | | | | |
| Pits----- | --- | --- | --- | --- |
| HeB: | | | | |
| Hersh----- | Sandy - Veg. Zone 3 | 3,500 | 3,300 | 3,000 |
| HeC: | | | | |
| Hersh----- | Sandy - Veg. Zone 3 | 3,500 | 3,300 | 3,000 |
| HeD: | | | | |
| Hersh----- | Sandy - Veg. Zone 3 | 3,500 | 3,300 | 3,000 |
| Hf: | | | | |
| Hobbs----- | Silty Overflow - Veg. Zone 3 | 4,500 | 4,000 | 3,800 |
| HgB: | | | | |
| Hobbs----- | Silty Overflow - Veg. Zone 3 | 4,500 | 4,000 | 3,800 |
| Ho: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| HoB: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| HoC: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| HoC2: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| Hp: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| HpB: | | | | |
| Holdrege----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| Hr: | | | | |
| Hord----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| InB: | | | | |
| Inavale----- | Sandy Lowland - Veg. Zone 3 | 3,500 | 3,000 | 2,200 |
| Ke: | | | | |
| Kenesaw----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| KgB: | | | | |
| Kenesaw----- | Silty - Veg. Zone 3 | 4,000 | 3,600 | 3,300 |
| Coly----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| Lf: | | | | |
| Lex----- | Subirrigated - Veg. Zone 3 | 5,200 | 4,900 | 4,600 |
| Lg: | | | | |
| Lex----- | Saline Subirrigated - Veg. Zone 3 | 3,800 | 3,400 | 3,000 |
| LoB: | | | | |
| Libory----- | Sandy Lowland - Veg. Zone 3 | 4,300 | 3,500 | 2,700 |
| M-W: | | | | |
| Miscellaneous Water----- | --- | --- | --- | --- |
| Ma: | | | | |
| Massie----- | --- | --- | --- | --- |
| Pm: | | | | |
| Platte----- | Subirrigated - Veg. Zone 3 | 5,000 | 4,600 | 4,200 |
| Ru: | | | | |
| Rusco----- | Silty Lowland - Veg. Zone 3 | 3,500 | 3,000 | 2,500 |
| Sc: | | | | |
| Scott----- | Clayey Overflow - Veg. Zone 3 | 3,900 | 3,300 | 2,300 |
| SmB: | | | | |
| Simeon----- | Shallow To Gravel - Veg. Zone 3 | 1,800 | 1,600 | 1,100 |
| To: | | | | |

RANGELAND PRODUCTIVITY--Continued
Kearney County, Nebraska

(Only the soils that support rangeland vegetation suitable for grazing are rated.) Refer to range site description to determine the percentage allowable of grasses, forbs, and shrubs for the range ecological site.

| Map symbol and soil name | Ecological site | Total dry-weight production | | |
|----------------------------------|--------------------------------|-----------------------------|-----------------|---------------------|
| | | Favorable year | Average year | Unfavorable year |
| | | Lb/acre | Lb/acre | Lb/acre |
| Tryon----- UcF: | Wet Subirrigated - Veg. Zone 3 | 5,800 | 5,500 | 5,300 |
| Uly----- Coly----- UsF: | Silty - Veg. Zone 3 | 3,700 | 3,200 | 2,700 |
| Ustorthents----- | Limy Upland - Veg. Zone 3 | 3,300 | 3,000 | 2,700 |
| VaB: | --- | --- | --- | --- |
| Valentine----- VaD: | Sandy - Veg. Zone 3 | 3,300 | 3,000 | 2,600 |
| Valentine----- VaF: | Sands - Veg. Zone 3 | 3,000 | 2,600 | 2,200 |
| Valentine----- VbD: | Sands - Veg. Zone 3 | 3,000 | 2,600 | 2,200 |
| Valentine----- Els----- W: | Sands - Veg. Zone 3 | 3,000 | 2,600 | 2,200 |
| Water----- Wa: | Subirrigated - Veg. Zone 3 | 5,500 | 5,300 | 5,000 |
| Wann----- | --- | --- | --- | --- |
| | Subirrigated - Veg. Zone 3 | 5,500 | 5,300 | 5,000 |

BUILDING SITE DEVELOPMENT
Kearney County, Nebraska

Building Site Development

Soil properties influence the development of building sites, including the selection of the site, the design of the structure, construction, performance after construction, and maintenance. These tables show the degree and kind of soil limitations that affect dwellings with and without basements, small commercial buildings, local roads and streets, shallow excavations, and lawns and landscaping.

The ratings in the tables are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect building site development. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Dwellings are single-family houses of three stories or less. For dwellings without basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. For dwellings with basements, the foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of about 7 feet. The ratings for dwellings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility. Compressibility is inferred from the Unified classification. The properties that affect the ease and amount of excavation include depth to a water table, ponding, flooding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Small commercial buildings are structures that are less than three stories high and do not have basements. The foundation is assumed to consist of spread footings of reinforced concrete built on undisturbed soil at a depth of 2 feet or at the depth of maximum frost penetration, whichever is deeper. The ratings are based on the soil properties that affect the capacity of the soil to support a load without movement and on the properties that affect excavation and construction costs. The properties that affect the load-supporting capacity include depth to a water table, ponding, flooding, subsidence, linear extensibility (shrink-swell potential), and compressibility (which is inferred from the Unified classification). The properties that affect the ease and amount of excavation include flooding, depth to a water table, ponding, slope, depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, and the amount and size of rock fragments.

Local roads and streets have an all-weather surface and carry automobile and light truck traffic all year. They have a subgrade of cut or fill soil material; a base of gravel, crushed rock, or soil material stabilized by lime or cement; and a surface of flexible material (asphalt), rigid material (concrete), or gravel with a binder. The ratings are based on the soil properties that affect the ease of excavation and grading and the traffic-supporting capacity. The properties that affect the ease of excavation and grading are depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, depth to a water table, ponding, flooding, the amount of large stones, and slope. The properties that affect the traffic-supporting capacity are soil strength (as inferred from the AASHTO group index number), subsidence, linear extensibility (shrink-swell potential), the potential for frost action, depth to a water table, and ponding.

Shallow excavations are trenches or holes dug to a maximum depth of 5 or 6 feet for graves, utility lines, open ditches, or other purposes. The ratings are based on the soil properties that influence the ease of digging and the resistance to sloughing. Depth to bedrock or a cemented pan, hardness of bedrock or a cemented pan, the amount of large stones, and dense layers influence the ease of digging, filling, and compacting. Depth to the seasonal high water table, flooding, and ponding may restrict the period when excavations can be made. Slope influences the ease of using machinery. Soil texture, depth to the water table, and linear extensibility (shrink-swell potential) influence the resistance to sloughing.

Lawns and landscaping require soils on which turf and ornamental trees and shrubs can be established and maintained. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer.

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Dwellings without basements | | Dwellings with basements | | Small commercial buildings | |
|-----------------------------|--------------------------|--|----------------------|--|--------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 0.07 |
| Bo: Boel----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 0.07 |
| Bu: Butler----- | 100 | Very limited Shrink-swell Depth to saturated zone | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Shrink-swell Depth to saturated zone | 1.00 1.00 |
| CaC: Coly----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.12 |
| CaD: Coly----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| CkB: Coly----- | 60 | Not limited | | Not limited | | Not limited | |
| Kenesaw----- | 40 | Not limited | | Not limited | | Not limited | |
| CoD2: Coly----- | 60 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| Uly----- | 40 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| De: Detroit----- | 100 | Very limited Shrink-swell | 1.00 | Very limited Shrink-swell | 1.00 | Very limited Shrink-swell | 1.00 |
| Fm: Fillmore----- | 100 | Very limited Shrink-swell Depth to saturated zone | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Shrink-swell Depth to saturated zone | 1.00 1.00 |
| Gb: Gibbon----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 0.07 |
| Gc: Gibbon----- | 100 | Very limited Flooding Depth to saturated zone Shrink-swell | 1.00 0.07 0.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone Shrink-swell | 1.00 0.07 0.00 |
| Go: Gothenburg----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Not limited | | Not limited | | Not limited | |
| HeC: Hersh----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.12 |
| HeD: Hersh----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| Hf: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Dwellings without basements | | Dwellings with basements | | Small commercial buildings | |
|------------------------------|--------------------------|--|--------------------------|--|--------------|--|--------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| Ho: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Not limited | | Somewhat limited Shrink-swell | 0.50 |
| HoB: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Not limited | | Somewhat limited Shrink-swell | 0.50 |
| HoC: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Not limited | | Somewhat limited Shrink-swell Slope | 0.50 0.12 |
| HoC2: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Not limited | | Somewhat limited Shrink-swell Slope | 0.50 0.12 |
| Hp: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HpB: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| Hr: Hord----- | 100 | Somewhat limited Shrink-swell | 0.06 | Somewhat limited Shrink-swell | 0.06 | Somewhat limited Shrink-swell | 0.06 |
| InB: Inavale----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| Ke: Kenesaw----- | 100 | Not limited | | Not limited | | Not limited | |
| KgB: Kenesaw----- | 55 | Not limited | | Not limited | | Not limited | |
| Lf: Coly----- | 45 | Not limited | | Not limited | | Not limited | |
| Lf: Lex----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.39 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 0.39 |
| Lg: Lex----- | 100 | Very limited Flooding Shrink-swell Depth to saturated zone | 1.00 0.50 0.07 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Shrink-swell Depth to saturated zone | 1.00 0.50 0.07 |
| LoB: Libory----- | 100 | Somewhat limited Depth to saturated zone | 0.07 | Very limited Depth to saturated zone | 1.00 | Somewhat limited Depth to saturated zone | 0.07 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 |
| Pm: Platte----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.39 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 0.39 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone Shrink-swell | 1.00 0.50 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 0.50 |
| Sc: Scott----- | 100 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 | Very limited Depth to saturated zone Shrink-swell | 1.00 1.00 |
| SmB: Simeon----- | 100 | Not limited | | Not limited | | Not limited | |
| To: Tryon----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Dwellings without basements | | Dwellings with basements | | Small commercial buildings | |
|-----------------------------|--------------------------|--|-------|--|--------------|--|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| UcF: Uly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Coly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| UsF: Ustorhents----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Not limited | | Not limited | | Not limited | |
| VaD: Valentine----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.48 |
| VaF: Valentine----- | 100 | Somewhat limited Slope | 0.84 | Somewhat limited Slope | 0.84 | Very limited Slope | 1.00 |
| VbD: Valentine----- | 65 | Not limited | | Not limited | | Somewhat limited Slope | 0.48 |
| Els----- | 35 | Somewhat limited Depth to saturated zone | 0.07 | Very limited Depth to saturated zone | 1.00 | Somewhat limited Depth to saturated zone | 0.07 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding Depth to saturated zone | 1.00 1.00 | Very limited Flooding | 1.00 |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Local roads and streets | | Shallow excavations | | Lawns and landscaping | |
|--------------------------|-----------------|------------------------------------|-------|---|-------|---|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Frost action | 1.00 | Very limited Cutbanks cave | 1.00 | Somewhat limited Depth to saturated zone | 0.03 |
| | | Flooding | 0.40 | Depth to saturated zone | 1.00 | | |
| | | Depth to saturated zone | 0.03 | | | | |
| Bo: Boel----- | 100 | Somewhat limited Frost action | 0.50 | Very limited Cutbanks cave | 1.00 | Somewhat limited Depth to saturated zone | 0.03 |
| | | Flooding | 0.40 | Depth to saturated zone | 1.00 | | |
| | | Depth to saturated zone | 0.03 | | | | |
| Bu: Butler----- | 100 | Very limited Shrink-swell | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Frost action | 1.00 | Too clayey | 0.50 | | |
| | | Depth to saturated zone | 1.00 | Cutbanks cave | 0.10 | | |
| CaC: Coly----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| CaD: Coly----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Somewhat limited Slope | 0.04 |
| | | Slope | 0.04 | Slope | 0.04 | | |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Frost action | 0.50 | Cutbanks cave | 0.10 | | |
| CkB: Coly----- | 60 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| Kenesaw----- | 40 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| CoD2: Coly----- | 60 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Somewhat limited Slope | 0.04 |
| | | Slope | 0.04 | Slope | 0.04 | | |
| Uly----- | 40 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Somewhat limited Slope | 0.04 |
| | | Slope | 0.04 | Slope | 0.04 | | |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Frost action | 0.50 | Cutbanks cave | 0.10 | | |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Frost action | 0.50 | Cutbanks cave | 0.10 | | |
| De: Detroit----- | 100 | Very limited Shrink-swell | 1.00 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| Fm: Fillmore----- | 100 | Very limited Shrink-swell | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Frost action | 1.00 | Too clayey | 0.50 | | |
| | | Depth to saturated zone | 1.00 | Cutbanks cave | 0.10 | | |
| Gb: Gibbon----- | 100 | Very limited Frost action | 1.00 | Very limited Depth to saturated zone | 1.00 | Somewhat limited Depth to saturated zone | 0.03 |
| | | Flooding | 0.40 | Cutbanks cave | 0.10 | | |
| | | Depth to saturated zone | 0.03 | | | | |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Local roads and streets | | Shallow excavations | | Lawns and landscaping | |
|-----------------------------|--------------------------|---------------------------------------|-------|--|-------|---------------------------------------|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Gc: Gibbon----- | 100 | Very limited Frost action | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Salinity | 1.00 |
| | | Flooding | 0.40 | Cutbanks cave | 0.10 | Depth to saturated zone | 0.03 |
| | | Depth to saturated zone | 0.03 | | | | |
| | | Shrink-swell | 0.00 | | | | |
| Go: Gothenburg----- | 100 | Very limited Flooding | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Flooding | 1.00 |
| | | Depth to saturated zone | 1.00 | Cutbanks cave | 1.00 | Droughty | 1.00 |
| | | Frost action | 0.50 | Flooding | 0.80 | Depth to saturated zone | 1.00 |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| HeC: Hersh----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| HeD: Hersh----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Somewhat limited Slope | 0.04 |
| | | Slope | 0.04 | Slope | 0.04 | | |
| Hf: Hobbs----- | 100 | Very limited Flooding | 1.00 | Somewhat limited Flooding | 0.60 | Somewhat limited Flooding | 0.60 |
| | | Frost action | 0.50 | Cutbanks cave | 0.10 | | |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Somewhat limited Flooding | 0.80 | Very limited Flooding | 1.00 |
| | | Frost action | 0.50 | Cutbanks cave | 0.10 | | |
| Ho: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| | | Frost action | 0.50 | | | | |
| HoB: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| | | Frost action | 0.50 | | | | |
| HoC: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| | | Frost action | 0.50 | | | | |
| HoC2: Holdrege----- | 100 | Somewhat limited Shrink-swell | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| | | Frost action | 0.50 | | | | |
| Hp: Holdrege----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| HpB: Holdrege----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| Hr: Hord----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| | | Shrink-swell | 0.06 | | | | |
| InB: Inavale----- | 100 | Somewhat limited Flooding | 0.40 | Very limited Cutbanks cave | 1.00 | Somewhat limited Droughty | 0.19 |
| Ke: Kenesaw----- | 100 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| KgB: Kenesaw----- | 55 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |
| Coly----- | 45 | Somewhat limited Frost action | 0.50 | Somewhat limited Cutbanks cave | 0.10 | Not limited | |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Local roads and streets | | Shallow excavations | | Lawns and landscaping | |
|------------------------------|--------------------------|--|----------------------------------|--|------------------------------|--|--------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Lf: Lex----- | 100 | Very limited Frost action Flooding Depth to saturated zone | 1.00 0.40 0.19 | Very limited Cutbanks cave Depth to saturated zone | 1.00 1.00 | Somewhat limited Depth to saturated zone | 0.19 |
| Lg: Lex----- | 100 | Very limited Frost action Shrink-swell Flooding Depth to saturated zone | 1.00 0.50 0.40 0.03 | Very limited Cutbanks cave Depth to saturated zone | 1.00 1.00 | Very limited Salinity Depth to saturated zone | 1.00 0.03 |
| LoB: Libory----- | 100 | Somewhat limited Depth to saturated zone | 0.03 | Very limited Cutbanks cave Depth to saturated zone | 1.00 1.00 | Somewhat limited Depth to saturated zone | 0.03 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Depth to saturated zone Frost action Shrink-swell | 1.00 1.00 1.00 | Very limited Depth to saturated zone Too clayey Cutbanks cave | 1.00 0.12 0.10 | Very limited Depth to saturated zone | 1.00 |
| Pm: Platte----- | 100 | Very limited Flooding Frost action Depth to saturated zone | 1.00 0.50 0.19 | Very limited Cutbanks cave Depth to saturated zone Flooding Depth to dense layer | 1.00 1.00 0.60 0.50 | Somewhat limited Flooding Droughty Depth to saturated zone | 0.60 0.35 0.19 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone Frost action Shrink-swell | 1.00 1.00 0.50 | Very limited Depth to saturated zone Cutbanks cave | 1.00 0.10 | Very limited Depth to saturated zone | 1.00 |
| Sc: Scott----- | 100 | Very limited Shrink-swell Depth to saturated zone Frost action | 1.00 1.00 1.00 | Very limited Depth to saturated zone Too clayey Cutbanks cave | 1.00 0.28 0.10 | Very limited Depth to saturated zone | 1.00 |
| SmB: Simeon----- | 100 | Not limited | | Very limited Cutbanks cave | 1.00 | Somewhat limited Droughty | 0.11 |
| To: Tryon----- | 100 | Very limited Depth to saturated zone Frost action Flooding | 1.00 0.50 0.40 | Very limited Depth to saturated zone Cutbanks cave | 1.00 1.00 | Very limited Depth to saturated zone Droughty | 1.00 0.56 |
| UcF: Uly----- | 60 | Very limited Slope Frost action | 1.00 0.50 | Very limited Slope Cutbanks cave | 1.00 0.10 | Very limited Slope | 1.00 |
| Coly----- | 40 | Very limited Slope Frost action | 1.00 0.50 | Very limited Slope Cutbanks cave | 1.00 0.10 | Very limited Slope | 1.00 |
| UsF: Ustorthents----- | 100 | Very limited Slope Frost action | 1.00 0.50 | Very limited Slope Cutbanks cave | 1.00 0.10 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Not limited | | Very limited Cutbanks cave | 1.00 | Somewhat limited Droughty | 0.71 |

BUILDING SITE DEVELOPMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Local roads and streets | | Shallow excavations | | Lawns and landscaping | |
|-----------------------------|--------------------------|--|--------------|---|--------------|--|--------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| VaD: Valentine----- | 100 | Not limited | | Very limited Cutbanks cave | 1.00 | Somewhat limited Droughty | 0.71 |
| VaF: Valentine----- | 100 | Somewhat limited Slope | 0.84 | Very limited Cutbanks cave Slope | 1.00 0.84 | Somewhat limited Slope Droughty | 0.84 0.71 |
| VbD: Valentine----- | 65 | Not limited | | Very limited Cutbanks cave | 1.00 | Somewhat limited Droughty | 0.71 |
| Els----- | 35 | Somewhat limited Frost action Depth to saturated zone | 0.50 0.03 | Very limited Cutbanks cave Depth to saturated zone | 1.00 1.00 | Somewhat limited Droughty Somewhat limited Droughty Depth to saturated zone | 0.74 0.03 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Very limited Frost action Flooding | 1.00 0.40 | Very limited Cutbanks cave Depth to saturated zone | 1.00 1.00 | Not limited | |

CONSTRUCTION MATERIALS
Kearney County, Nebraska

Construction Materials

These tables give information about the soils as potential sources of gravel, sand, topsoil, reclamation material, and roadfill. Normal compaction, minor processing, and other standard construction practices are assumed.

The soils are rated good, fair, or poor as potential sources of topsoil, reclamation material, and roadfill. The features that limit the soils as sources of these materials are specified in the tables. The numerical ratings given after the specified features indicate the degree to which the features limit the soils as sources of topsoil, reclamation material, or roadfill. The lower the number, the greater the limitation.

The soils are rated as a probable or improbable source of sand and gravel. A rating of probable means that the source material is likely to be in or below the soil. The numerical ratings in these columns indicate the degree of probability. The number 0.00 indicates that the soil is an improbable source. A number between 0.00 and 1.00 indicates the degree to which the soil is a probable source of sand or gravel.

Sand and gravel are natural aggregates suitable for commercial use with a minimum of processing. They are used in many kinds of construction. Specifications for each use vary widely. In the first table, only the probability of finding material in suitable quantity is evaluated. The suitability of the material for specific purposes is not evaluated, nor are factors that affect excavation of the material. The properties used to evaluate the soil as a source of sand or gravel are gradation of grain sizes (as indicated by the Unified classification of the soil), the thickness of suitable material, and the content of rock fragments. If the lowest layer of the soil contains sand or gravel, the soil is rated as a probable source regardless of thickness. The assumption is that the sand or gravel layer below the depth of observation exceeds the minimum thickness.

Topsoil is used to cover an area so that vegetation can be established and maintained. The upper 40 inches of a soil is evaluated for use as topsoil. Also evaluated is the reclamation potential of the borrow area. The ratings are based on the soil properties that affect plant growth; the ease of excavating, loading, and spreading the material; and reclamation of the borrow area. Toxic substances, soil reaction, and the properties that are inferred from soil texture, such as available water capacity and fertility, affect plant growth. The ease of excavating, loading, and spreading is affected by rock fragments, slope, depth to a water table, soil texture, and thickness of suitable material. Reclamation of the borrow area is affected by slope, depth to a water table, rock fragments, depth to bedrock or a cemented pan, and toxic material.

The surface layer of most soils is generally preferred for topsoil because of its organic matter content. Organic matter greatly increases the absorption and retention of moisture and nutrients for plant growth.

Reclamation material is used in areas that have been drastically disturbed by surface mining or similar activities. When these areas are reclaimed, layers of soil material or unconsolidated geological material, or both, are replaced in a vertical sequence. The reconstructed soil favors plant growth. The ratings in the table do not apply to quarries and other mined areas that require an offsite source of reconstruction material. The ratings are based on the soil properties that affect erosion and stability of the surface and the productive potential of the reconstructed soil. These properties include the content of sodium, salts, and calcium carbonate; reaction; available water capacity; erodibility; texture; content of rock fragments; and content of organic matter and other features that affect fertility.

Roadfill is soil material that is excavated in one place and used in road embankments in another place. In this table, the soils are rated as a source of roadfill for low embankments, generally less than 6 feet high and less exacting in design than higher embankments.

The ratings are for the whole soil, from the surface to a depth of about 5 feet. It is assumed that soil layers will be mixed when the soil material is excavated and spread.

The ratings are based on the amount of suitable material and on soil properties that affect the ease of excavation and the performance of the material after it is in place. The thickness of the suitable material is a major consideration. The ease of excavation is affected by large stones, depth to a water table, and slope. How well the soil performs in place after it has been compacted and drained is determined by its strength (as inferred from the AASHTO classification of the soil) and linear extensibility (shrink-swell potential).

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of gravel | | Potential source of sand | |
|--------------------------|------------------|--|--------------|--|--------------|
| | | Rating class | Value | Rating class | Value |
| Ad: Alda----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.09 0.98 |
| Bo: Boel----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.91 |
| Bu: Butler----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CaC: Coly----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CaD: Coly----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CaF: Coly----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CkB: Coly----- | 60 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Kenesaw----- | 40 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CoD2: Coly----- | 60 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Uly----- | 40 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| CoF2: Coly----- | 60 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Uly----- | 40 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| De: Detroit----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Fm: Fillmore----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Gb: Gibbon----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Gc: Gibbon----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of gravel | | Potential source of sand | |
|--------------------------|------------------|--|--------------|--|--------------|
| | | Rating class | Value | Rating class | Value |
| Go: Gothenburg----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Bottom layer | 0.16 |
| GP: Pits----- | 100 | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Bottom layer Thickest layer | 0.09 0.09 |
| HeC: Hersh----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Bottom layer Thickest layer | 0.09 0.09 |
| HeD: Hersh----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Bottom layer Thickest layer | 0.09 0.09 |
| Hf: Hobbs----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| HgB: Hobbs----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Ho: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| HoB: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| HoC: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| HoC2: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Hp: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| HpB: Holdrege----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Hr: Hord----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| InB: Inavale----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.70 0.98 |
| Ke: Kenesaw----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of gravel | | Potential source of sand | |
|------------------------------|------------------|--|--------------|--|--------------|
| | | Rating class | Value | Rating class | Value |
| KgB: Kenesaw----- | 55 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Coly----- | 45 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Lf: Lex----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.00 0.91 |
| Lg: Lex----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.00 0.91 |
| LoB: Libory----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Bottom layer Thickest layer | 0.00 0.62 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Pm: Platte----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.00 0.97 |
| Ru: Rusco----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Sc: Scott----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| SmB: Simeon----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.10 0.91 |
| To: Tryon----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.65 |
| UcF: Uly----- | 60 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| Coly----- | 40 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| UsF: Ustorthents----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Poor Bottom layer Thickest layer | 0.00 0.00 |
| VaB: Valentine----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.70 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The ratings given for the thickest layer are for the thickest layer above and excluding the bottom layer. The numbers in the value columns range from 0.00 to 0.99. The greater the value, the greater the likelihood that the bottom layer or thickest layer of the soil is a source of sand or gravel. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of gravel | | Potential source of sand | |
|--------------------------|------------------|--|--------------|--|--------------|
| | | Rating class | Value | Rating class | Value |
| VaD: Valentine----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.70 |
| VaF: Valentine----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.70 |
| VbD: Valentine----- | 65 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.70 |
| Els----- | 35 | Poor Bottom layer Thickest layer | 0.00 0.00 | Good Thickest layer | 0.65 |
| W: Water----- | 100 | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Poor Bottom layer Thickest layer | 0.00 0.00 | Fair Thickest layer Bottom layer | 0.07 0.10 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of reclamation material | | Potential source of roadfill | | Potential source of topsoil | |
|--------------------------|------------------|---|----------------------|--|--------------|---|--------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Fair Low content of organic matter Sodium content | 0.12 0.97 | Not Rated Depth to saturated zone | 0.76 | Fair Depth to saturated zone Sodium content | 0.76 0.98 |
| Bo: Boel----- | 100 | Poor Too sandy Low content of organic matter | 0.00 0.12 | Not Rated Depth to saturated zone | 0.76 | Poor Too sandy Depth to saturated zone | 0.00 0.76 |
| Bu: Butler----- | 100 | Poor Too clayey Too acid Water erosion | 0.00 0.84 0.99 | Not Rated Depth to saturated zone Shrink-swell | 0.00 0.13 | Poor Too Clayey Depth to saturated zone | 0.00 0.00 |
| CaC: Coly----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| CaD: Coly----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Fair Slope | 0.96 |
| CaF: Coly----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated Slope | 0.98 | Poor Slope | 0.00 |
| CkB: Coly----- | 60 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| Kenesaw----- | 40 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| CoD2: Coly----- | 60 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Fair Slope | 0.96 |
| Uly----- | 40 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Fair Slope | 0.96 |
| CoF2: Coly----- | 60 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated Slope | 0.98 | Poor Slope | 0.00 |
| Uly----- | 40 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated Slope | 0.98 | Poor Slope | 0.00 |
| De: Detroit----- | 100 | Poor Too clayey Water erosion | 0.00 0.99 | Not Rated Shrink-swell | 0.66 | Poor Too Clayey | 0.00 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of reclamation material | | Potential source of roadfill | | Potential source of topsoil | |
|--------------------------|------------------|--|------------------------------|--|--------------|---|------------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Fm: Fillmore----- | 100 | Poor Too clayey Too acid Water erosion | 0.00 0.95 0.99 | Not Rated Depth to saturated zone Shrink-swell | 0.00 0.21 | Poor Too Clayey Depth to saturated zone | 0.00 0.00 |
| Gb: Gibbon----- | 100 | Fair Low content of organic matter | 0.88 | Not Rated Depth to saturated zone | 0.76 | Fair Depth to saturated zone | 0.76 |
| Gc: Gibbon----- | 100 | Fair Salinity Low content of organic matter Water erosion Sodium content | 0.88 0.88 0.90 0.97 | Not Rated Depth to saturated zone Shrink-swell | 0.76 0.93 | Fair Salinity Depth to saturated zone Sodium content | 0.50 0.76 0.98 |
| Go: Gothenburg----- | 100 | Poor Droughty Low content of organic matter Too sandy | 0.00 0.12 0.50 | Not Rated Depth to saturated zone | 0.00 | Poor Depth to saturated zone Rock fragments Too sandy Hard to reclaim | 0.00 0.03 0.50 0.98 |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Fair Low content of organic matter | 0.12 | Not Rated | | Good | |
| HeC: Hersh----- | 100 | Fair Low content of organic matter | 0.12 | Not Rated | | Good | |
| HeD: Hersh----- | 100 | Fair Low content of organic matter | 0.12 | Not Rated | | Fair Slope | 0.96 |
| Hf: Hobbs----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| HgB: Hobbs----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| Ho: Holdrege----- | 100 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| HoB: Holdrege----- | 100 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| HoC: Holdrege----- | 100 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of reclamation material | | Potential source of roadfill | | Potential source of topsoil | |
|------------------------------|------------------|--|------------------------------|--------------------------------------|-------|---|------------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HoC2: Holdrege----- | 100 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| Hp: Holdrege----- | 100 | Fair Water erosion | 0.90 | Not Rated | | Good | |
| HpB: Holdrege----- | 100 | Fair Water erosion | 0.90 | Not Rated | | Good | |
| Hr: Hord----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| InB: Inavale----- | 100 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.81 | Not Rated | | Poor Too sandy | 0.00 |
| Ke: Kenesaw----- | 100 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| KgB: Kenesaw----- | 55 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated | | Good | |
| Coly----- | 45 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated | | Good | |
| Lf: Lex----- | 100 | Poor Too sandy Low content of organic matter Water erosion | 0.00 0.12 0.99 | Not Rated Depth to saturated zone | 0.53 | Poor Too sandy Rock fragments Depth to saturated zone Hard to reclaim | 0.00 0.03 0.53 0.98 |
| Lg: Lex----- | 100 | Fair Low content of organic matter Sodium content Salinity Water erosion | 0.12 0.78 0.88 0.99 | Not Rated Depth to saturated zone | 0.76 | Poor Salinity Depth to saturated zone Sodium content | 0.00 0.76 0.78 |
| LoB: Libory----- | 100 | Poor Wind erosion Too sandy Low content of organic matter Water erosion | 0.00 0.00 0.12 0.90 | Not Rated Depth to saturated zone | 0.76 | Poor Too sandy Depth to saturated zone | 0.00 0.76 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of reclamation material | | Potential source of roadfill | | Potential source of topsoil | |
|--------------------------|------------------|--|------------------------------|--|--------------|--|--------------------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ma: Massie----- | 100 | Poor Too clayey Too acid Water erosion | 0.00 0.84 0.99 | Not Rated Depth to saturated zone Shrink-swell | 0.00 0.00 | Poor Depth to saturated zone Too Clayey | 0.00 0.00 |
| Pm: Platte----- | 100 | Poor Too sandy Low content of organic matter Droughty | 0.00 0.12 0.14 | Not Rated Depth to saturated zone | 0.53 | Poor Hard to reclaim Too sandy Rock fragments Depth to saturated zone Hard to reclaim | 0.00 0.00 0.03 0.53 0.98 |
| Ru: Rusco----- | 100 | Fair Low content of organic matter Water erosion Too clayey | 0.12 0.90 0.95 | Not Rated Depth to saturated zone | 0.00 | Poor Depth to saturated zone Too Clayey | 0.00 0.69 |
| Sc: Scott----- | 100 | Poor Too clayey Water erosion Too acid | 0.00 0.90 0.95 | Not Rated Depth to saturated zone Shrink-swell | 0.00 0.06 | Poor Too Clayey Depth to saturated zone | 0.00 0.00 |
| SmB: Simeon----- | 100 | Poor Too sandy Low content of organic matter Droughty | 0.00 0.12 0.87 | Not Rated | | Poor Too sandy | 0.00 |
| To: Tryon----- | 100 | Poor Too sandy Droughty Low content of organic matter | 0.00 0.45 0.88 | Not Rated Depth to saturated zone | 0.00 | Poor Too sandy Depth to saturated zone | 0.00 0.00 |
| UcF: Uly----- | 60 | Fair Low content of organic matter Water erosion | 0.12 0.90 | Not Rated Slope | 0.98 | Poor Slope | 0.00 |
| Coly----- | 40 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated Slope | 0.98 | Poor Slope | 0.00 |
| UsF: Ustorthents----- | 100 | Fair Low content of organic matter Water erosion | 0.88 0.90 | Not Rated Slope | 0.02 | Poor Slope | 0.00 |
| VaB: Valentine----- | 100 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.23 | Not Rated | | Poor Too sandy | 0.00 |

CONSTRUCTION MATERIALS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 0.99. The smaller the value, the greater the limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct. of map unit | Potential source of reclamation material | | Potential source of roadfill | | Potential source of topsoil | |
|--------------------------|------------------|--|------------------------------|--------------------------------------|-------|---|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| VaD: Valentine----- | 100 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.23 | Not Rated | | Poor Too sandy | 0.00 |
| VaF: Valentine----- | 100 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.23 | Not Rated | | Poor Too sandy Slope | 0.00 0.16 |
| VbD: Valentine----- | 65 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.23 | Not Rated | | Poor Too sandy | 0.00 |
| Els----- | 35 | Poor Too sandy Wind erosion Low content of organic matter Droughty | 0.00 0.00 0.12 0.22 | Not Rated Depth to saturated zone | 0.76 | Poor Too sandy Depth to saturated zone | 0.00 0.76 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Fair Low content of organic matter Sodium content | 0.88 0.97 | Not Rated Depth to saturated zone | 0.91 | Fair Depth to saturated zone Rock fragments Sodium content | 0.91 0.97 0.98 |

RECREATIONAL INTERPRETATIONS
Kearney County, Nebraska

Recreation

The soils of the survey area are rated in the following tables according to limitations that affect their suitability for recreation. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the recreational uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

The ratings in the tables are based on restrictive soil features, such as wetness, slope, and texture of the surface layer. Susceptibility to flooding is considered. Not considered in the ratings, but important in evaluating a site, are the location and accessibility of the area, the size and shape of the area and its scenic quality, vegetation, access to water, potential water impoundment sites, and access to public sewer lines. The capacity of the soil to absorb septic tank effluent and the ability of the soil to support vegetation also are important. Soils that are subject to flooding are limited for recreational uses by the duration and intensity of flooding and the season when flooding occurs. In planning recreational facilities, onsite assessment of the height, duration, intensity, and frequency of flooding is essential.

The information in this table can be supplemented by other information in this survey, for example, interpretations for building site development, construction materials, sanitary facilities, and water management.

Camp areas require site preparation, such as shaping and leveling the tent and parking areas, stabilizing roads and intensively used areas, and installing sanitary facilities and utility lines. Camp areas are subject to heavy foot traffic and some vehicular traffic. The ratings are based on the soil properties that affect the ease of developing camp areas and the performance of the areas after development. Slope, stoniness, and depth to bedrock or a cemented pan are the main concerns affecting the development of camp areas.

The soil properties that affect the performance of the areas after development are those that influence trafficability and promote the growth of vegetation, especially in heavily used areas. For good trafficability, the surface of camp areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Picnic areas are subject to heavy foot traffic. Most vehicular traffic is confined to access roads and parking areas. The ratings are based on the soil properties that affect the ease of developing picnic areas and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of picnic areas. For good trafficability, the surface of picnic areas should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Playgrounds require soils that are nearly level, are free of stones, and can withstand intensive foot traffic. The ratings are based on the soil properties that affect the ease of developing playgrounds and that influence trafficability and the growth of vegetation after development. Slope and stoniness are the main concerns affecting the development of playgrounds. For good trafficability, the surface of the playgrounds should absorb rainfall readily, remain firm under heavy foot traffic, and not be dusty when dry. The soil properties that influence trafficability are texture of the surface layer, depth to a water table, ponding, flooding, permeability, and large stones. The soil properties that affect the growth of plants are depth to bedrock or a cemented pan, permeability, and toxic substances in the soil.

Paths and trails for hiking and horseback riding should require little or no slope modification through cutting and filling. The ratings are based on the soil properties that affect trafficability and erodibility. These properties are stoniness, depth to a water table, ponding, flooding, slope, and texture of the surface layer.

Golf fairways are subject to heavy foot traffic and some light vehicular traffic. Cutting or filling may be required. Irrigation is not considered in the ratings. The ratings are based on the soil properties that affect plant growth and trafficability after vegetation is established. The properties that affect plant growth are reaction; depth to a water table; ponding; depth to bedrock or a cemented pan; the available water capacity in the upper 40 inches; the content of salts, sodium, or calcium carbonate; and sulfidic materials. The properties that affect trafficability are flooding, depth to a water table, ponding, slope, stoniness, and the amount of sand, clay, or organic matter in the surface layer. The suitability of the soil for traps, tees, roughs, and greens is not considered in the ratings.

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Camp areas | | Picnic areas | | Playgrounds | |
|-----------------------------|--------------------------|--|--------------------------|--|----------------------|--|--------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Somewhat limited Depth to saturated zone | 0.03 | Somewhat limited Depth to saturated zone | 0.07 |
| Bo: Boel----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Somewhat limited Depth to saturated zone | 0.03 | Somewhat limited Depth to saturated zone | 0.07 |
| Bu: Butler----- | 100 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 |
| CaC: Coly----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.87 |
| CaD: Coly----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| CkB: Coly----- | 60 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| Kenesaw----- | 40 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| CoD2: Coly----- | 60 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| Uly----- | 40 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Slope | 1.00 |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| De: Detroit----- | 100 | Somewhat limited Restricted permeability | 0.39 | Somewhat limited Restricted permeability | 0.39 | Somewhat limited Restricted permeability | 0.39 |
| Fm: Fillmore----- | 100 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 | Very limited Restricted permeability Depth to saturated zone | 1.00 1.00 |
| Gb: Gibbon----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.07 | Somewhat limited Depth to saturated zone | 0.03 | Somewhat limited Depth to saturated zone | 0.07 |
| Gc: Gibbon----- | 100 | Very limited Flooding Salinity Depth to saturated zone | 1.00 1.00 0.07 | Very limited Salinity Depth to saturated zone | 1.00 0.03 | Very limited Salinity Depth to saturated zone | 1.00 0.07 |
| Go: Gothenburg----- | 100 | Very limited Flooding Depth to saturated zone Too sandy | 1.00 1.00 0.42 | Very limited Depth to saturated zone Too sandy Flooding | 1.00 0.42 0.40 | Very limited Flooding Depth to saturated zone Too sandy | 1.00 1.00 0.42 |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| HeC: Hersh----- | 100 | Not limited | | Not limited | | Somewhat limited | |

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Camp areas | | Picnic areas | | Playgrounds | |
|------------------------------|--------------------------|--|----------------------|--|--------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HeD: Hersh----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Slope | 0.87 |
| Hf: Hobbs----- | 100 | Very limited Flooding | 1.00 | Not limited | | Very limited Slope | 1.00 |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Somewhat limited Flooding | 0.40 | Somewhat limited Flooding | 0.60 |
| Ho: Holdrege----- | 100 | Not limited | | Not limited | | Very limited Flooding Slope | 1.00 0.00 |
| HoB: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HoC: Holdrege----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| HoC2: Holdrege----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.87 |
| Hp: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HpB: Holdrege----- | 100 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| Hr: Hord----- | 100 | Not limited | | Not limited | | Not limited | |
| InB: Inavale----- | 100 | Very limited Flooding Too sandy | 1.00 0.96 | Somewhat limited Too sandy | 0.96 | Somewhat limited Too sandy Slope | 0.96 0.00 |
| Ke: Kenesaw----- | 100 | Not limited | | Not limited | | Not limited | |
| KgB: Kenesaw----- | 55 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| Coly----- | 45 | Not limited | | Not limited | | Somewhat limited Slope | 0.00 |
| Lf: Lex----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.39 | Somewhat limited Depth to saturated zone | 0.19 | Somewhat limited Depth to saturated zone | 0.39 |
| Lg: Lex----- | 100 | Very limited Flooding Salinity Depth to saturated zone | 1.00 1.00 0.07 | Very limited Salinity Depth to saturated zone | 1.00 0.03 | Very limited Salinity Depth to saturated zone | 1.00 0.07 |
| LoB: Libory----- | 100 | Somewhat limited Too sandy Depth to saturated zone | 0.94 0.07 | Somewhat limited Too sandy Depth to saturated zone | 0.94 0.03 | Somewhat limited Too sandy Depth to saturated zone Slope | 0.94 0.07 0.00 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Depth to saturated zone Restricted permeability | 1.00 1.00 | Very limited Depth to saturated zone Restricted permeability | 1.00 1.00 | Very limited Depth to saturated zone Restricted permeability | 1.00 1.00 |
| Pm: Platte----- | 100 | Very limited Flooding Depth to saturated zone | 1.00 0.39 | Somewhat limited Depth to saturated zone | 0.19 | Somewhat limited Flooding Depth to saturated zone | 0.60 0.39 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| Sc: Scott----- | 100 | Very limited | | Very limited | | Very limited | |

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Camp areas | | Picnic areas | | Playgrounds | |
|-----------------------------|--------------------------|---|--------------|---|--------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| SmB: Simeon----- | 100 | Depth to saturated zone Restricted permeability | 1.00 | Depth to saturated zone Restricted permeability | 1.00 | Depth to saturated zone Restricted permeability | 1.00 |
| | | Not limited | 1.00 | Not limited | 1.00 | Somewhat limited Slope | 0.00 |
| To: Tryon----- | 100 | Very limited Flooding | 1.00 | Very limited Depth to saturated zone Too sandy | 1.00 | Very limited Depth to saturated zone Too sandy | 1.00 |
| | | Depth to saturated zone Too sandy | 0.95 | Too sandy | 0.95 | Too sandy | 0.95 |
| UcF: Uly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Coly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| UsF: Ustorthents----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Too sandy | 0.96 | Somewhat limited Too sandy Slope | 0.96 0.00 |
| VaD: Valentine----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Too sandy | 0.96 | Very limited Slope Too sandy | 1.00 0.96 |
| VaF: Valentine----- | 100 | Somewhat limited Too sandy Slope | 0.96 0.84 | Somewhat limited Too sandy Slope | 0.96 0.84 | Very limited Slope Too sandy | 1.00 0.96 |
| VbD: Valentine----- | 65 | Somewhat limited Too sandy | 0.96 | Somewhat limited Too sandy | 0.96 | Very limited Slope Too sandy | 1.00 0.96 |
| Els----- | 35 | Somewhat limited Too sandy Depth to saturated zone | 0.95 0.07 | Somewhat limited Too sandy Depth to saturated zone | 0.95 0.03 | Somewhat limited Too sandy Depth to saturated zone Slope | 0.95 0.07 0.00 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Very limited Flooding | 1.00 | Not limited | | Not limited | |

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Paths and trails | | Golf fairways | |
|-----------------------------|--------------------------|---|----------------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Not limited | | Somewhat limited Depth to saturated zone | 0.03 |
| Bo: Boel----- | 100 | Not limited | | Somewhat limited Depth to saturated zone | 0.03 |
| Bu: Butler----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| CaC: Coly----- | 100 | Not limited | | Not limited | |
| CaD: Coly----- | 100 | Not limited | | Somewhat limited Slope | 0.04 |
| CaF: Coly----- | 100 | Somewhat limited Slope | 0.02 | Very limited Slope | 1.00 |
| CkB: Coly----- | 60 | Not limited | | Not limited | |
| Kenesaw----- | 40 | Not limited | | Not limited | |
| CoD2: Coly----- | 60 | Not limited | | Somewhat limited Slope | 0.04 |
| Uly----- | 40 | Not limited | | Somewhat limited Slope | 0.04 |
| CoF2: Coly----- | 60 | Somewhat limited Slope | 0.02 | Very limited Slope | 1.00 |
| Uly----- | 40 | Somewhat limited Slope | 0.02 | Very limited Slope | 1.00 |
| De: Detroit----- | 100 | Not limited | | Not limited | |
| Fm: Fillmore----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| Gb: Gibbon----- | 100 | Not limited | | Somewhat limited Depth to saturated zone | 0.03 |
| Gc: Gibbon----- | 100 | Not limited | | Very limited Salinity Depth to saturated zone | 1.00 0.03 |
| Go: Gothenburg----- | 100 | Very limited Depth to saturated zone Too sandy Flooding | 1.00 0.42 0.40 | Very limited Flooding Droughty Depth to saturated zone | 1.00 1.00 1.00 |
| GP: Pits----- | 100 | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Not limited | | Not limited | |
| HeC: Hersh----- | 100 | Not limited | | Not limited | |
| HeD: Hersh----- | 100 | Not limited | | Somewhat limited Slope | 0.04 |
| Hf: Hobbs----- | 100 | Not limited | | Somewhat limited Flooding | 0.60 |
| HgB: Hobbs----- | 100 | Somewhat limited Flooding | 0.40 | Very limited Flooding | 1.00 |
| Ho: Holdrege----- | 100 | Not limited | | Not limited | |
| HoB: Holdrege----- | 100 | Not limited | | Not limited | |
| HoC: Holdrege----- | 100 | Not limited | | Not limited | |

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Paths and trails | | Golf fairways | |
|------------------------------|--------------------------|---|--------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HoC2: Holdrege----- | 100 | Not limited | | Not limited | |
| Hp: Holdrege----- | 100 | Not limited | | Not limited | |
| HpB: Holdrege----- | 100 | Not limited | | Not limited | |
| Hr: Hord----- | 100 | Not limited | | Not limited | |
| InB: Inavale----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Droughty | 0.19 |
| Ke: Kenesaw----- | 100 | Not limited | | Not limited | |
| KgB: Kenesaw----- | 55 | Not limited | | Not limited | |
| Coly----- | 45 | Not limited | | Not limited | |
| Lf: Lex----- | 100 | Not limited | | Somewhat limited Depth to saturated zone | 0.19 |
| Lg: Lex----- | 100 | Not limited | | Very limited Salinity Depth to saturated zone | 1.00 0.03 |
| LoB: Libory----- | 100 | Somewhat limited Too sandy | 0.94 | Somewhat limited Depth to saturated zone | 0.03 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| Pm: Platte----- | 100 | Not limited | | Somewhat limited Flooding Droughty Depth to saturated zone | 0.60 0.35 0.19 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| Sc: Scott----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| SmB: Simeon----- | 100 | Not limited | | Somewhat limited Droughty | 0.11 |
| To: Tryon----- | 100 | Very limited Depth to saturated zone Too sandy | 1.00 0.95 | Very limited Depth to saturated zone Droughty | 1.00 0.56 |
| UcF: Uly----- | 60 | Somewhat limited Slope | 0.02 | Very limited Slope | 1.00 |
| Coly----- | 40 | Somewhat limited Slope | 0.02 | Very limited Slope | 1.00 |
| UsF: Ustorthents----- | 100 | Somewhat limited Slope | 0.98 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Droughty | 0.71 |
| VaD: Valentine----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Droughty | 0.71 |
| VaF: Valentine----- | 100 | Somewhat limited Too sandy | 0.96 | Somewhat limited Slope Droughty | 0.84 0.71 |
| VbD: Valentine----- | 65 | Somewhat limited Too sandy | 0.96 | Somewhat limited Droughty | 0.71 |
| Els----- | 35 | Somewhat limited | | Somewhat limited | |

RECREATIONAL INTERPRETATIONS--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Paths and trails | | Golf fairways | |
|-----------------------------|--------------------------|---------------------------------------|-------|--|--------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| | | Too sandy | 0.95 | Droughty Depth to saturated zone | 0.74 0.03 |
| W: Water----- | 100 | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Not limited | | Not limited | |

WILDLIFE INTERPRETATIONS
Kearney County, Nebraska

Use and Explanation of Wildlife Interpretations

Soils directly affect the kind and amount of vegetation that is available to wildlife as food and cover. They also affect the development of water impoundments. The kind and abundance of wildlife that populate an area depend largely on the amount and distribution of food, cover, water, and living space. If any one of these elements is missing, inadequate, or inaccessible, wildlife will be scarce or will not inhabit the area. If the soils have the potential, wildlife habitat can be created or improved by planting appropriate vegetation, properly managing the existing plant cover, and fostering the natural establishment of desirable plants.

In the Wildlife Interpretations table, the soils in the survey area are rated according to their potential for providing habitat for various kinds of wildlife. This information can be used in planning parks, wildlife refuges, nature study areas, and other developments for wildlife; in selecting soils that are suitable for establishing, improving, or maintaining specific elements of wildlife habitat; and in determining the intensity of management needed for each element of the habitat.

Suitability Ratings

The potential of the soil is rated good, fair, poor, or very poor.

Good - means that the element of wildlife habitat or the kind of habitat is easily created, improved, or maintained. Few or no limitations affect management, and satisfactory results can be expected if the soil is used for the designated purpose.

Fair - means that the element of wildlife habitat or kind of habitat can be created, improved, or maintained in most places. Moderately intensive management is required for satisfactory results.

Poor - means that limitations are severe for the designated element or kind of wildlife habitat. Habitat can be created, improved, or maintained in most places, but management is difficult and requires intensive effort.

Very Poor - means that limitations are very severe for the designated element or kind of wildlife habitat. Habitat is difficult to create, improve, or maintain in most places, and management is difficult and requires intensive effort.

Description of Wildlife Habitat Elements

Openland habitat consists of croplands, pastures, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. The kind of wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, killdeer, cottontail rabbit, red fox, and coyote.

Woodland habitat consists of hardwood or conifers, or a mixture of these and associated grasses, legumes and wild herbaceous plants. Examples of wildlife attracted to this habitat are wild turkey, thrushes, woodpeckers, owl, tree squirrels, raccoon, and deer.

Wetland habitat consists of water-tolerant plants in open, marshy or swampy, shallow water areas. Examples of wildlife attracted to this habitat are ducks, geese, herons, bitterns, rails, kingfishers, shorebirds, muskrat, mink, and beaver.

The elements of wildlife habitat are described in the following paragraphs.

Grain and seed crops are domestic grains and seed-producing herbaceous plants. Soil properties and features that affect the growth of grain and seed crops are depth of the root zone, texture of the surface layer, available water capacity, wetness, slope, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of grain and seed crops are corn, wheat, oats, and barley.

Grasses and legumes are domestic perennial grasses and herbaceous legumes. Soil properties and features that affect the growth of grasses and legumes are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, flooding, and slope. Soil temperature and soil moisture also are considerations. Examples of grasses and legumes are fescue, lovegrass, bromegrass, clover, and alfalfa.

Wild herbaceous plants are native or naturally established grasses and forbs, including weeds. Soil properties and features that affect the growth of these plants are depth of the root zone, texture of the surface layer, available water capacity, wetness, surface stoniness, and flooding. Soil temperature and soil moisture also are considerations. Examples of wild herbaceous plants are bluestem, goldenrod, beggarweed, wheatgrass, and grama.

Hardwood trees and woody understory produce nuts or other fruit, buds, catkins, twigs, bark, and foliage. Soil properties and features that affect the growth of hardwood trees and shrubs are depth of the root zone, available water capacity, and wetness. Examples of these plants are oak, poplar, cherry, sweetgum, apple, hawthorn, dogwood, hickory, blackberry, and blueberry. Examples of fruit-producing shrubs that are suitable for planting on soils rated good are Russian-olive, autumn-olive, and crabapple.

Coniferous plants furnish browse and seeds. Soil properties and features that affect the growth of coniferous trees, shrubs, and ground cover are depth of the root zone, available water capacity, and wetness. Examples of coniferous plants are pine, spruce, fir, cedar, and juniper.

Shrubs are bushy woody plants that produce fruit, buds, twigs, bark, and foliage. Soil properties and features that affect the growth of shrubs are depth of the root zone, available water capacity, salinity, and soil moisture. Examples of shrubs are fragrant sumac, chokecherry, American plum, sand plum, and garden currant.

Wetland plants are annual and perennial wild herbaceous plants that grow on moist or wet sites. Submerged or floating aquatic plants are excluded. Soil properties and features affecting wetland plants are texture of the surface layer, wetness, reaction, salinity, slope, and surface stoniness. Examples of wetland plants are smartweed, wild millet, saltgrass, cordgrass, rushes, sedges, and cattails.

Shallow water areas have an average depth of less than 5 feet. Some are naturally wet areas. Others are created by dams, levees, or other water-control structures. Soil properties and features affecting shallow water areas are depth to bedrock, wetness, surface stoniness, slope, and permeability. Examples of shallow water areas are marshes, waterfowl feeding areas, and ponds.

The habitat for various kinds of wildlife is described in the following paragraphs.

Habitat for openland wildlife consists of cropland, pasture, meadows, and areas that are overgrown with grasses, herbs, shrubs, and vines. These areas produce grain and seed crops, grasses and legumes, and wild herbaceous plants. Wildlife attracted to these areas include bobwhite quail, pheasant, meadowlark, field sparrow, cottontail, red fox and coyote.

Habitat for woodland wildlife consists of areas of deciduous and/or coniferous plants and associated grasses, legumes, and wild herbaceous plants. Wildlife attracted to these areas include wild turkey, thrushes, woodpeckers, squirrels, gray fox, raccoon, and deer.

Habitat for wetland wildlife consists of open, marshy or swampy shallow water areas. Some of the wildlife attracted to such areas are ducks, geese, herons, shore birds, muskrat, mink, and beaver.

Habitat for rangeland wildlife consists of areas of shrubs and wild herbaceous plants. Wildlife attracted to rangeland include antelope, deer, cottontail rabbit, prairie chicken, meadowlark, quail, and pheasant.

WILDLIFE INTERPRETATIONS
Kearney County, Nebraska

| Map symbol and soil name | Potential for habitat elements | | | | | | | | Potential as habitat for-- | | | |
|-----------------------------|--------------------------------|---------------------------|-----------------------------------|------------------------|---------------------------|--------|-------------------|---------------------------|--------------------------------|--------------------------------|--------------------------|---------------------------------|
| | Grain and seed crops | Grasses and legumes | Wild herba- ceous plants | Hard- wood trees | Conif- erous plants | Shrubs | Wetland plants | Shallow water areas | Open- land wild- life | Wood- land wild- life | Wetland wild- life | Range- land wild- life |
| Ad: ALDA----- | Fair | Fair | Fair | Good | Good | Good | Fair | Fair | Fair | Good | Fair | Good |
| Bo: BOEL----- | Fair | Fair | Good | Good | Good | Good | Fair | Fair | Fair | Good | Poor | Fair |
| Bu: BUTLER----- | Good | Good | Good | --- | Good | Good | Fair | Fair | Good | --- | Fair | Good |
| CaC: COLY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| CaD: COLY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| CaF: COLY----- | Poor | Fair | Fair | Fair | Fair | Fair | Very poor | Very poor | Fair | Fair | Very poor | Fair |
| CkB: COLY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| KENESAW----- | Good | Good | Good | Good | Good | Good | Very poor | Very poor | Good | Good | Very poor | Good |
| CoD2: COLY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| ULY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Good |
| CoF2: COLY----- | Poor | Fair | Fair | Fair | Fair | Fair | Very poor | Very poor | Fair | Fair | Very poor | Fair |
| ULY----- | Poor | Fair | Good | Good | Fair | Fair | Very poor | Very poor | Poor | Good | Very poor | Fair |
| De: DETROIT----- | Good | Good | Good | --- | --- | Good | Good | Good | Good | --- | Good | Good |
| Fm: FILLMORE----- | Fair | Good | Fair | Fair | Fair | Fair | Good | Fair | Fair | Fair | Good | Fair |
| Gb: GIBBON----- | Good | Good | Good | Good | Fair | Good | Fair | Good | Good | Good | Fair | Good |
| Gc: GIBBON----- | Poor | Poor | Poor | Good | Fair | Poor | Fair | Fair | Poor | Fair | Fair | Poor |
| Go: GOTHENBURG----- | Very poor | Very poor | Fair | Poor | Fair | Fair | Fair | Good | Poor | Poor | Fair | Fair |
| GP: PITS----- | Very poor | Very poor | Poor | Poor | Poor | Poor | Very poor | Fair | Very poor | Very poor | Poor | Poor |
| HeB: HERSH----- | Fair | Good | Good | Good | Good | Good | Very poor | Very poor | Fair | Good | Very poor | Good |
| HeC: HERSH----- | Fair | Good | Good | Good | Good | Good | Very poor | Very poor | Fair | Good | Very poor | Good |
| HeD: HERSH----- | Fair | Good | Good | Good | Good | Good | Very poor | Very poor | Fair | Good | Very poor | Good |
| Hf: HOBBS----- | Good | Good | Good | Good | Good | Good | Poor | Poor | Good | Good | Poor | Good |
| HgB: HOBBS----- | Poor | Fair | Fair | Fair | Fair | Fair | Very poor | Very poor | Fair | Fair | Very poor | Fair |
| Ho: HOLDREGE----- | Good | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Good | Good | Very poor | Fair |

WILDLIFE INTERPRETATIONS--Continued
Kearney County, Nebraska

| Map symbol and soil name | Potential for habitat elements | | | | | | | | Potential as habitat for-- | | | |
|-------------------------------------|--------------------------------|---------------------------|-----------------------------------|------------------------|---------------------------|--------------|-------------------|---------------------------|--------------------------------|--------------------------------|--------------------------|---------------------------------|
| | Grain and seed crops | Grasses and legumes | Wild herba- ceous plants | Hard- wood trees | Conif- erous plants | Shrubs | Wetland plants | Shallow water areas | Open- land wild- life | Wood- land wild- life | Wetland wild- life | Range- land wild- life |
| HoB: HOLDREGE----- | Good | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Good | Good | Very poor | Fair |
| HoC: HOLDREGE----- | Fair | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| HoC2: HOLDREGE----- | Fair | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| Hp: HOLDREGE----- | Good | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Good | Good | Very poor | Fair |
| HpB: HOLDREGE----- | Good | Good | Fair | Good | Fair | Fair | Very poor | Very poor | Good | Good | Very poor | Fair |
| Hr: HORD----- | Good | Good | Good | Good | Good | Good | Very poor | Very poor | Good | Good | Very poor | Good |
| InB: INAVAL----- | Fair | Fair | Good | Fair | Fair | Fair | Very poor | Very poor | Fair | Fair | Very poor | Good |
| Ke: KENESAW----- | Good | Good | Good | Good | Good | Good | Very poor | Very poor | Good | Good | Very poor | Good |
| KgB: KENESAW----- | Good | Good | Good | Good | Good | Good | Very poor | Very poor | Good | Good | Very poor | Good |
| COLY----- | Fair | Good | Good | Good | Fair | Fair | Very poor | Very poor | Fair | Good | Very poor | Fair |
| Lf: LEX----- | Fair | Fair | Good | Fair | Good | Good | Fair | Fair | Fair | Fair | Fair | Good |
| Lg: LEX----- | Poor | Poor | Poor | Good | Fair | Poor | Fair | Fair | Poor | Fair | Fair | Poor |
| LoB: LIBORY----- | Fair | Fair | Good | Good | Good | Good | Poor | Very poor | Fair | Good | Very poor | Good |
| M-W: MISCELLANEOUS WATER----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Ma: MASSIE----- | Very poor | Very poor | Very poor | Very poor | Very poor | Very poor | Good | Good | Very poor | Very poor | Good | Very poor |
| Pm: PLATTE----- | Fair | Good | Fair | Poor | Fair | Good | Fair | Good | Fair | Poor | Good | Fair |
| Ru: RUSCO----- | Good | Good | Poor | Good | Good | Good | Good | Good | Fair | Good | Good | Fair |
| Sc: SCOTT----- | Poor | Fair | Fair | Fair | Fair | Poor | Good | Good | Fair | Fair | Good | Fair |
| SmB: SIMEON----- | Poor | Poor | Fair | Poor | Poor | Poor | Very poor | Very poor | Poor | Poor | Very poor | Fair |
| To: TRYON----- | Very poor | Poor | Fair | Poor | Poor | Fair | Good | Good | Poor | Poor | Good | Fair |
| UcF: ULY----- | Poor | Fair | Good | Good | Fair | Fair | Very poor | Very poor | Poor | Good | Very poor | Fair |
| COLY----- | Poor | Fair | Fair | Fair | Fair | Fair | Very poor | Very poor | Fair | Fair | Very poor | Fair |
| UsF: USTORTHE----- | Very poor | Very poor | Fair | Good | Good | Fair | Poor | Very poor | Poor | Fair | Very poor | Fair |

WILDLIFE INTERPRETATIONS--Continued
Kearney County, Nebraska

| Map symbol and soil name | Potential for habitat elements | | | | | | | | Potential as habitat for-- | | | |
|-----------------------------|--------------------------------|---------------------------|-----------------------------------|------------------------|---------------------------|--------|-------------------|---------------------------|--------------------------------|--------------------------------|--------------------------|---------------------------------|
| | Grain and seed crops | Grasses and legumes | Wild herba- ceous plants | Hard- wood trees | Conif- erous plants | Shrubs | Wetland plants | Shallow water areas | Open- land wild- life | Wood- land wild- life | Wetland wild- life | Range- land wild- life |
| VaB: VALENTINE----- | Fair | Good | Fair | Poor | Fair | Fair | Very poor | Very poor | Fair | Poor | Very poor | Fair |
| VaD: VALENTINE----- | Poor | Fair | Fair | Poor | Poor | Poor | Very poor | Very poor | Fair | Poor | Very poor | Fair |
| VaF: VALENTINE----- | Poor | Fair | Fair | Poor | Poor | Poor | Very poor | Very poor | Fair | Poor | Very poor | Fair |
| VbD: VALENTINE----- | Poor | Fair | Fair | Poor | Poor | Poor | Very poor | Very poor | Fair | Poor | Very poor | Fair |
| ELS----- | Poor | Fair | Fair | Fair | Fair | Fair | Good | Fair | Fair | Fair | Fair | Fair |
| W: WATER----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Wa: WANN----- | Good | Good | Good | Good | Fair | Good | Poor | Fair | Good | Good | Fair | Good |

YIELDS PER ACRE OF PASTURE AND HAYLAND
Kearney County, Nebraska

Use and Explanation of Pastureland and Hayland Interpretations

This subsection provides information concerning the suitability of soils for the production of pasture and hayland. This subsection may contain pasture and hayland suitability groupings, land capability and yield estimates, yield estimates for individual grasses or legumes, or other information pertaining to the production of forage.

Pasture and Hayland Suitability Groupings

Soils are placed in pasture and hayland groups according to their suitability for the production of forage. The soils in each group are enough alike to be suited to the same grasses or legumes, to have similar limitations and hazards, to require similar management, and to have similar productivity and other responses to management. Thus, the pasture and hayland suitability group is a convenient way of grouping the soils for their management. If used, these groupings are identified and described in other reports in the subsection.

Yield Estimates

The average yields per acre that can be expected of the principal pasture or hayland crops, under a high level of management, are presented in this subsection. In any given year, yields may be higher or lower than those indicated in the tables because of variations in rainfall or other climatic factors. The yields are based mainly on the experience and records of farmers, conservationists, and extension agents. Available yield data from nearby counties and results of field trials and demonstrations are also considered.

Under good management, proper grazing is essential for the production of high quality forage, stand survival, and erosion control. Proper grazing helps plants maintain sufficient and generally vigorous top growth during the growing season. Brush control is essential in many areas, and weed control generally is needed. Rotation grazing and renovation are also important management practices.

The Pasture and Hayland table show yield estimates in tons per acre and animal unit months for pasture and hayland groups. An animal unit month is the amount of forage required by one animal unit (AU) for 30 days. On animal unit (AU) is one (1000 pound) mature cow and a calf up to weaning age (usually six months of age) or their equivalent. The Natural Resources Conservation Service uses 900 pounds of air dry forage as the amount needed to meet this requirement. To maintain a healthy and vigorous plant community, the degree of use should never be greater than 50 percent. Therefore only 25 percent of the total biomass grown is considered consumed by the grazing animal. Animal Unit Months can be converted to air dry pounds per acre production by multiplying the AUM by 30 days, then by 30 pounds per day, and then by four. This figure is the amount of total forage production.

Planners of management systems for individual fields or farms should consider the detailed information given in the description of each soil in the Nontechnical Description section. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

| Map symbol and soil name | Land capability | | Alfalfa hay | |
|-----------------------------|--------------------|-----|-------------|------|
| | N | I | N | I |
| | | | Tons | Tons |
| Ad: Alda----- | 3w | 3w | 3.60 | 5.00 |
| Bo: Boel----- | 3w | 3w | 3.40 | 4.00 |
| Bu: Butler----- | 2w | 2w | 2.90 | 5.00 |
| CaC: Coly----- | 3e | 3e | 2.00 | 5.00 |
| CaD: Coly----- | 4e | 4e | 1.60 | 3.50 |
| CaF: Coly----- | 6e | --- | --- | --- |
| CkB: Coly----- | 2e | 2e | 3.60 | 5.80 |
| Kenesaw----- | 2e | 2e | 3.60 | 5.80 |
| CoD2: Coly----- | 4e | 4e | 1.80 | 3.00 |
| Uly----- | 4e | 4e | 1.80 | 3.00 |
| CoF2: Coly----- | 6e | --- | --- | --- |
| Uly----- | 6e | --- | --- | --- |
| De: Detroit----- | 1 | 1 | 3.60 | 6.00 |
| Fm: Fillmore----- | 3w | 4w | 2.20 | --- |
| Gb: Gibbon----- | 2w | 2w | 4.20 | 5.80 |
| Gc: Gibbon----- | 4s | 3s | 2.30 | 3.50 |
| Go: Gothenburg----- | 7s | --- | --- | --- |
| GP: Pits----- | 8s | --- | --- | --- |
| HeB: Hersh----- | 3e | 2e | 2.20 | 5.00 |
| HeC: Hersh----- | 3e | 3e | 1.80 | 4.20 |
| HeD: Hersh----- | 4e | 4e | 1.20 | 3.50 |
| Hf: Hobbs----- | 2w | 2w | 4.80 | 6.00 |
| HgB: Hobbs----- | 6w | --- | --- | --- |
| Ho: Holdrege----- | 2c | 1 | 3.10 | 6.50 |
| HoB: Holdrege----- | 2e | 2e | 2.90 | 6.20 |
| HoC: Holdrege----- | 3e | 3e | 2.80 | 5.40 |
| HoC2: Holdrege----- | 3e | 3e | 2.00 | 5.20 |
| Hp: Holdrege----- | 2c | 1 | 3.10 | 6.50 |

(Yields in the "N" columns are for nonirrigated soils; those in the "I" columns are for irrigated soils. Yields are those that can be expected under a high level of nonirrigated and irrigated management by component. Absence of a yield indicates that the soil is not suited to the crop or the crop generally is not grown on the soil)
Animal-unit-month: The amount of forage or feed required to feed one animal unit (one cow, one horse, one mule, five sheep, or five goats) for 30 days.

| Map symbol and soil name | Land capability | | Alfalfa hay | |
|----------------------------------|--------------------|-----|-------------|------|
| | N | I | N | I |
| | | | Tons | Tons |
| HpB: Holdrege----- | 2e | 2e | 2.90 | 6.20 |
| Hr: Hord----- | 2c | 1 | 3.60 | 6.50 |
| InB: Inavale----- | 4e | 3e | 1.80 | 5.00 |
| Ke: Kenesaw----- | 2c | 1 | 2.80 | 6.10 |
| KgB: Kenesaw----- | 2e | 2e | 2.80 | 6.00 |
| Coly----- | 2e | 2e | 2.80 | 6.00 |
| Lf: Lex----- | 3w | 3w | 4.20 | 5.50 |
| Lg: Lex----- | 4s | 3s | 2.30 | 3.50 |
| LoB: Libory----- | 3e | 3e | 3.60 | 5.00 |
| M-W: Miscellaneous Water----- | --- | --- | --- | --- |
| Ma: Massie----- | 8w | --- | --- | --- |
| Pm: Platte----- | 4w | 4w | 2.40 | 2.90 |
| Ru: Rusco----- | 2w | 2w | 3.80 | 5.00 |
| Sc: Scott----- | 4w | --- | --- | --- |
| Smb: Simeon----- | 6s | 4s | --- | 3.00 |
| To: Tryon----- | 5w | --- | --- | --- |
| UcF: Uly----- | 6e | --- | --- | --- |
| Coly----- | 6e | --- | --- | --- |
| UsF: Ustorthents----- | 7s | --- | --- | --- |
| VaB: Valentine----- | 4e | 4e | 1.00 | 3.60 |
| VaD: Valentine----- | 6e | 4e | --- | 3.00 |
| VaF: Valentine----- | 6e | --- | --- | --- |
| VbD: Valentine----- | 6e | 4e | --- | 3.40 |
| Els----- | 4w | 4w | --- | 3.40 |
| W: Water----- | --- | --- | --- | --- |
| Wa: Wann----- | 2w | 2w | 4.20 | 5.80 |

CONSERVATION TREE AND SHRUB MANAGEMENT
Kearney County, Nebraska

A Conservation Tree/Shrub Suitability Group (CTSG), formerly Windbreak Suitability Group, is a physiographic unit or area having similar climatic and edaphic characteristics that control the selection and height growth of trees and shrubs.

In this table, the Conservation Tree and Shrub Grouping is expressed as a group index number. The group index for Conservation Tree and Shrub groups (CTSG) are a guide for species best suited for different kinds of soil and for prediction height, growth, and effectiveness. The groupings can be used when selection woody plants for windbreaks, wildlife plantings riparian buffers, reforestation, other environmental plantings, recreation, landscaping, wetland restoration or enhancement and critical area plantings. CTSG's are developed to assure satisfactory species selection and adaptation to specific conditions of soil, climate and physiography. CTSG's are a guide for selection species best suited for different kinds of soil and prediction height growth and effectiveness.

All soil series mapped in the state have been placed in 10 groups of similar soil characteristics. Groups 1, 2, 3, 4, 6, and 9 are further divided into subgroups. In addition, all groups provide information by Major Land Resource Areas.

Each tree or shrub species has certain climatic and physiographic limits. Within these parameters a tree or shrub may be well or poorly suited because of soil characteristics. Each tree or shrub also has definable potentials of height growth depending on the factors just mentioned. Accurate definitions of potential heights are necessary for proper windbreak planning and design.

Windbreaks protect livestock, buildings, roads and yards from wind and snow. They also protect fruit trees and gardens, and they furnish habitat for wildlife. Several rows of low-growing and high-growing broadleaf and coniferous trees and shrubs provide the most protection.

Field windbreaks are narrow plantings made at right angles to the prevailing wind and at specific intervals across the field. The interval depends on the erodibility of the soil. Field windbreaks protect cropland and crops from wind, help to keep snow on the fields, and provide food and cover for wildlife.

Environmental plantings help to beautify and screen houses and other buildings and to abate noise. The plants, mostly evergreen shrubs and trees, are closely spaced. To ensure plant survival, a healthy planting stock of suitable species should be planted properly on a well prepared site and maintained in good condition.

Windbreaks are often planted on land that did not grow trees originally. Knowledge of how trees perform on such land can be gained only by observing and recording their performance where trees have been planted and survived. The problem is compounded by the fact that many favorite windbreak species are not indigenous to the areas in which they are planted.

The Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups shows the adapted species listing for each group index number. Showing the height that locally grown trees and shrubs are expected to reach in 20 years on various soils. The estimates are based on measurements and observation of established plantings that have been given adequate care. This information should be used to determine the placement of a windbreak, the area protected and the arrangement of species.

A number of attributes are included in the CTSG species tables for each group number found in this section of the Field Office Technical Guide. These attributes were rated subjectively and assigned a relative value to further assist those unfamiliar with individual species characteristics or desirability for the intended use. Definitions and explanations can be found. Additional information on planning windbreaks and screens and planting and caring for trees and shrubs can be obtained from the local office of the Natural Resources Conservation Service or of the Cooperative Extension Service or from a commercial nursery. See part 537 of the National Forestry Manual for additional information.

In the Tree and Shrub Management table interpretive ratings are given for various aspects of forest and conservation tree and shrub management. Some rating class terms indicate the degree to which the soils are suited to a specified forest management practice. Well suited indicates that the soil has features that are favorable for the specified practice and has no limitations. Good performance can be expected, and little or no maintenance is needed. Moderately well suited indicates that the soil has features that are moderately favorable for the specified practice. One or more soil properties are less than desirable and fair performance can be expected. Some maintenance is needed. Poorly suited indicates that the soil has one or more properties that are unfavorable for the specified practice. Overcoming the unfavorable properties requires special design, extra maintenance, and costly alteration. Unsited indicates that the expected performance of the soil is unacceptable for the specified practice or that extreme measures are needed to overcome the undesirable soil properties.

The paragraphs that follow indicate the soil properties considered in rating the soils for forest and conservation tree and shrub management practices. More detailed information about the criteria used in the ratings is available in the "National Forestry Manual," which is available in local offices of the Natural Resources Conservation Service or on the Internet. Also, in the Kansas Field Office Technical Guide Notice KS-230, Conservation Tree and Shrub Plantings Suitability Groups.

Ratings in the columns suitability for hand planting and suitability for mechanical planting are based on slope, depth to a restrictive layer, content of sand, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, moderately well suited, poorly suited, or unsited to these methods of planting. It is assumed that necessary site preparation is completed before seedlings are planted.

Ratings in the column suitability for mechanical site preparation (surface) are based on slope, depth to a restrictive layer, plasticity index, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 1-foot is considered in the ratings.

Ratings in the column suitability for mechanical site preparation (deep) are based on slope, depth to a restrictive layer, rock fragments on or below the surface, depth to a water table, and ponding. The soils are described as well suited, poorly suited, or unsited to this management activity. The part of the soil from the surface to a depth of about 3 feet is considered in the ratings.

Ratings in the column potential for seedling mortality are based on flooding, ponding, depth to a water table, content of lime, reaction, salinity, available water capacity, soil moisture regime, soil temperature regime, aspect, and slope. The soils are described as having a low, moderate, or high potential for seedling mortality. See the National Forestry Manual, Subpart B for criteria used in rating management concerns. Specific information on plants and yields can be obtained from the local office of the Natural Resources Conservation Service or the Cooperative Extension Service.

CONSERVATION TREE AND SHRUB MANAGEMENT
Kearney County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Wind break Group | Suitability for hand planting | Suitability for mechanical planting | Suitability for mechanical site preparation (surface) | Suitability for mechanical site preparation (deep) | Potential for seedling mortality |
|--------------------------|------------------|------------------------------------|-------------------------------------|---|--|------------------------------------|
| | | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features |
| Ad: Alda----- | | Well suited | Well suited | Well suited | Well suited | Moderate Soil reaction |
| Bo: Boel----- | | Well suited | Well suited | Well suited | Well suited | Moderate Soil reaction |
| Bu: Butler----- | | Poorly suited Stickiness | Poorly suited Stickiness | Poorly suited Stickiness | Well suited | High Wetness |
| CaC: Coly----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Moderate Soil reaction |
| CaD: Coly----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Moderate Soil reaction |
| CaF: Coly----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Moderate Soil reaction |
| CkB: Coly----- | | Well suited | Well suited | Well suited | Well suited | Moderate Soil reaction Low |
| Kenesaw----- | | Well suited | Well suited | Well suited | Well suited | Moderate |
| CoD2: Coly----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Moderate Soil reaction Low |
| Uly----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Moderate Soil reaction Low |
| CoF2: Coly----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Moderate Soil reaction Low |
| Uly----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Moderate Soil reaction Low |
| De: Detroit----- | | Moderately suited Stickiness | Moderately suited Stickiness | Well suited | Well suited | Low |
| Fm: Fillmore----- | | Poorly suited Stickiness | Poorly suited Stickiness | Poorly suited Stickiness | Well suited | High Wetness |
| Gb: Gibbon----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Gc: Gibbon----- | | Well suited | Well suited | Well suited | Well suited | High Salinity |
| Go: Gothenburg----- | | Moderately suited Sandiness | Moderately suited Sandiness | Well suited | Well suited | High Wetness |
| GP: Pits----- | | Not rated | Not rated | Not rated | Not rated | Not rated |
| HeB: Hersh----- | | Well suited | Well suited | Well suited | Well suited | Low |
| HeC: Hersh----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Low |
| HeD: Hersh----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Low |
| Hf: Hobbs----- | | Well suited | Well suited | Well suited | Well suited | Low |
| HgB: Hobbs----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Ho: Holdrege----- | | Well suited | Well suited | Well suited | Well suited | Low |
| HoB: Holdrege----- | | Well suited | Well suited | Well suited | Well suited | Low |
| HoC: Holdrege----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Low |

CONSERVATION TREE AND SHRUB MANAGEMENT
Kearney County,
Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Wind break Group | Suitability for hand planting | Suitability for mechanical planting | Suitability for mechanical site preparation (surface) | Suitability for mechanical site preparation (deep) | Potential for seedling mortality |
|------------------------------|------------------|------------------------------------|---|---|--|------------------------------------|
| | | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features |
| HoC2: Holdrege----- | | Well suited | Moderately suited Slope | Well suited | Well suited | Low |
| Hp: Holdrege----- | | Well suited | Well suited | Well suited | Well suited | Low |
| HpB: Holdrege----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Hr: Hord----- | | Well suited | Well suited | Well suited | Well suited | Low |
| InB: Inavale----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Ke: Kenesaw----- | | Well suited | Well suited | Well suited | Well suited | Low |
| KgB: Kenesaw----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Coly----- | | Well suited | Well suited | Well suited | Well suited | Moderate Soil reaction |
| Lf: Lex----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Lg: Lex----- | | Well suited | Well suited | Well suited | Well suited | High Soil reaction Salinity |
| LOB: Libory----- | | Well suited | Well suited | Well suited | Well suited | Low |
| M-W: Miscellaneous Water- | | Not rated | Not rated | Not rated | Not rated | Not rated |
| Ma: Massie----- | | Poorly suited Stickiness | Poorly suited Stickiness | Poorly suited Stickiness | Well suited | High Wetness |
| Pm: Platte----- | | Well suited | Well suited | Well suited | Well suited | Low |
| Ru: Rusco----- | | Moderately suited Stickiness | Moderately suited Stickiness | Well suited | Well suited | High Wetness |
| Sc: Scott----- | | Poorly suited Stickiness | Poorly suited Stickiness | Poorly suited Stickiness | Well suited | High Wetness |
| SmB: Simeon----- | | Moderately suited Sandiness | Moderately suited Sandiness | Well suited | Well suited | Low |
| To: Tryon----- | | Well suited | Well suited | Well suited | Well suited | High Wetness |
| UcF: Uly----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Low |
| Coly----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Moderate Soil reaction |
| UsF: Ustorthents----- | | Well suited | Poorly suited Slope | Poorly suited Slope | Poorly suited Slope | Moderate Soil reaction |
| VaB: Valentine----- | | Moderately suited Sandiness | Moderately suited Sandiness | Well suited | Well suited | Low |
| VaD: Valentine----- | | Moderately suited Sandiness | Moderately suited Sandiness Slope | Well suited | Well suited | Low |
| VaF: Valentine----- | | Moderately suited Sandiness | Moderately suited Slope Sandiness | Well suited | Well suited | Low |
| VbD: Valentine----- | | Moderately suited Sandiness | Moderately suited Sandiness Slope | Well suited | Well suited | Low |
| Els----- | | Well suited | Well suited | Well suited | Well suited | Low |
| W: Water----- | | Not rated | Not rated | Not rated | Not rated | Not rated |

CONSERVATION TREE AND SHRUB MANAGEMENT
 Kearney County,
 Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. Pines and spruces are prone to disease problems. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Wind break Group | Suitability for hand planting | Suitability for mechanical planting | Suitability for mechanical site preparation (surface) | Suitability for mechanical site preparation (deep) | Potential for seedling mortality |
|--------------------------|------------------|------------------------------------|-------------------------------------|---|--|------------------------------------|
| | | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features | Rating class and limiting features |
| Wa: Wann----- | | Well suited | Well suited | Well suited | Well suited | Low |

ENGINEERING INDEX PROPERTIES
Kearney County, Nebraska

Engineering Index Properties table gives the engineering classifications and the range of index properties for the layers of each soil in the survey area. Depth to the upper and lower boundaries of each layer is indicated. Texture is given in the standard terms used by the U.S. Department of Agriculture. These terms are defined according to percentages of sand, silt, and clay in the fraction of the soil that is less than 2 millimeters in diameter. Loam, for example, is soil that is 7 to 27 percent clay, 28 to 50 percent silt, and less than 52 percent sand. If the content of particles coarser than sand is 15 percent or more, an appropriate modifier is added, for example, gravelly. Textural terms are defined in the Glossary.

Classification of the soils is determined according to the Unified soil classification system (ASTM, 1998) and the system adopted by the American Association of State Highway and Transportation Officials (AASHTO, 1998). The Unified system classifies soils according to properties that affect their use as construction material. Soils are classified according to particle-size distribution of the fraction less than 3 inches in diameter and according to plasticity index, liquid limit, and organic matter content. Sandy and gravelly soils are identified as GW, GP, GM, GC, SW, SP, SM, and SC; silty and clayey soils as ML, CL, OL, MH, CH, and OH; and highly organic soils as PT. Soils exhibiting engineering properties of two groups can have a dual classification, for example, CL-ML.

The AASHTO system classifies soils according to those properties that affect roadway construction and maintenance. In this system, the fraction of a mineral soil that is less than 3 inches in diameter is classified in one of seven groups from A-1 through A-7 on the basis of particle-size distribution, liquid limit, and plasticity index. Soils in group A-1 are coarse grained and low in content of fines (silt and clay). At the other extreme, soils in group A-7 are fine grained. Highly organic soils are classified in group A-8 on the basis of visual inspection. If laboratory data are available, the A-1, A-2, and A-7 groups are further classified as A-1-a, A-1-b, A-2-4, A-2-5, A-2-6, A-2-7, A-7-5, or A-7-6. As an additional refinement, the suitability of a soil as subgrade material can be indicated by a group index number. Group index numbers range from 0 for the best subgrade material to 20 or higher for the poorest. The AASHTO classification for soils tested, with group index numbers in parentheses, is given in Engineering Index Properties table.

Rock fragments larger than 10 inches in diameter and 3 to 10 inches in diameter are indicated as a percentage of the total soil on a dry-weight basis. The percentages are estimates determined mainly by converting volume percentage in the field to weight percentage. Percentage (of soil particles) passing designated sieves is the percentage of the soil fraction less than 3 inches in diameter based on an oven-dry weight. The sieves, numbers 4, 10, 40, and 200 (USA Standard Series), have openings of 4.76, 2.00, 0.420, and 0.074 millimeters, respectively. Estimates are based on laboratory tests of soils sampled in the survey area and in nearby areas and on estimates made in the field.

Liquid limit and plasticity index (Atterberg limits) indicate the plasticity characteristics of a soil. The estimates are based on test data from the survey area or from nearby areas and on field examination. The estimates of particle-size distribution, liquid limit, and plasticity index are generally rounded to the nearest 5 percent. Thus, if the ranges of gradation and Atterberg limits extend a marginal amount (1 or 2 percentage points) across classification boundaries, the classification in the marginal zone is generally omitted in the table.

ENGINEERING INDEX PROPERTIES--Continued
Kearney County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

| Map symbol and soil name | Depth In | USDA texture | Classification | | Fragments | | Percentage passing sieve number-- | | | | Liquid limit Pct | Plas- ticity index |
|-----------------------------|---------------------------------|---|--|--|------------------|------------------|--------------------------------------|---------------------------|--------------------------------------|--------------------------------------|----------------------------------|---------------------------------|
| | | | Unified | AASHTO | >10 inches | 3-10 inches | 4 | 10 | 40 | 200 | | |
| | | | | | Pct | Pct | | | | | | |
| Ad: Alda----- | 0-12 12-30 30-60 | Loam Fine sandy loam Coarse sand | ML, CL-ML, CL SC-SM, SM SC-SM, SM, SP, SP-SM | A-4 A-2, A-4 A-1, A-2, A-3 | 0 0 0 | 0 0 0 | 90-100 95-100 70-100 | 85-100 95-100 65-95 | 85-100 70-100 30-95 | 50-75 30-50 2-15 | 20-35 15-26 15-20 | 3-10 NP-7 NP-5 |
| Bo: Boel----- | 0-17 17-54 54-60 | Fine sandy loam Fine sand Loamy fine sand | SC-SM, SM SM, SC-SM, SP SP, SM, SC-SM | A-2, A-4 A-3, A-2 A-3, A-2 | 0 0 0 | 0 0 0 | 100 100 100 | 100 95-100 95-100 | 85-95 85-95 85-95 | 20-40 0-25 0-25 | 15-20 10-20 10-20 | NP-5 NP-5 NP-5 |
| Bu: Butler----- | 0-12 12-35 35-41 41-60 | Silt loam Silty clay Silty clay loam Silt loam | CL, CL-ML CH CH, CL CL | A-4, A-6 A-7 A-6, A-7 A-6, A-7 | 0 0 0 0 | 0 0 0 0 | 100 100 100 100 | 100 100 100 100 | 100 100 100 100 | 95-100 95-100 95-100 95-100 | 20-35 50-70 35-60 30-45 | 5-15 30-45 15-35 10-25 |
| CaC: Coly----- | 0-5 5-60 | Silt loam Silt loam | CL, CL-ML, ML CL, ML, CL-ML | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| CaD: Coly----- | 0-5 5-60 | Silt loam Silt loam | CL, CL-ML, ML ML, CL-ML, CL | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| CaF: Coly----- | 0-4 4-60 | Silt loam Silt loam | ML, CL-ML, CL CL-ML, CL, ML | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| CkB: Coly----- | 0-6 6-60 | Silt loam Silt loam | ML, CL-ML, CL ML, CL, CL-ML | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| Kenesaw----- | 0-10 10-23 23-60 | Silt loam Silt loam Silt loam Silt loam | CL, CL-ML CL, CL-ML ML, CL-ML, CL ML, CL-ML, CL | A-4, A-6 A-4, A-6 A-4, A-6 A-4, A-6 | 0 0 0 0 | 0 0 0 0 | 100 100 100 100 | 100 100 100 100 | 95-100 95-100 90-100 95-100 | 85-100 85-100 85-100 80-100 | 25-35 18-35 20-35 | 5-15 2-13 2-12 |
| CoD2: Coly----- | 0-5 5-60 | Silt loam Silt loam | ML, CL, CL-ML ML, CL-ML, CL | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| Uly----- | 0-7 7-28 28-60 | Silt loam Silt loam Silt loam | ML, CL ML, CL CL, ML | A-4, A-6 A-4, A-6 A-4, A-6 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 100 100 100 | 95-100 95-100 95-100 | 20-40 25-40 25-40 | 2-20 3-15 3-15 |
| CoF2: Coly----- | 0-6 6-60 | Silt loam Silt loam | ML, CL-ML, CL CL, CL-ML, ML | A-4, A-6, A-7 A-4, A-6 | 0 0 | 0 0 | 100 100 | 100 100 | 85-100 85-100 | 85-100 85-100 | 20-45 20-40 | 2-20 2-15 |
| Uly----- | 0-7 7-28 28-60 | Silt loam Silt loam Silt loam | CL, ML CL, ML CL, ML | A-4, A-6 A-4, A-6 A-4, A-6 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 100 100 100 | 95-100 95-100 95-100 | 20-40 25-40 25-40 | 2-20 3-15 3-15 |
| De: Detroit----- | 0-16 16-38 38-60 | Silt loam Silty clay loam Silty clay loam | CL CH CL | A-6 A-7 A-6, A-7 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 95-100 95-100 95-100 | 85-95 90-100 85-100 | 35-40 50-60 35-50 | 15-25 30-40 15-30 |
| Fm: Fillmore----- | 0-12 12-35 35-44 44-60 | Silt loam Silty clay Silty clay loam Silt loam | CL, CL-ML, ML CH CH, CL CL, CH | A-4, A-6 A-7 A-6, A-7 A-6, A-7 | 0 0 0 0 | 0 0 0 0 | 100 100 100 100 | 100 100 100 100 | 95-100 100 100 100 | 95-100 95-100 95-100 95-100 | 20-40 50-75 35-60 25-75 | 2-20 30-45 20-40 10-45 |
| Gb: Gibbon----- | 0-14 14-37 37-60 | Loam Silt loam Stratified very fine sandy loam to silt loam | CL, CL-ML CL CL, ML, SC, SM | A-4, A-6 A-6 A-4 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 85-95 90-100 70-95 | 60-75 80-90 35-90 | 25-40 25-38 15-25 | 5-20 12-20 NP-8 |
| Gc: Gibbon----- | 0-10 10-40 40-60 | Loam Silt loam Silty clay loam | CL CL CL | A-4, A-6 A-6, A-7 A-7 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 90-100 95-100 95-100 | 70-90 80-95 85-95 | 25-35 30-45 40-50 | 8-15 10-20 20-28 |
| Go: Gothenburg----- | 0-3 3-8 8-60 | Loamy sand Sand Stratified coarse sand to gravelly coarse sand | SM SM, SP, SP-SM SM, SP, SP-SM | A-2 A-2, A-3 A-1, A-2, A-3 | 0 0 0 | 0 0 0 | 100 100 70-100 | 95-100 80-100 50-95 | 50-90 65-80 25-65 | 15-35 3-15 0-15 | 0-20 0-20 0-20 | NP NP NP |
| GP: Pits----- | 0-60 | Gravelly sand | SP-SM, SP, SM, GP-GM | A-1, A-2, A-3 | --- | 0-5 | 45-100 | 40-100 | 0-80 | 0-40 | 0-14 | NP |
| HeB: Hersh----- | 0-7 7-11 11-60 | Fine sandy loam Fine sandy loam Fine sandy loam | SC-SM, SM, ML, SC SM, SC-SM, CL-ML, ML ML, SM | A-4 A-4 A-4 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 70-85 80-100 90-100 | 40-55 40-65 40-65 | 15-25 15-20 15-20 | NP-10 NP-5 NP-5 |
| HeC: Hersh----- | 0-7 7-11 11-60 | Fine sandy loam Fine sandy loam Fine sandy loam | ML, SC, SC- SM, SM CL-ML, ML, SC-SM, SM SM, ML | A-4 A-4 A-4 | 0 0 0 | 0 0 0 | 100 100 100 | 100 100 100 | 70-85 80-100 90-100 | 40-55 40-65 40-65 | 15-25 15-20 15-20 | NP-10 NP-5 NP-5 |

ENGINEERING INDEX PROPERTIES--Continued
Kearney County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

| Map symbol and soil name | Depth | USDA texture | Classification | | Fragments | | Percentage passing sieve number-- | | | | Liquid limit | Plasticity index |
|----------------------------------|-------|--|----------------------|---------------|------------|-------------|-----------------------------------|--------|--------|--------|--------------|------------------|
| | | | Unified | AASHTO | >10 inches | 3-10 inches | 4 | 10 | 40 | 200 | | |
| | | | | | Pct | Pct | | | | | | |
| HeD: Hersh----- | 0-7 | Fine sandy loam | ML, SC, SC-SM, SM | A-4 | 0 | 0 | 100 | 100 | 70-85 | 40-55 | 15-25 | NP-10 |
| | 7-11 | Fine sandy loam | SM, CL-ML, SC-SM, ML | A-4 | 0 | 0 | 100 | 100 | 80-100 | 40-65 | 15-20 | NP-5 |
| Hf: Hobbs----- | 11-60 | Fine sandy loam | ML, SM | A-4 | 0 | 0 | 100 | 100 | 90-100 | 40-65 | 15-20 | NP-5 |
| | 0-6 | Silt loam | CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-40 | 5-20 |
| | 6-34 | Stratified silt loam | CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-40 | 5-20 |
| HgB: Hobbs----- | 34-60 | Silt loam | CL-ML, MH, CL | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 80-100 | 25-55 | 5-25 |
| | 0-9 | Silt loam | CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-40 | 5-20 |
| | 9-34 | Stratified silt loam | CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-40 | 5-20 |
| Ho: Holdrege----- | 34-60 | Silt loam | CL, CL-ML, MH | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 80-100 | 25-55 | 5-25 |
| | 0-13 | Silt loam | CL, CL-ML, ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 13-25 | Silty clay loam | CH, CL | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 25-31 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 31-60 | Silt loam | ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| HoB: Holdrege----- | 0-13 | Silt loam | ML, CL, CL-ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 13-25 | Silty clay loam | CH, CL | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 25-31 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 31-60 | Silt loam | CL, ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| HoC: Holdrege----- | 0-13 | Silt loam | CL-ML, CL, ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 13-25 | Silty clay loam | CL, CH | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 25-31 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 31-60 | Silt loam | CL, ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| HoC2: Holdrege----- | 0-13 | Silt loam | ML, CL-ML, CL | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 13-25 | Silty clay loam | CH, CL | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 25-31 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 31-60 | Silt loam | ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| Hp: Holdrege----- | 0-27 | Silt loam | CL-ML, CL, ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 27-43 | Silty clay loam | CH, CL | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 43-47 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 47-60 | Silt loam | CL, ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| HpB: Holdrege----- | 0-27 | Silt loam | ML, CL-ML, CL | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-45 | 2-20 |
| | 27-43 | Silty clay loam | CL, CH | A-6, A-7 | 0 | 0 | 100 | 100 | 98-100 | 90-100 | 30-55 | 15-35 |
| | 43-47 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 95-100 | 25-40 | 9-17 |
| | 47-60 | Silt loam | ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 90-100 | 30-40 | 5-15 |
| Hr: Hord----- | 0-13 | Silt loam | CL-ML, ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 20-35 | 3-18 |
| | 13-43 | Silt loam | CL | A-4, A-6 | 0 | 0 | 100 | 100 | 98-100 | 85-100 | 25-40 | 8-23 |
| | 43-60 | Silt loam | CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 100 | 100 | 85-100 | 25-40 | 6-21 |
| InB: Inavale----- | 0-5 | Loamy fine sand | SM, SC-SM, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 85-95 | 5-35 | 15-25 | NP-5 |
| | 5-10 | Loamy fine sand | SP-SM, SM, SC-SM | A-2, A-3 | 0 | 0 | 100 | 90-100 | 65-85 | 5-30 | 15-25 | NP-5 |
| | 10-60 | Fine sand | SP-SM, SM, SC-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 70-90 | 5-30 | 15-25 | NP-5 |
| Ke: Kenesaw----- | 0-8 | Silt loam | CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-35 | 5-15 |
| | 8-22 | Silt loam | ML, CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 90-100 | 85-100 | 18-35 | 2-13 |
| | 22-60 | Silt loam | ML, CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 80-100 | 20-35 | 2-12 |
| KgB: Kenesaw----- | 0-7 | Silt loam | CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 100 | 95-100 | 85-100 | 25-35 | 5-15 |
| | 7-18 | Silt loam | ML, CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 100 | 90-100 | 85-100 | 18-35 | 2-13 |
| | 18-60 | Silt loam | CL, CL-ML, ML | A-6, A-4 | 0 | 0 | 100 | 100 | 95-100 | 80-100 | 20-35 | 2-12 |
| Coly----- | 0-5 | Silt loam | CL, CL-ML, ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 85-100 | 85-100 | 20-45 | 2-20 |
| | 5-60 | Silt loam | ML, CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 85-100 | 85-100 | 20-45 | 2-15 |
| Lf: Lex----- | 0-11 | Loam | CL, CL-ML, ML | A-4, A-6 | 0 | 0 | 95-100 | 95-100 | 85-100 | 60-95 | 22-35 | 3-15 |
| | 11-24 | Stratified sandy loam to silty clay loam | CL, CL-ML, ML | A-4, A-6, A-7 | 0 | 0 | 95-100 | 95-100 | 85-100 | 60-90 | 20-45 | 3-25 |
| Lg: Lex----- | 24-60 | Coarse sand | SM, SP-SM, SP | A-1, A-2, A-3 | 0 | 0 | 60-100 | 50-95 | 30-65 | 3-14 | 15-20 | NP-5 |
| | 0-12 | Loam | CL, CL-ML, ML | A-4, A-6 | 0 | 0 | 95-100 | 95-100 | 85-100 | 60-95 | 20-35 | 3-15 |
| | 12-31 | Silt loam | CL | A-6 | 0 | 0 | 95-100 | 95-100 | 90-100 | 70-85 | 30-40 | 12-22 |
| | 31-60 | Coarse sand | SM, SP, SP-SM | A-1, A-2, A-3 | 0 | 0 | 80-100 | 70-100 | 35-70 | 3-15 | 10-20 | NP-5 |
| LoB: Libory----- | 0-16 | Loamy fine sand | SM | A-2, A-4 | 0 | 0 | 100 | 100 | 65-85 | 15-45 | 10-20 | NP-5 |
| | 16-24 | Loamy fine sand | SP-SM, SM | A-2 | 0 | 0 | 100 | 100 | 55-80 | 12-35 | 10-20 | NP-5 |
| | 24-60 | Silt loam | CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 85-100 | 60-95 | 20-40 | 4-24 |
| M-W: Miscellaneous Water----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Ma: Massie----- | 0-10 | Silty clay loam | CL | A-6, A-7 | 0 | 0 | 100 | 100 | 100 | 95-100 | 30-45 | 10-25 |
| | 10-60 | Silty clay | CL, CH | A-7 | 0 | 0 | 100 | 100 | 100 | 95-100 | 45-70 | 20-45 |

ENGINEERING INDEX PROPERTIES--Continued
Kearney County, Nebraska

(Absence of an entry indicates that the data were not estimated.)

| Map symbol and soil name | Depth | USDA texture | Classification | | Fragments | | Percentage passing sieve number-- | | | | Liquid limit | Plas- ticity index |
|-----------------------------|-------|--|-------------------------|---------------|-----------|--------|--------------------------------------|--------|--------|--------|-----------------|--------------------------|
| | | | Unified | AASHTO | >10 | 3-10 | 4 | 10 | 40 | 200 | | |
| | | | | | inches | inches | | | | | | |
| | In | | | | Pct | Pct | | | | | Pct | |
| Pm: Platte----- | 0-6 | Loam | CL-ML, CL | A-4, A-6 | 0 | 0 | 100 | 95-100 | 85-100 | 60-95 | 22-35 | 4-15 |
| | 6-12 | Loam | ML, CL-ML, SM, SC-SM | A-4 | 0 | 0 | 100 | 95-100 | 75-95 | 45-75 | 10-20 | NP-5 |
| | 12-60 | Gravelly coarse sand | SM, SP-SM | A-1, A-2, A-3 | 0 | 0 | 70-100 | 50-95 | 25-65 | 0-15 | 10-20 | NP-5 |
| Ru: Rusco----- | 0-5 | Silt loam | CL-ML, ML, CL | A-4, A-6 | 0 | 0 | 100 | 95-100 | 85-100 | 60-100 | 20-40 | 3-15 |
| | 5-30 | Silty clay loam | CL | A-6, A-7 | 0 | 0 | 100 | 100 | 90-100 | 85-100 | 30-45 | 15-30 |
| | 30-60 | Silt loam | ML, CL-ML, CL | A-4, A-6 | 0 | 0 | 95-100 | 95-100 | 90-100 | 70-100 | 20-35 | 3-15 |
| Sc: Scott----- | 0-6 | Silt loam | CL, CL-ML, ML | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 100 | 95-100 | 20-45 | 2-20 |
| | 6-37 | Clay | CL, CH | A-7 | 0 | 0 | 100 | 100 | 100 | 95-100 | 41-75 | 20-45 |
| | 37-48 | Silty clay loam | CH, CL | A-6, A-7 | 0 | 0 | 100 | 100 | 100 | 95-100 | 35-60 | 20-40 |
| | 48-60 | Silt loam | CL | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 90-100 | 90-100 | 25-50 | 8-24 |
| SmB: Simeon----- | 0-6 | Sandy loam | SM | A-2, A-4 | 0 | 0 | 95-100 | 90-100 | 60-85 | 30-40 | 10-20 | NP-5 |
| | 6-60 | Coarse sand | SP-SM, SP, SM | A-1, A-2, A-3 | 0 | 0 | 90-100 | 80-100 | 35-95 | 0-30 | 10-20 | NP-5 |
| To: Tryon----- | 0-4 | Loamy fine sand | SP-SM, SM | A-2 | 0 | 0 | 100 | 100 | 85-100 | 10-30 | 10-20 | NP-5 |
| | 4-60 | Fine sand | SP-SM, SP, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 51-95 | 3-30 | 5-15 | NP-5 |
| UcF: Uly----- | 0-9 | Silt loam | ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 100 | 95-100 | 20-40 | 2-20 |
| | 9-27 | Silt loam | ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 100 | 95-100 | 25-40 | 3-15 |
| | 27-60 | Silt loam | CL, ML | A-4, A-6 | 0 | 0 | 100 | 100 | 100 | 95-100 | 25-40 | 3-15 |
| Coly----- | 0-4 | Silt loam | ML, CL-ML, CL | A-4, A-6, A-7 | 0 | 0 | 100 | 100 | 85-100 | 85-100 | 20-45 | 2-20 |
| | 4-60 | Silt loam | CL-ML, ML, CL | A-4, A-6 | 0 | 0 | 100 | 100 | 85-100 | 85-100 | 20-40 | 2-15 |
| UsF: Ustorthents---- | 0-60 | Silt loam | ML, CL, CL-ML | A-4, A-6 | 0 | 0 | 100 | 90-100 | 80-100 | 55-95 | 25-35 | 5-15 |
| VaB: Valentine----- | 0-5 | Loamy fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 95-100 | 2-35 | 15-20 | NP-5 |
| | 5-8 | Loamy fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 90-100 | 2-35 | 15-20 | NP-5 |
| | 8-60 | Fine sand | SP-SM, SP, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 70-100 | 2-25 | 15-20 | NP-5 |
| VaD: Valentine----- | 0-5 | Loamy fine sand | SP, SP-SM, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 95-100 | 2-35 | 15-20 | NP-5 |
| | 5-8 | Loamy fine sand | SP, SP-SM, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 90-100 | 2-35 | 15-20 | NP-5 |
| | 8-60 | Fine sand | SM, SP-SM, SP | A-2, A-3 | 0 | 0 | 100 | 100 | 70-100 | 2-25 | 15-20 | NP-5 |
| VaF: Valentine----- | 0-5 | Loamy fine sand | SP, SP-SM, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 95-100 | 2-35 | 15-20 | NP-5 |
| | 5-8 | Loamy fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 90-100 | 2-35 | 15-20 | NP-5 |
| | 8-60 | Fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 70-100 | 2-25 | 15-20 | NP-5 |
| VbD: Valentine----- | 0-5 | Loamy fine sand | SP-SM, SP, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 95-100 | 2-35 | 15-20 | NP-5 |
| | 5-8 | Loamy fine sand | SP-SM, SP, SM | A-2, A-3 | 0 | 0 | 100 | 100 | 90-100 | 2-35 | 15-20 | NP-5 |
| | 8-60 | Fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 100 | 100 | 70-100 | 2-25 | 15-20 | NP-5 |
| Els----- | 0-6 | Loamy fine sand | SM | A-2 | 0 | 0 | 95-100 | 85-100 | 60-100 | 25-35 | --- | NP |
| | 6-10 | Loamy sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 90-100 | 90-100 | 70-100 | 4-30 | --- | NP |
| | 10-60 | Fine sand | SM, SP, SP-SM | A-2, A-3 | 0 | 0 | 90-100 | 90-100 | 70-100 | 4-30 | --- | NP |
| W: Water----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Wa: Wann----- | 0-14 | Fine sandy loam | SM, SC-SM | A-2, A-4 | 0 | 0 | 95-100 | 95-100 | 70-100 | 30-50 | 15-25 | NP-5 |
| | 14-48 | Fine sandy loam | SC-SM, SM | A-2, A-4 | 0 | 0 | 95-100 | 75-100 | 60-100 | 20-50 | 15-25 | NP-5 |
| | 48-60 | Stratified sandy clay loam to fine sand | SM | A-2, A-4 | 0 | 0 | 95-100 | 95-100 | 70-100 | 15-40 | 15-20 | NP-3 |

Physical Properties

This table shows estimates of some physical characteristics and features that affect soil behavior. These estimates are given for the layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils.

Depth to the upper and lower boundaries of each layer is indicated.

Particle size is the effective diameter of a soil particle as measured by sedimentation, sieving, or micrometric methods. Particle sizes are expressed as classes with specific effective diameter class limits. The broad classes are sand, silt, and clay, ranging from the larger to the smaller.

Sand as a soil separate consists of mineral soil particles that are 0.05 millimeter to 2 millimeters in diameter. In this table, the estimated sand content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Silt as a soil separate consists of mineral soil particles that are 0.002 to 0.05 millimeter in diameter. In this table, the estimated silt content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

Clay as a soil separate consists of mineral soil particles that are less than 0.002 millimeter in diameter. In this table, the estimated clay content of each soil layer is given as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of sand, silt, and clay affects the physical behavior of a soil. Particle size is important for engineering and agronomic interpretations, for determination of soil hydrologic qualities, and for soil classification.

The amount and kind of clay affect the fertility and physical condition of the soil and the ability of the soil to adsorb cations and to retain moisture. They influence shrink-swell potential, permeability, plasticity, the ease of soil dispersion, and other soil properties. The amount and kind of clay in a soil also affect tillage and earthmoving operations.

Moist bulk density is the weight of soil (oven-dry) per unit volume. Volume is measured when the soil is at field moisture capacity, that is, the moisture content at 1/3- or 1/10-bar (33kPa or 10kPa) moisture tension. Weight is determined after the soil is dried at 105 degrees C. In the table, the estimated moist bulk density of each soil horizon is expressed in grams per cubic centimeter of soil material that is less than 2 millimeters in diameter. Bulk density data are used to compute shrink-swell potential, available water capacity, total pore space, and other soil properties. The moist bulk density of a soil indicates the pore space available for water and roots. Depending on soil texture, a bulk density of more than 1.4 can restrict water storage and root penetration. Moist bulk density is influenced by texture, kind of clay, content of organic matter, and soil structure.

Saturated hydraulic conductivity refers to the ability of a soil to transmit water or air. The term "permeability," as used in soil surveys, indicates saturated hydraulic conductivity (K_{sat}). The estimates in the table indicate the rate of water movement, in micrometers per second (um/sec), when the soil is saturated. They are based on soil characteristics observed in the field, particularly structure, porosity, and texture. Permeability is considered in the design of soil drainage systems and septic tank absorption fields.

Available water capacity refers to the quantity of water that the soil is capable of storing for use by plants. The capacity for water storage is given in inches of water per inch of soil for each soil layer. The capacity varies, depending on soil properties that affect retention of water. The most important properties are the content of organic matter, soil texture, bulk density, and soil structure. Available water capacity is an important factor in the choice of plants or crops to be grown and in the design and management of irrigation systems. Available water capacity is not an estimate of the quantity of water actually available to plants at any given time.

Linear extensibility refers to the change in length of an unconfined clod as moisture content is decreased from a moist to a dry state. It is an expression of the volume change between the water content of the clod at 1/3- or 1/10-bar tension (33kPa or 10kPa tension) and oven dryness. The volume change is reported in the table as percent change for the whole soil. Volume change is influenced by the amount and type of clay minerals in the soil.

Linear extensibility is used to determine the shrink-swell potential of soils. The shrink-swell potential is low if the soil has a linear extensibility of less than 3 percent; moderate if 3 to 6 percent; high if 6 to 9 percent; and very high if more than 9 percent. If the linear extensibility is more than 3, shrinking and swelling can cause damage to buildings, roads, and other structures and to plant roots. Special design commonly is needed.

Organic matter is the plant and animal residue in the soil at various stages of decomposition. In this table, the estimated content of organic matter is expressed as a percentage, by weight, of the soil material that is less than 2 millimeters in diameter.

The content of organic matter in a soil can be maintained by returning crop residue to the soil. Organic matter has a positive effect on available water capacity, water infiltration, soil organism activity, and tilth. It is a source of nitrogen and other nutrients for crops and soil organisms.

Erosion factors are shown in this table as the K factor (K_w and K_f) and the T factor. Erosion factor K indicates the susceptibility of a soil to sheet and rill erosion by water. Factor K is one of six factors used in the Universal Soil Loss Equation (USLE) and the Revised Universal Soil Loss Equation (RUSLE) to predict the average annual rate of soil loss by sheet and rill erosion in tons per acre per year. The estimates are based primarily on percentage of silt, sand, and organic matter and on soil structure and permeability. Values of K range from 0.02 to 0.69. Other factors being equal, the higher the value, the more susceptible the soil is to sheet and rill erosion by water.

Erosion factor K_w indicates the erodibility of the whole soil. The estimates are modified by the presence of rock fragments.

Erosion factor K_f indicates the erodibility of the fine-earth fraction, or the material less than 2 millimeters in size.

Erosion factor T is an estimate of the maximum average annual rate of soil erosion by wind or water that can occur without affecting crop productivity over a sustained period. The rate is in tons per acre per year.

Wind erodibility groups are made up of soils that have similar properties affecting their susceptibility to wind erosion in cultivated areas. The soils assigned to group 1 are the most susceptible to wind erosion, and those assigned to group 8 are the least susceptible. The groups are as follows:

1. Coarse sands, sands, fine sands, and very fine sands.

PHYSICAL PROPERTIES OF THE SOILS
Kearney County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

| Map symbol and soil name | Depth In | Sand Pct | Silt Pct | Clay Pct | Moist bulk density g/cc | Permeability (Ksat) in/hr | Available water capacity In/in | Linear extensibility Pct | Organic matter Pct | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|-------------|-------------|-------------|-------------|----------------------------|------------------------------|-----------------------------------|-----------------------------|-----------------------|-----------------|----|---|------------------------|------------------------|
| | | | | | | | | | | K | KF | T | | |

2. Loamy coarse sands, loamy sands, loamy fine sands, loamy very fine sands, ash material, and sapric soil material.
3. Coarse sandy loams, sandy loams, fine sandy loams, and very fine sandy loams.
- 4L. Calcareous loams, silt loams, clay loams, and silty clay loams.
4. Clays, silty clays, noncalcareous clay loams, and silty clay loams that are more than 35 percent clay.
5. Noncalcareous loams and silt loams that are less than 20 percent clay and sandy clay loams, sandy clays, and hemic soil material.
6. Noncalcareous loams and silt loams that are more than 20 percent clay and noncalcareous clay loams that are less than 35 percent clay.
7. Silts, noncalcareous silty clay loams that are less than 35 percent clay, and fibric soil material.
8. Soils that are not subject to wind erosion because of coarse fragments on the surface or because of surface wetness.

Wind erodibility index is a numerical value indicating the susceptibility of soil to wind erosion, or the tons per acre per year that can be expected to be lost to wind erosion. There is a close correlation between wind erosion and the texture of the surface layer, the size and durability of surface clods, rock fragments, organic matter, and a calcareous reaction. Soil moisture and frozen soil layers also influence wind erosion.

PHYSICAL PROPERTIES OF THE SOILS
Kearney County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

| Map symbol and soil name | Depth | Sand | Silt | Clay | Moist bulk density | Permeability (Ksat) | Available water capacity | Linear extensibility | Organic matter | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|-------|------|------|-------|--------------------|---------------------|--------------------------|----------------------|----------------|-----------------|-----|---|------------------------|------------------------|
| | | | | | | | | | | K | Kf | T | | |
| | In | Pct | Pct | Pct | g/cc | in/hr | In/in | Pct | Pct | | | | | |
| Ad: | | | | | | | | | | | | | | |
| Alda----- | 0-12 | 43 | 38 | 12-25 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 0.0-2.9 | 2.0-4.0 | .28 | .28 | 4 | 5 | 56 |
| | 12-30 | 66 | 28 | 3-10 | 1.40-1.60 | 2.00-6.00 | 0.15-0.17 | 0.0-2.9 | 0.5-1.0 | .20 | .20 | | | |
| | 30-60 | 92 | 6 | 0-2 | 1.50-1.70 | 19.98-19.98 | 0.02-0.04 | 0.0-2.9 | 0.0-0.5 | .10 | .15 | | | |
| Bo: | | | | | | | | | | | | | | |
| Boel----- | 0-17 | 67 | 20 | 8-18 | 1.50-1.70 | 2.00-6.00 | 0.16-0.18 | 0.0-2.9 | 1.0-2.0 | .20 | .20 | 3 | 3 | 86 |
| | 17-54 | 96 | 1 | 0-6 | 1.50-1.60 | 5.95-19.98 | 0.05-0.10 | 0.0-2.9 | 0.0-0.5 | .20 | .20 | | | |
| | 54-60 | 96 | 1 | 0-6 | 1.50-1.60 | 5.95-19.98 | 0.05-0.10 | 0.0-2.9 | 0.0-0.5 | .20 | .20 | | | |
| Bu: | | | | | | | | | | | | | | |
| Butler----- | 0-12 | 25 | 53 | 18-27 | 1.20-1.40 | 0.60-2.00 | 0.22-0.24 | 3.0-5.9 | 2.0-4.0 | .37 | .37 | 3 | 6 | 48 |
| | 12-35 | 5 | 45 | 45-55 | 1.10-1.20 | 0.01-0.06 | 0.11-0.13 | 9.0-12.0 | 1.0-2.0 | .37 | .37 | | | |
| | 35-41 | 7 | 54 | 32-45 | 1.10-1.30 | 0.20-0.60 | 0.14-0.20 | 6.0-8.9 | 0.5-1.0 | .37 | .37 | | | |
| | 41-60 | 23 | 50 | 20-35 | 1.20-1.40 | 0.60-2.00 | 0.18-0.22 | 3.0-5.9 | 0.0-0.5 | .37 | .37 | | | |
| CaC: | | | | | | | | | | | | | | |
| Coly----- | 0-5 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 5-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| CaD: | | | | | | | | | | | | | | |
| Coly----- | 0-5 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 5-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| CaF: | | | | | | | | | | | | | | |
| Coly----- | 0-4 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 4-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| ChB: | | | | | | | | | | | | | | |
| Coly----- | 0-6 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 6-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| Kenesaw----- | 0-10 | 11 | 67 | 18-25 | 1.20-1.40 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 10-23 | 14 | 71 | 10-20 | 1.20-1.30 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 23-60 | 14 | 73 | 8-18 | 1.30-1.40 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| CoD2: | | | | | | | | | | | | | | |
| Coly----- | 0-5 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 5-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| Uly----- | 0-7 | 11 | 67 | 17-27 | 1.20-1.45 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 7-28 | 9 | 66 | 20-30 | 1.25-1.45 | 0.60-2.00 | 0.18-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 28-60 | 10 | 68 | 18-27 | 1.25-1.45 | 0.60-2.00 | 0.16-0.20 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| CoF2: | | | | | | | | | | | | | | |
| Coly----- | 0-6 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 6-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| Uly----- | 0-7 | 11 | 67 | 17-27 | 1.20-1.45 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 7-28 | 9 | 66 | 20-30 | 1.25-1.45 | 0.60-2.00 | 0.18-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 28-60 | 10 | 68 | 18-27 | 1.25-1.45 | 0.60-2.00 | 0.16-0.20 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| De: | | | | | | | | | | | | | | |
| Detroit----- | 0-16 | 24 | 51 | 22-27 | 1.25-1.40 | 0.60-2.00 | 0.22-0.24 | 0.0-2.9 | 2.0-4.0 | .37 | .37 | 5 | 6 | 48 |
| | 16-38 | 7 | 53 | 35-45 | 1.35-1.50 | 0.06-0.20 | 0.12-0.18 | 6.0-8.9 | 1.0-2.0 | .37 | .37 | | | |
| | 38-60 | 20 | 54 | 18-35 | 1.30-1.50 | 0.20-0.60 | 0.18-0.22 | 3.0-5.9 | 0.0-0.5 | .37 | .37 | | | |
| Fm: | | | | | | | | | | | | | | |
| Fillmore----- | 0-12 | 25 | 53 | 18-27 | 1.30-1.40 | 0.60-2.00 | 0.21-0.24 | 0.0-2.9 | 2.0-4.0 | .37 | .37 | 3 | 6 | 48 |
| | 12-35 | 5 | 45 | 45-55 | 1.10-1.30 | 0.00-0.06 | 0.11-0.14 | 9.0-12.0 | 1.0-2.0 | .37 | .37 | | | |
| | 35-44 | 4 | 60 | 30-50 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 6.0-8.9 | 0.5-1.0 | .37 | .37 | | | |
| | 44-60 | 18 | 55 | 20-45 | 1.30-1.50 | 0.20-2.00 | 0.10-0.22 | 1.0-8.9 | 0.0-0.5 | .43 | .43 | | | |
| Gb: | | | | | | | | | | | | | | |
| Gibbon----- | 0-14 | 42 | 42 | 8-25 | 1.50-1.60 | 0.60-2.00 | 0.21-0.23 | 0.0-2.9 | 2.0-4.0 | .28 | .28 | 5 | 4L | 86 |
| | 14-37 | 9 | 67 | 20-27 | 1.30-1.50 | 0.60-2.00 | 0.18-0.22 | 0.0-3.0 | 0.5-1.0 | .32 | .32 | | | |
| | 37-60 | | | 15-25 | 1.50-1.70 | 0.57-5.95 | 0.16-0.20 | 0.0-2.9 | 0.5-1.0 | .32 | .32 | | | |
| Gc: | | | | | | | | | | | | | | |
| Gibbon----- | 0-10 | 37 | 43 | 15-25 | 1.50-1.60 | 0.60-2.00 | 0.20-0.22 | 0.0-2.9 | 1.0-3.0 | .28 | .28 | 5 | 4L | 86 |
| | 10-40 | 9 | 65 | 20-32 | 1.20-1.30 | 0.60-2.00 | 0.22-0.24 | 1.0-5.9 | 0.5-1.0 | .43 | .43 | | | |
| | 40-60 | 8 | 56 | 30-43 | 1.35-1.50 | 0.20-0.60 | 0.12-0.18 | 3.0-9.0 | 0.0-0.5 | .43 | .43 | | | |
| Go: | | | | | | | | | | | | | | |
| Gothenburg--- | 0-3 | 79 | 16 | 2-8 | 1.50-1.60 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 8 | 0 |
| | 3-8 | 96 | 1 | 1-5 | 1.50-1.70 | 5.95-19.98 | 0.06-0.08 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| | 8-60 | | | 0-3 | 1.55-1.70 | 19.98-19.98 | 0.02-0.04 | 0.0-2.9 | 0.0-0.5 | .05 | .10 | | | |
| GP: | | | | | | | | | | | | | | |
| Pits----- | 0-60 | 95 | 1 | 0-8 | 1.70-2.00 | 6.00-20.00 | 0.02-0.09 | 0.0-2.9 | 0.0-0.5 | .10 | .17 | 2 | 8 | 0 |
| HeB: | | | | | | | | | | | | | | |
| Hersh----- | 0-7 | 66 | 20 | 10-18 | 1.30-1.50 | 2.00-6.00 | 0.16-0.18 | 0.0-2.9 | 0.5-2.0 | .24 | .24 | 5 | 3 | 86 |
| | 7-11 | 67 | 20 | 8-18 | 1.30-1.50 | 2.00-6.00 | 0.15-0.18 | 0.0-2.9 | 0.5-1.0 | .24 | .24 | | | |
| | 11-60 | 66 | 20 | 10-18 | 1.20-1.50 | 2.00-6.00 | 0.15-0.17 | 0.0-2.9 | 0.0-0.5 | .24 | .24 | | | |
| HeC: | | | | | | | | | | | | | | |
| Hersh----- | 0-7 | 66 | 20 | 10-18 | 1.30-1.50 | 2.00-6.00 | 0.16-0.18 | 0.0-2.9 | 0.5-2.0 | .24 | .24 | 5 | 3 | 86 |
| | 7-11 | 67 | 20 | 8-18 | 1.30-1.50 | 2.00-6.00 | 0.15-0.18 | 0.0-2.9 | 0.5-1.0 | .24 | .24 | | | |
| | 11-60 | 66 | 20 | 10-18 | 1.20-1.50 | 2.00-6.00 | 0.15-0.17 | 0.0-2.9 | 0.0-0.5 | .24 | .24 | | | |
| HeD: | | | | | | | | | | | | | | |
| Hersh----- | 0-7 | 66 | 20 | 10-18 | 1.30-1.50 | 2.00-6.00 | 0.16-0.18 | 0.0-2.9 | 0.5-2.0 | .24 | .24 | 5 | 3 | 86 |
| | 7-11 | 67 | 20 | 8-18 | 1.30-1.50 | 2.00-6.00 | 0.15-0.18 | 0.0-2.9 | 0.5-1.0 | .24 | .24 | | | |
| | 11-60 | 66 | 20 | 10-18 | 1.20-1.50 | 2.00-6.00 | 0.15-0.17 | 0.0-2.9 | 0.0-0.5 | .24 | .24 | | | |
| Hf: | | | | | | | | | | | | | | |
| Hobbs----- | 0-6 | 11 | 68 | 15-27 | 1.20-1.40 | 0.60-2.00 | 0.21-0.24 | 0.0-2.9 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 6-34 | | | 15-27 | 1.20-1.40 | 0.60-2.00 | 0.18-0.20 | 0.0-2.9 | 0.5-1.0 | .32 | .32 | | | |
| | 34-60 | 10 | 68 | 15-30 | 1.20-1.40 | 0.60-2.00 | 0.18-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| HgB: | | | | | | | | | | | | | | |
| Hobbs----- | 0-9 | 11 | 68 | 15-27 | 1.20-1.40 | 0.60-2.00 | 0.21-0.24 | 0.0-2.9 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 9-34 | | | 15-27 | 1.20-1.40 | 0.60-2.00 | 0.18-0.20 | 0.0-2.9 | 0.5-1.0 | .32 | .32 | | | |
| | 34-60 | 10 | 68 | 15-30 | 1.20-1.40 | 0.60-2.00 | 0.18-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |

PHYSICAL PROPERTIES OF THE SOILS
Kearney County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

| Map symbol and soil name | Depth In | Sand Pct | Silt Pct | Clay Pct | Moist bulk density g/cc | Permea- bility (Ksat) in/hr | Available water capacity In/in | Linear extensi- bility Pct | Organic matter Pct | Erosion factors | | | Wind erodi- bility group | Wind erodi- bility index |
|-------------------------------------|-------------|-------------|-------------|-------------|----------------------------------|--------------------------------------|---|-------------------------------------|--------------------------|-----------------|-----|-----|-----------------------------------|-----------------------------------|
| | | | | | | | | | | K | Kf | T | | |
| Ho: Holdrege----- | 0-13 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 13-25 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 25-31 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 31-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| HoB: Holdrege----- | 0-13 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 13-25 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 25-31 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 31-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| HoC: Holdrege----- | 0-13 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 13-25 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 25-31 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 31-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| HoC2: Holdrege----- | 0-13 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 13-25 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 25-31 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 31-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| Hp: Holdrege----- | 0-27 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 27-43 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 43-47 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 47-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| HpB: Holdrege----- | 0-27 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.22-0.24 | 1.0-3.0 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 27-43 | 7 | 62 | 28-35 | 1.20-1.40 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-2.0 | .43 | .43 | | | |
| | 43-47 | 9 | 66 | 20-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.20 | 1.0-4.0 | 0.0-0.5 | .43 | .43 | | | |
| | 47-60 | 9 | 67 | 20-27 | 1.40-1.60 | 0.60-2.00 | 0.20-0.22 | 1.0-3.0 | 0.0-0.5 | .43 | .43 | | | |
| Hr: Hord----- | 0-13 | 11 | 67 | 17-27 | 1.30-1.40 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 2.0-4.0 | .32 | .32 | 5 | 6 | 48 |
| | 13-43 | 9 | 64 | 20-35 | 1.35-1.45 | 0.60-2.00 | 0.17-0.22 | 0.0-5.9 | 0.5-1.0 | .43 | .43 | | | |
| | 43-60 | 9 | 67 | 18-30 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| InB: Inavale----- | 0-5 | 87 | 7 | 2-10 | 1.50-1.60 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 2 | 134 |
| | 5-10 | 87 | 7 | 3-10 | 1.50-1.60 | 5.95-19.98 | 0.06-0.11 | 0.0-2.9 | 0.0-0.5 | .17 | .17 | | | |
| | 10-60 | 92 | 1 | 3-10 | 1.50-1.60 | 5.95-19.98 | 0.05-0.10 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| Ke: Kenesaw----- | 0-8 | 11 | 67 | 18-25 | 1.20-1.40 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 8-22 | 14 | 71 | 10-20 | 1.20-1.30 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 22-60 | 14 | 73 | 8-18 | 1.30-1.40 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| KgB: Kenesaw----- | 0-7 | 11 | 67 | 18-25 | 1.20-1.40 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 7-18 | 14 | 71 | 10-20 | 1.20-1.30 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 18-60 | 14 | 73 | 8-18 | 1.30-1.40 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| Coly----- | 0-5 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 5-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| Lf: Lex----- | 0-11 | 42 | 37 | 15-27 | 1.25-1.45 | 0.60-2.00 | 0.20-0.22 | 0.0-2.9 | 1.0-3.0 | .28 | .28 | 4 | 4L | 86 |
| | 11-24 | | | 18-32 | 1.30-1.70 | 0.20-2.00 | 0.15-0.22 | 0.0-4.8 | 0.5-1.0 | .37 | .37 | | | |
| | 24-60 | 90 | 6 | 2-5 | 1.50-1.70 | 19.98-19.98 | 0.02-0.06 | 0.0-2.9 | 0.0-0.5 | .05 | .10 | | | |
| Lg: Lex----- | 0-12 | 42 | 38 | 15-25 | 1.40-1.60 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .28 | .28 | 4 | 4L | 86 |
| | 12-31 | 20 | 50 | 27-32 | 1.40-1.60 | 0.20-2.00 | 0.15-0.19 | 3.0-5.9 | 0.5-1.0 | .37 | .37 | | | |
| | 31-60 | 90 | 4 | 2-10 | 1.50-1.70 | 19.98-19.98 | 0.02-0.05 | 0.0-2.9 | 0.0-0.5 | .05 | .10 | | | |
| LoB: Libory----- | 0-16 | 86 | 7 | 2-12 | 1.50-1.70 | 5.95-19.98 | 0.07-0.12 | 0.0-2.9 | 1.0-3.0 | .17 | .17 | 5 | 2 | 134 |
| | 16-24 | 86 | 7 | 2-12 | 1.50-1.70 | 5.95-19.98 | 0.06-0.11 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | | | |
| | 24-60 | 24 | 52 | 15-32 | 1.20-1.40 | 0.60-2.00 | 0.17-0.22 | 0.0-4.8 | 0.0-0.5 | .43 | .43 | | | |
| M-W: Miscellaneous Water----- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| Ma: Massie----- | 0-10 | 20 | 48 | 27-40 | 1.40-1.50 | 0.20-0.60 | 0.21-0.24 | 3.0-5.9 | 2.0-8.0 | .37 | .37 | 3 | 7 | 38 |
| | 10-60 | 7 | 48 | 35-55 | 1.20-1.40 | 0.00-0.06 | 0.09-0.20 | 6.0-12.0 | 0.5-2.0 | .37 | .37 | | | |
| Pm: Platte----- | 0-6 | 44 | 41 | 10-20 | 1.50-1.70 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .28 | .28 | 3 | 4L | 86 |
| | 6-12 | 46 | 42 | 7-18 | 1.70-1.90 | 0.60-2.00 | 0.15-0.19 | 0.0-2.9 | 0.0-0.5 | .28 | .28 | | | |
| | 12-60 | 92 | 7 | 0-3 | 1.90-2.00 | 19.98-19.98 | 0.02-0.04 | 0.0-2.9 | 0.0-0.5 | .05 | .10 | | | |
| Ru: Rusco----- | 0-5 | 11 | 67 | 17-27 | 1.30-1.40 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 5 | 56 |
| | 5-30 | 7 | 62 | 28-35 | 1.20-1.30 | 0.20-0.60 | 0.18-0.20 | 3.0-5.9 | 0.5-1.0 | .43 | .43 | | | |
| | 30-60 | 11 | 69 | 15-25 | 1.40-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| Sc: Scott----- | 0-6 | 26 | 53 | 15-27 | 1.25-1.40 | 0.60-2.00 | 0.21-0.24 | 0.0-2.9 | 2.0-4.0 | .37 | .37 | 3 | 6 | 48 |
| | 6-37 | 23 | 29 | 40-55 | 1.20-1.40 | 0.00-0.06 | 0.08-0.16 | 7.0-12.0 | 1.0-2.0 | .37 | .37 | | | |
| | 37-48 | 19 | 48 | 27-40 | 1.15-1.40 | 0.20-0.60 | 0.18-0.20 | 6.0-8.9 | 0.5-1.0 | .43 | .43 | | | |
| | 48-60 | 24 | 50 | 18-35 | 1.30-1.50 | 0.60-2.00 | 0.14-0.22 | 3.0-5.9 | 0.5-1.0 | .43 | .43 | | | |
| SmB: Simeon----- | 0-6 | 68 | 20 | 6-18 | 1.30-1.50 | 2.00-6.00 | 0.13-0.15 | 0.0-2.9 | 0.5-1.0 | .24 | .24 | 5 | 3 | 86 |
| | 6-60 | 90 | 4 | 2-10 | 1.50-1.70 | 5.95-19.98 | 0.05-0.10 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| To: Tryon----- | 0-4 | 87 | 7 | 3-10 | 1.40-1.60 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 4.0-8.0 | .17 | .17 | 5 | 8 | 0 |
| | 4-60 | 95 | 1 | 1-7 | 1.50-1.70 | 5.95-19.98 | 0.06-0.08 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | | | |

PHYSICAL PROPERTIES OF THE SOILS
Kearney County, Nebraska: Published

(Single entries under "Sand and Silt" are a representative percentage are calculated using an algorithm. Entries under "Erosion factors--T" apply to the entire profile. Entries under "Wind erodibility group" and "Wind erodibility index" apply only to the surface layer)

| Map symbol and soil name | Depth | Sand | Silt | Clay | Moist bulk density | Permeability (Ksat) | Available water capacity | Linear extensibility | Organic matter | Erosion factors | | | Wind erodibility group | Wind erodibility index |
|--------------------------|-------|------|------|-------|--------------------|---------------------|--------------------------|----------------------|----------------|-----------------|-----|---|------------------------|------------------------|
| | | | | | | | | | | K | Kf | T | | |
| | In | Pct | Pct | Pct | g/cc | in/hr | In/in | Pct | Pct | | | | | |
| UcF: | | | | | | | | | | | | | | |
| Uly----- | 0-9 | 11 | 67 | 17-27 | 1.20-1.45 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-3.0 | .32 | .32 | 5 | 6 | 48 |
| | 9-27 | 9 | 66 | 20-30 | 1.25-1.45 | 0.60-2.00 | 0.18-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| | 27-60 | 10 | 68 | 18-27 | 1.25-1.45 | 0.60-2.00 | 0.16-0.20 | 0.0-2.9 | 0.0-0.5 | .43 | .43 | | | |
| Coly----- | 0-4 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.20-0.24 | 0.0-2.9 | 1.0-2.0 | .43 | .43 | 5 | 4L | 86 |
| | 4-60 | 11 | 68 | 18-24 | 1.30-1.50 | 0.60-2.00 | 0.17-0.22 | 0.0-2.9 | 0.5-1.0 | .43 | .43 | | | |
| UsF: | | | | | | | | | | | | | | |
| Ustorhents-- | 0-60 | | | 18-35 | 1.10-1.65 | 0.60-2.00 | 0.16-0.23 | 0.0-5.9 | 0.5-1.0 | .43 | .43 | 5 | 4L | 86 |
| VaB: | | | | | | | | | | | | | | |
| Valentine---- | 0-5 | 87 | 7 | 2-10 | 1.35-1.55 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 2 | 134 |
| | 5-8 | 87 | 7 | 2-10 | 1.55-1.75 | 5.95-19.98 | 0.09-0.11 | 0.0-2.9 | 0.0-0.5 | .17 | .17 | | | |
| | 8-60 | 96 | 1 | 0-6 | 1.60-1.80 | 5.95-19.98 | 0.05-0.07 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| VaD: | | | | | | | | | | | | | | |
| Valentine---- | 0-5 | 87 | 7 | 2-10 | 1.35-1.55 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 2 | 134 |
| | 5-8 | 87 | 7 | 2-10 | 1.55-1.75 | 5.95-19.98 | 0.09-0.11 | 0.0-2.9 | 0.0-0.5 | .17 | .17 | | | |
| | 8-60 | 96 | 1 | 0-6 | 1.60-1.80 | 5.95-19.98 | 0.05-0.07 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| VaF: | | | | | | | | | | | | | | |
| Valentine---- | 0-5 | 87 | 7 | 2-10 | 1.35-1.55 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 2 | 134 |
| | 5-8 | 87 | 7 | 2-10 | 1.55-1.75 | 5.95-19.98 | 0.09-0.11 | 0.0-2.9 | 0.0-0.5 | .17 | .17 | | | |
| | 8-60 | 96 | 1 | 0-6 | 1.60-1.80 | 5.95-19.98 | 0.05-0.07 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| VbD: | | | | | | | | | | | | | | |
| Valentine---- | 0-5 | 87 | 7 | 2-10 | 1.35-1.55 | 5.95-19.98 | 0.10-0.12 | 0.0-2.9 | 0.5-1.0 | .17 | .17 | 5 | 2 | 134 |
| | 5-8 | 87 | 7 | 2-10 | 1.55-1.75 | 5.95-19.98 | 0.09-0.11 | 0.0-2.9 | 0.0-0.5 | .17 | .17 | | | |
| | 8-60 | 96 | 1 | 0-6 | 1.60-1.80 | 5.95-19.98 | 0.05-0.07 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| Els----- | 0-6 | 87 | 7 | 3-10 | 1.55-1.60 | 5.95-19.98 | 0.08-0.13 | 0.0-2.9 | 0.5-3.0 | .17 | .17 | 5 | 2 | 134 |
| | 6-10 | 95 | 1 | 0-8 | 1.50-1.60 | 5.95-19.98 | 0.05-0.08 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| | 10-60 | 95 | 1 | 0-8 | 1.50-1.60 | 5.95-19.98 | 0.04-0.07 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |
| W: | | | | | | | | | | | | | | |
| Water----- | --- | | | --- | --- | --- | --- | --- | --- | --- | --- | - | --- | 0 |
| Wa: | | | | | | | | | | | | | | |
| Wann----- | 0-14 | 64 | 26 | 5-15 | 1.30-1.50 | 2.00-6.00 | 0.13-0.18 | 0.0-2.9 | 1.0-3.0 | .20 | .20 | 5 | 3 | 86 |
| | 14-48 | 63 | 26 | 3-18 | 1.50-1.70 | 2.00-6.00 | 0.11-0.17 | 0.0-2.9 | 0.5-1.0 | .28 | .28 | | | |
| | 48-60 | | | 3-22 | 1.35-1.80 | 2.00-6.00 | 0.05-0.17 | 0.0-2.9 | 0.0-0.5 | .15 | .15 | | | |

CHEMICAL PROPERTIES OF THE SOILS
Kearney County, Nebraska

The Chemical Properties table shows estimates of some characteristics and features that affect soil behavior. These estimates are given for the major layers of each soil in the survey area. The estimates are based on field observations and on test data for these and similar soils. Depth to the upper and lower boundaries of each layer is indicated.

Cation-exchange capacity is the total amount of extractable bases that can be held by the soil, expressed in terms of milliequivalents per 100 grams of soil at neutrality (pH 7.0) or at some other stated pH value. Soils having a low cation-exchange capacity hold fewer cations and may require more frequent applications of fertilizer than soils having a high cation-exchange capacity. Soils having a high cation-exchange capacity can retain cations. The ability to retain cations helps to prevent the pollution of ground water.

Soil reaction is a measure of acidity or alkalinity and is expressed as a range in pH values. The range in pH of each major horizon is based on many field tests. For many soils, values have been verified by laboratory analyses. Soil reaction is important in selecting crops and other plants, in evaluating soil amendments for fertility and stabilization, and in determining the risk of corrosion.

Calcium carbonate equivalent is the percent of carbonates, by weight, in the fraction of the soil less than 2 millimeters in size. The availability of plant nutrients is influenced by the amount of carbonates in the soil. Incorporating nitrogen fertilizer into calcareous soils helps to prevent nitrite accumulation and ammonium-N volatilization.

Gypsum is expressed as a percent, by weight, of hydrated calcium sulfates in the fraction of the soil less than 20 millimeters in size. Gypsum is partially soluble in water and can be dissolved and removed by water. Soils that have a high content of gypsum may collapse if the gypsum is removed by percolating water.

Salinity is a measure of soluble salts in the soil at saturation. It is expressed as the electrical conductivity of the saturation extract, in millimhos per centimeter at 25 degrees C. Estimates are based on field and laboratory measurements at representative sites of nonirrigated soils. The salinity of irrigated soils is affected by the quality of the irrigation water and by the frequency of water application. Hence, the salinity of soils in individual fields can differ greatly from the value given in the table. Salinity affects the suitability of a soil for crop production, the stability of soil if used as construction material, and the potential of the soil to corrode metal and concrete.

Sodium adsorption ratio (SAR) is a measure of the amount of sodium (Na) relative to calcium (Ca) and magnesium (Mg) in the water extract from saturated soil paste. It is the ratio of the Na concentration divided by the square root of one-half of the Ca + Mg concentration. Soils that have SAR values of 13 or more may be characterized by an increased dispersion of organic matter and clay particles, reduced permeability and aeration, and a general degradation of soil structure.

CHEMICAL PROPERTIES OF THE SOILS--Continued
 Kearney County, Nebraska

| Map symbol and soil name | Depth | Cation- exchange capacity | Soil reaction | Calcium carbonate | Gypsum | Salinity | Sodium adsorption ratio |
|-----------------------------|-------|---------------------------------|------------------|----------------------|--------|----------|-------------------------------|
| | In | meq/100g | pH | Pct | Pct | mmhos/cm | |
| Ad: | | | | | | | |
| Alda----- | 0-12 | 10-30 | 6.6-8.4 | 0-10 | 0 | 0.0-4.0 | 0-9 |
| | 12-30 | 5.0-10 | 7.4-8.4 | 1-15 | 0 | 0.0-4.0 | 0-9 |
| | 30-60 | 0.0-5.0 | 6.6-8.4 | 0-5 | 0 | 0.0-4.0 | 0-9 |
| Bo: | | | | | | | |
| Boel----- | 0-17 | 8.0-17 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 17-54 | 0.0-4.0 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 54-60 | 0.0-4.0 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| Bu: | | | | | | | |
| Butler----- | 0-12 | 18-27 | 5.1-6.5 | 0 | 0 | 0 | 0 |
| | 12-35 | 30-40 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| | 35-41 | 20-35 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 41-60 | 10-30 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| CaC: | | | | | | | |
| Coly----- | 0-5 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 5-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| CaD: | | | | | | | |
| Coly----- | 0-5 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 5-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| CaF: | | | | | | | |
| Coly----- | 0-4 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 4-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| CkB: | | | | | | | |
| Coly----- | 0-6 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 6-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| Kenesaw----- | 0-10 | 9.0-17 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 10-23 | 7.0-13 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 23-60 | 5.0-13 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| CoD2: | | | | | | | |
| Coly----- | 0-5 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 5-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| Uly----- | 0-7 | 10-20 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 7-28 | 14-25 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 28-60 | 12-20 | 7.4-8.4 | 1-15 | 0 | 0 | 0 |
| CoF2: | | | | | | | |
| Coly----- | 0-6 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 6-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| Uly----- | 0-7 | 10-20 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 7-28 | 14-25 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 28-60 | 12-20 | 7.4-8.4 | 1-15 | 0 | 0 | 0 |
| De: | | | | | | | |
| Detroit----- | 0-16 | 10-25 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 16-38 | 15-30 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 38-60 | 10-20 | 6.6-8.4 | 0-10 | 0 | 0 | 0-1 |
| Fm: | | | | | | | |
| Fillmore----- | 0-12 | 15-22 | 5.1-6.5 | 0 | 0 | 0 | 0 |
| | 12-35 | 32-40 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| | 35-44 | 22-30 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 44-60 | 12-40 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| Gb: | | | | | | | |
| Gibbon----- | 0-14 | 8.0-22 | 7.4-8.4 | 0-5 | 0 | 0.0-2.0 | 0 |
| | 14-37 | 14-20 | 7.4-8.4 | 5-15 | 0 | 0.0-2.0 | 0-5 |
| | 37-60 | 10-18 | 7.9-8.4 | 5-15 | 0 | 0.0-2.0 | 0-5 |
| Gc: | | | | | | | |
| Gibbon----- | 0-10 | 11-21 | 7.4-8.4 | 0-5 | 0 | 4.0-16.0 | 0-10 |
| | 10-40 | 14-23 | 7.4-8.4 | 5-15 | 0 | 4.0-8.0 | 0-10 |
| | 40-60 | 21-30 | 7.4-8.4 | 5-15 | 0 | 0.0-4.0 | 0-10 |
| Go: | | | | | | | |
| Gothenburg----- | 0-3 | 0.0-10 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 3-8 | 0.0-5.0 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 8-60 | 0.0-5.0 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| GP: | | | | | | | |
| Pits----- | 0-60 | 0.0-5.0 | 6.6-8.4 | 0 | 0 | 0 | 0 |
| HeB: | | | | | | | |
| Hersh----- | 0-7 | 10-15 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 7-11 | 8.0-12 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 11-60 | 8.0-12 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| HeC: | | | | | | | |
| Hersh----- | 0-7 | 10-15 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 7-11 | 8.0-12 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 11-60 | 8.0-12 | 6.6-7.8 | 0 | 0 | 0 | 0 |

CHEMICAL PROPERTIES OF THE SOILS--Continued
Kearney County, Nebraska

| Map symbol and soil name | Depth | Cation- exchange capacity | Soil reaction | Calcium carbonate | Gypsum | Salinity | Sodium adsorption ratio |
|-----------------------------|-------|---------------------------------|------------------|----------------------|--------|----------|-------------------------------|
| | In | meq/100g | pH | Pct | Pct | mmhos/cm | |
| HeD: | | | | | | | |
| Hersh----- | 0-7 | 10-15 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 7-11 | 8.0-12 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 11-60 | 8.0-12 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| Hf: | | | | | | | |
| Hobbs----- | 0-6 | 15-30 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 6-34 | 10-20 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 34-60 | 20-40 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| HgB: | | | | | | | |
| Hobbs----- | 0-9 | 15-30 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 9-34 | 10-20 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 34-60 | 20-40 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| Ho: | | | | | | | |
| Holdrege----- | 0-13 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 13-25 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 25-31 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 31-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| HoB: | | | | | | | |
| Holdrege----- | 0-13 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 13-25 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 25-31 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 31-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| HoC: | | | | | | | |
| Holdrege----- | 0-13 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 13-25 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 25-31 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 31-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| HoC2: | | | | | | | |
| Holdrege----- | 0-13 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 13-25 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 25-31 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 31-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| Hp: | | | | | | | |
| Holdrege----- | 0-27 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 27-43 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 43-47 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 47-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| HpB: | | | | | | | |
| Holdrege----- | 0-27 | 16-24 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 27-43 | 19-26 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 43-47 | 14-21 | 6.6-7.8 | 0 | 0 | 0 | 0 |
| | 47-60 | 14-19 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| Hr: | | | | | | | |
| Hord----- | 0-13 | 10-20 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 13-43 | 10-30 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 43-60 | 10-35 | 7.4-8.4 | 0-5 | 0 | 0 | 0 |
| InB: | | | | | | | |
| Inavale----- | 0-5 | 2.0-10 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| | 5-10 | 1.0-10 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| | 10-60 | 1.0-10 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| Ke: | | | | | | | |
| Kenesaw----- | 0-8 | 9.0-17 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 8-22 | 7.0-13 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 22-60 | 5.0-13 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| KgB: | | | | | | | |
| Kenesaw----- | 0-7 | 9.0-17 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 7-18 | 7.0-13 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 18-60 | 5.0-13 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| Coly----- | 0-5 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 5-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| Lf: | | | | | | | |
| Lex----- | 0-11 | 11-22 | 7.4-8.4 | 1-10 | 0 | 0.0-4.0 | 0-6 |
| | 11-24 | 12-23 | 6.1-8.4 | 0-5 | 0 | 0.0-4.0 | 0-6 |
| | 24-60 | 2.0-5.0 | 6.1-7.8 | 0-5 | 0 | 0.0-4.0 | 0-6 |
| Lg: | | | | | | | |
| Lex----- | 0-12 | 11-21 | 7.9-9.0 | 1-10 | 0 | 4.0-16.0 | 0-13 |
| | 12-31 | 18-23 | 7.9-9.0 | 1-10 | 0 | 4.0-16.0 | 0-13 |
| | 31-60 | 1.0-7.0 | 7.9-8.4 | 1-10 | 0 | 0.0-4.0 | 0-13 |

CHEMICAL PROPERTIES OF THE SOILS--Continued
Kearney County, Nebraska

| Map symbol and soil name | Depth | Cation- exchange capacity | Soil reaction | Calcium carbonate | Gypsum | Salinity | Sodium adsorption ratio |
|-----------------------------|-------|---------------------------------|------------------|----------------------|--------|----------|-------------------------------|
| | In | meq/100g | pH | Pct | Pct | mmhos/cm | |
| LoB: | | | | | | | |
| Libory----- | 0-16 | 2.0-12 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 16-24 | 1.0-9.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 24-60 | 10-22 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| M-W: | | | | | | | |
| Miscellaneous Water----- | --- | --- | --- | --- | --- | --- | --- |
| Ma: | | | | | | | |
| Massie----- | 0-10 | 10-40 | 5.1-6.5 | 0 | 0 | 0 | 0 |
| | 10-60 | 25-40 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| Pm: | | | | | | | |
| Platte----- | 0-6 | 5.0-20 | 6.6-8.4 | 0-10 | 0 | 0.0-2.0 | 0 |
| | 6-12 | 4.0-20 | 6.6-8.4 | 0-5 | 0 | 0.0-2.0 | 0 |
| | 12-60 | 0.0-3.0 | 6.6-8.4 | 0-5 | 0 | 0.0-2.0 | 0 |
| Ru: | | | | | | | |
| Rusco----- | 0-5 | 10-30 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 5-30 | 20-30 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 30-60 | 10-20 | 7.4-8.4 | 1-10 | 0 | 0 | 0 |
| Sc: | | | | | | | |
| Scott----- | 0-6 | 20-30 | 5.1-6.5 | 0 | 0 | 0 | 0 |
| | 6-37 | 20-35 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| | 37-48 | 20-35 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 48-60 | 25-35 | 6.6-8.4 | 0-5 | 0 | 0 | 0 |
| SmB: | | | | | | | |
| Simeon----- | 0-6 | 0.0-10 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 6-60 | 0.0-5.0 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| To: | | | | | | | |
| Tryon----- | 0-4 | 10-25 | 5.6-8.4 | 0-5 | 0 | 0 | 0 |
| | 4-60 | 1.0-5.0 | 5.6-7.8 | 0 | 0 | 0 | 0 |
| UcF: | | | | | | | |
| Uly----- | 0-9 | 10-20 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 9-27 | 14-25 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| | 27-60 | 12-20 | 7.4-8.4 | 1-15 | 0 | 0 | 0 |
| | 0-4 | 14-19 | 7.4-8.4 | 1-5 | 0 | 0 | 0 |
| | 4-60 | 13-18 | 7.4-8.4 | 5-10 | 0 | 0 | 0 |
| UsF: | | | | | | | |
| Ustorthents---- | 0-60 | 10-25 | 7.4-8.4 | 1-15 | 0 | 0.0-2.0 | 0 |
| VaB: | | | | | | | |
| Valentine----- | 0-5 | 2.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 5-8 | 1.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 8-60 | 0.0-5.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| VaD: | | | | | | | |
| Valentine----- | 0-5 | 2.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 5-8 | 1.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 8-60 | 0.0-5.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| VaF: | | | | | | | |
| Valentine----- | 0-5 | 2.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 5-8 | 1.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 8-60 | 0.0-5.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| VbD: | | | | | | | |
| Valentine----- | 0-5 | 2.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 5-8 | 1.0-8.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 8-60 | 0.0-5.0 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 0-6 | 5.0-10 | 5.6-7.3 | 0 | 0 | 0 | 0 |
| | 6-10 | 0.0-5.0 | 6.1-7.3 | 0 | 0 | 0 | 0 |
| | 10-60 | 0.0-5.0 | 6.1-7.8 | 0 | 0 | 0 | 0 |
| W: | | | | | | | |
| Water----- | --- | --- | --- | --- | --- | --- | --- |
| Wa: | | | | | | | |
| Wann----- | 0-14 | 4.0-14 | 6.6-8.4 | 0-5 | 0 | 0.0-2.0 | 0-5 |
| | 14-48 | 2.0-14 | 7.4-9.0 | 0-5 | 0 | 0.0-2.0 | 0-10 |
| | 48-60 | 2.0-16 | 7.4-9.0 | 0-5 | 0 | 0.0-2.0 | 0-10 |

WATER FEATURES
Kearney County, Nebraska

The Water Features table gives estimates of various water features. The estimates are used in land use planning that involves engineering considerations. Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The four hydrologic soil groups are:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

The months in the table indicate the portion of the year in which the feature is most likely to be a concern.

Water table refers to a saturated zone in the soil. The Water Features table indicates, by month, depth to the top (upper limit) and base (lower limit) of the saturated zone in most years. Estimates of the upper and lower limits are based mainly on observations of the water table at selected sites and on evidence of a saturated zone, namely grayish colors or mottles (redoximorphic features) in the soil. A saturated zone that lasts for less than a month is not considered a water table. Ponding is standing water in a closed depression. Unless a drainage system is installed, the water is removed only by percolation, transpiration, or evaporation. The Water Features table indicates surface water depth and the duration and frequency of ponding. Duration is expressed as very brief if less than 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, rare, occasional, and frequent. None means that ponding is not probable; rare that it is unlikely but possible under unusual weather conditions (the chance of ponding is nearly 0 percent to 5 percent in any year); occasional that it occurs, on the average, once or less in 2 years (the chance of ponding is 5 to 50 percent in any year); and frequent that it occurs, on the average, more than once in 2 years (the chance of ponding is more than 50 percent in any year).

Flooding, the temporary inundation of an area, is caused by overflowing streams, by runoff from adjacent slopes, or by tides. Water standing for short periods after rainfall or snowmelt is not considered flooding, and water standing in swamps and marshes is considered ponding rather than flooding.

Duration and frequency are estimated. Duration is expressed as extremely brief if 0.1 hour to 4 hours, very brief if 4 hours to 2 days, brief if 2 to 7 days, long if 7 to 30 days, and very long if more than 30 days. Frequency is expressed as none, very rare, rare, occasional, frequent, and very frequent. None means that flooding is not probable; very rare that it is very unlikely but possible under extremely unusual weather conditions (the chance of flooding is less than 1 percent in any year); rare that it is unlikely but possible under unusual weather conditions (the chance of flooding is 1 to 5 percent in any year); occasional that it occurs infrequently under normal weather conditions (the chance of flooding is 5 to 50 percent in any year); frequent that it is likely to occur often under normal weather conditions (the chance of flooding is more than 50 percent in any year but is less than 50 percent in all months in any year); and very frequent that it is likely to occur very often under normal weather conditions (the chance of flooding is more than 50 percent in all months of any year).

The information is based on evidence in the soil profile, namely thin strata of gravel, sand, silt, or clay deposited by floodwater; irregular decrease in organic matter content with increasing depth; and little or no horizon development.

Also considered are local information about the extent and levels of flooding and the relation of each soil on the landscape to historic floods. Information on the extent of flooding based on soil data is less specific than that provided by detailed engineering surveys that delineate flood-prone areas at specific flood frequency levels.

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

| Map symbol and soil name | Hydro- logic group | Month | Soil Saturation | | Ponding | | | Flooding | | |
|-----------------------------|--------------------------|----------|-----------------|----------------|---------------------------|----------|-----------|----------|-----------|------|
| | | | Upper limit | Lower limit | Surface water depth | Duration | Frequency | Duration | Frequency | |
| | | | Ft | Ft | Ft | | | | | |
| Ad: Alda----- | C | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | | February | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | | March | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | | April | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | | May | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | | June | --- | --- | --- | --- | --- | Brief | Rare | |
| | | July | --- | --- | --- | --- | --- | Brief | Rare | |
| | | November | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | | December | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| Bo: Boel----- | | A | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | | February | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | | March | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare |
| | April | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | May | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | June | | --- | --- | --- | --- | --- | Brief | Rare | |
| | November | | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | December | | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| Bu: Butler----- | D | | March | 0.5-1.5 | 1.0-2.0 | --- | --- | --- | --- | None |
| | | | April | 0.5-1.5 | 1.0-2.0 | --- | --- | --- | --- | None |
| | | | May | 0.5-1.5 | 1.0-2.0 | --- | --- | --- | --- | None |
| | | | June | 0.5-1.5 | 1.0-2.0 | --- | --- | --- | --- | None |
| | | July | 0.5-1.5 | 1.0-2.0 | --- | --- | --- | --- | None | |
| CaC: Coly----- | | B | | --- | --- | --- | --- | --- | --- | |
| CaD: Coly----- | | B | | --- | --- | --- | --- | --- | --- | |
| CaF: Coly----- | B | | --- | --- | --- | --- | --- | --- | | |
| CkB: Coly----- | B | | --- | --- | --- | --- | --- | --- | | |
| Kenesaw----- | B | | --- | --- | --- | --- | --- | --- | | |
| CoD2: Coly----- | B | | --- | --- | --- | --- | --- | --- | | |
| Uly----- | B | | --- | --- | --- | --- | --- | --- | | |
| CoF2: Coly----- | B | | --- | --- | --- | --- | --- | --- | | |
| Uly----- | B | | --- | --- | --- | --- | --- | --- | | |
| De: Detroit----- | C | | --- | --- | --- | --- | --- | --- | | |
| Fm: Fillmore----- | D | March | 0.0-2.0 | 1.0-3.0 | 0.0-0.5 | Brief | --- | --- | None | |
| | | April | 0.0-2.0 | 1.0-3.0 | 0.0-0.5 | Brief | --- | --- | None | |
| | | May | 0.0-2.0 | 1.0-3.0 | 0.0-0.5 | Brief | --- | --- | None | |
| | | June | 0.0-2.0 | 1.0-3.0 | 0.0-0.5 | Brief | --- | --- | None | |
| | | July | 0.0-2.0 | 1.0-3.0 | 0.0-0.5 | Brief | --- | --- | None | |
| Gb: Gibbon----- | | B | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | February | | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | March | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | April | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | May | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | June | | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | July | | --- | --- | --- | --- | --- | Brief | Rare | |
| | November | | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | December | | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| Gc: Gibbon----- | B | | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | | February | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | | March | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare |
| | | April | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | | May | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | | June | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | |
| | | July | --- | --- | --- | --- | --- | Brief | Rare | |
| | | November | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |
| | | December | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | |

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

| Map symbol and soil name | Hydro- logic group | Month | Soil Saturation | | Ponding | | | Flooding | |
|-----------------------------|--------------------------|-----------|-----------------|----------------|---------------------------|----------|-----------|------------|------------|
| | | | Upper limit | Lower limit | Surface water depth | Duration | Frequency | Duration | Frequency |
| | | | Ft | Ft | Ft | | | | |
| Go: Gothenburg----- | D | January | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | February | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | March | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | April | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | May | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | June | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| | | July | --- | --- | --- | --- | --- | Brief | Frequent |
| | | November | 0.0-1.5 | >6.0 | --- | --- | --- | --- | None |
| | | December | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Frequent |
| GP: Pits----- | A | | --- | --- | --- | --- | --- | --- | --- |
| HeB: Hersh----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HeC: Hersh----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HeD: Hersh----- | B | | --- | --- | --- | --- | --- | --- | --- |
| Hf: Hobbs----- | B | April | --- | --- | --- | --- | --- | Brief | Occasional |
| | | May | --- | --- | --- | --- | --- | Brief | Occasional |
| | | June | --- | --- | --- | --- | --- | Brief | Occasional |
| | | July | --- | --- | --- | --- | --- | Brief | Occasional |
| | | August | --- | --- | --- | --- | --- | Brief | Occasional |
| | | September | --- | --- | --- | --- | --- | Brief | Occasional |
| HgB: Hobbs----- | B | April | --- | --- | --- | --- | --- | Brief | Frequent |
| | | May | --- | --- | --- | --- | --- | Brief | Frequent |
| | | June | --- | --- | --- | --- | --- | Brief | Frequent |
| | | July | --- | --- | --- | --- | --- | Brief | Frequent |
| | | August | --- | --- | --- | --- | --- | Brief | Frequent |
| | | September | --- | --- | --- | --- | --- | Brief | Frequent |
| Ho: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HoB: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HoC: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HoC2: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| Hp: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| HpB: Holdrege----- | B | | --- | --- | --- | --- | --- | --- | --- |
| Hr: Hord----- | B | | --- | --- | --- | --- | --- | --- | --- |
| InB: Inavale----- | A | January | --- | --- | --- | --- | --- | Very brief | Rare |
| | | February | --- | --- | --- | --- | --- | Very brief | Rare |
| | | March | --- | --- | --- | --- | --- | Very brief | Rare |
| | | April | --- | --- | --- | --- | --- | Very brief | Rare |
| | | May | --- | --- | --- | --- | --- | Very brief | Rare |
| | | June | --- | --- | --- | --- | --- | Very brief | Rare |
| | | July | --- | --- | --- | --- | --- | Very brief | Rare |
| Ke: Kenesaw----- | B | | --- | --- | --- | --- | --- | --- | --- |
| KgB: Kenesaw----- | B | | --- | --- | --- | --- | --- | --- | --- |
| Coly----- | B | | --- | --- | --- | --- | --- | --- | --- |
| Lf: | | | --- | --- | --- | --- | --- | --- | --- |

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

| Map symbol and soil name | Hydro-logic group | Month | Soil Saturation | | Ponding | | | Flooding | |
|-----------------------------|-------------------|----------|-----------------|-------------|---------------------|----------|-----------|------------|-----------|
| | | | Upper limit | Lower limit | Surface water depth | Duration | Frequency | Duration | Frequency |
| Lex----- Lg: Lex----- | B | January | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | February | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | March | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | April | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Rare |
| | | May | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Rare |
| | | June | --- | --- | --- | --- | --- | Brief | Rare |
| | | July | --- | --- | --- | --- | --- | Brief | Rare |
| | | November | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | December | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | February | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | March | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| April | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | | |
| May | 1.5-3.0 | >6.0 | --- | --- | --- | Brief | Rare | | |
| June | --- | --- | --- | --- | --- | Brief | Rare | | |
| July | --- | --- | --- | --- | --- | Brief | Rare | | |
| November | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | | |
| December | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None | | |
| LoB: Libory----- | A | March | 1.5-3.0 | 1.5-3.0 | --- | --- | --- | --- | None |
| April | | 1.5-3.0 | 1.5-3.0 | --- | --- | --- | --- | None | |
| May | | 1.5-3.0 | 1.5-3.0 | --- | --- | --- | --- | None | |
| June | | 1.5-3.0 | 1.5-3.0 | --- | --- | --- | --- | None | |
| March | | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None |
| April | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None | |
| May | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None | |
| June | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None | |
| July | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None | |
| August | 0.0-1.0 | 0.5-1.5 | 0.0-2.0 | Very long | --- | --- | --- | None | |
| Pm: Platte----- | B | February | 1.0-3.0 | >6.0 | --- | --- | --- | --- | None |
| March | | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Occasional | |
| April | | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Occasional | |
| May | | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Occasional | |
| June | | 1.0-3.0 | >6.0 | --- | --- | --- | Brief | Occasional | |
| July | | --- | --- | --- | --- | --- | Brief | Occasional | |
| August | | --- | --- | --- | --- | --- | Brief | Occasional | |
| September | | --- | --- | --- | --- | --- | Brief | Occasional | |
| October | | --- | --- | --- | --- | --- | Brief | Occasional | |
| March | | 0.0 | >6.0 | 0.0-0.5 | Brief | --- | --- | --- | None |
| April | 0.0 | >6.0 | 0.0-0.5 | Brief | --- | --- | --- | None | |
| May | 0.0 | >6.0 | 0.0-0.5 | Brief | --- | --- | --- | None | |
| June | 0.0 | >6.0 | 0.0-0.5 | Brief | --- | --- | --- | None | |
| July | 0.0 | >6.0 | 0.0-0.5 | Brief | --- | --- | --- | None | |
| Sc: Scott----- | D | March | 0.0-1.0 | 1.0-2.0 | 0.0-1.0 | Long | --- | --- | None |
| April | | 0.0-1.0 | 1.0-2.0 | 0.0-1.0 | Long | --- | --- | None | |
| May | | 0.0-1.0 | 1.0-2.0 | 0.0-1.0 | Long | --- | --- | None | |
| June | | 0.0-1.0 | 1.0-2.0 | 0.0-1.0 | Long | --- | --- | None | |
| July | | 0.0-1.0 | 1.0-2.0 | 0.0-1.0 | Long | --- | --- | None | |
| March | | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Rare | |
| February | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Rare | | |
| March | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Rare | | |
| April | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Rare | | |
| May | 0.0-1.5 | >6.0 | --- | --- | --- | Brief | Rare | | |
| June | --- | --- | --- | --- | --- | Brief | Rare | | |
| July | --- | --- | --- | --- | --- | Brief | Rare | | |
| November | 0.0-1.5 | >6.0 | --- | --- | --- | --- | None | | |
| December | 0.0-1.5 | >6.0 | --- | --- | --- | --- | None | | |
| UcF: Uly----- | B | | | | | | | | |
| Coly----- | B | | | | | | | | |
| UsF: Ustorthents----- | B | | | | | | | | |
| VaB: Valentine----- | A | | | | | | | | |

(Depths of layers are in feet. See text for definitions of terms used in this table. Estimates of the frequency of ponding and flooding apply to the whole year rather than to individual months. Absence of an entry indicates that the feature is not a concern or that data were not estimated.)

| Map symbol and soil name | Hydro- logic group | Month | Soil Saturation | | Ponding | | | Flooding | |
|-----------------------------|--------------------------|----------|-----------------|----------------|---------------------------|----------|-----------|----------|-----------|
| | | | Upper limit | Lower limit | Surface water depth | Duration | Frequency | Duration | Frequency |
| VaD: Valentine----- | A | | --- | --- | --- | --- | --- | --- | --- |
| VaF: Valentine----- | A | | --- | --- | --- | --- | --- | --- | --- |
| VbD: Valentine----- | A | | --- | --- | --- | --- | --- | --- | --- |
| Els----- | A | | --- | --- | --- | --- | --- | --- | --- |
| | | January | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | February | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | March | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | April | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | May | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | November | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| | | December | 1.5-3.0 | >6.0 | --- | --- | --- | --- | None |
| W: Water----- | --- | | --- | --- | --- | --- | --- | --- | --- |
| Wa: Wann----- | B | | --- | --- | --- | --- | --- | --- | --- |
| | | March | 1.5-3.5 | >6.0 | --- | --- | --- | Brief | Rare |
| | | April | 1.5-3.5 | >6.0 | --- | --- | --- | Brief | Rare |
| | | May | 1.5-3.5 | >6.0 | --- | --- | --- | Brief | Rare |
| | | June | 1.5-3.5 | >6.0 | --- | --- | --- | Brief | Rare |
| | | July | 1.5-3.5 | >6.0 | --- | --- | --- | Brief | Rare |

The following table gives estimates of various soil features. The estimates are used in land use planning that involves engineering considerations.

A restrictive layer is a nearly continuous layer that has one or more physical, chemical, or thermal properties that significantly impede the movement of water and air through the soil or that restrict roots or otherwise provide an unfavorable root environment. Examples are bedrock, cemented layers, dense layers, and frozen layers. The table indicates the hardness and thickness of the restrictive layer, both of which significantly affect the ease of excavation. Depth to top is the vertical distance from the soil surface to the upper boundary of the restrictive layer.

Potential for frost action is the likelihood of upward or lateral expansion of the soil caused by the formation of segregated ice lenses (frost heave) and the subsequent collapse of the soil and loss of strength on thawing. Frost action occurs when moisture moves into the freezing zone of the soil. Temperature, texture, density, permeability, content of organic matter, and depth to the water table are the most important factors considered in evaluating the potential for frost action. It is assumed that the soil is not insulated by vegetation or snow and is not artificially drained. Silty and highly structured, clayey soils that have a high water table in winter are the most susceptible to frost action. Well drained, very gravelly, or very sandy soils are the least susceptible. Frost heave and low soil strength during thawing cause damage to pavements and other rigid structures.

Risk of corrosion pertains to potential soil-induced electrochemical or chemical action that corrodes or weakens uncoated steel or concrete. The rate of corrosion of uncoated steel is related to such factors as soil moisture, particle-size distribution, acidity, and electrical conductivity of the soil. The rate of corrosion of concrete is based mainly on the sulfate and sodium content, texture, moisture content, and acidity of the soil. Special site examination and design may be needed if the combination of factors results in a severe hazard of corrosion. The steel or concrete in installations that intersect soil boundaries or soil layers is more susceptible to corrosion than the steel or concrete in installations that are entirely within one kind of soil or within one soil layer.

For uncoated steel, the risk of corrosion, expressed as low, moderate, or high, is based on soil drainage class, total acidity, electrical resistivity near field capacity, and electrical conductivity of the saturation extract.

For concrete, the risk of corrosion also is expressed as low, moderate, or high. It is based on soil texture, acidity, and amount of sulfates in the saturation extract.

SOIL FEATURES--Continued
Kearney County, Nebraska

| Map symbol and soil name | Restrictive layer | | | | Potential for Frost action | Risk of corrosion | |
|-----------------------------|-------------------|-----------------|-----------|----------|----------------------------------|-------------------|----------|
| | Kind | Depth to top | Thickness | Hardness | | Uncoated Steel | Concrete |
| | | In | In | | | | |
| Ad: | | | | | | | |
| Alda----- | --- | --- | --- | --- | High | Moderate | Low |
| Bo: | | | | | | | |
| Boel----- | --- | --- | --- | --- | Moderate | High | Low |
| Bu: | | | | | | | |
| Butler----- | --- | --- | --- | --- | High | High | Low |
| CaC: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| CaD: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| CaF: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| CkB: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| Kenesaw----- | --- | --- | --- | --- | Moderate | High | Low |
| CoD2: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| Uly----- | --- | --- | --- | --- | Moderate | High | Low |
| CoF2: | | | | | | | |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| Uly----- | --- | --- | --- | --- | Moderate | High | Low |
| De: | | | | | | | |
| Detroit----- | --- | --- | --- | --- | Low | High | Low |
| Fm: | | | | | | | |
| Fillmore----- | --- | --- | --- | --- | High | High | Low |
| Gb: | | | | | | | |
| Gibbon----- | --- | --- | --- | --- | High | High | Low |
| Gc: | | | | | | | |
| Gibbon----- | --- | --- | --- | --- | High | High | High |
| Go: | | | | | | | |
| Gothenburg----- | --- | --- | --- | --- | Moderate | Moderate | Low |
| GP: | | | | | | | |
| Pits----- | --- | --- | --- | --- | Low | Low | Low |
| HeB: | | | | | | | |
| Hersh----- | --- | --- | --- | --- | Moderate | Low | Low |
| HeC: | | | | | | | |
| Hersh----- | --- | --- | --- | --- | Moderate | Low | Low |
| HeD: | | | | | | | |
| Hersh----- | --- | --- | --- | --- | Moderate | Low | Low |
| Hf: | | | | | | | |
| Hobbs----- | --- | --- | --- | --- | Moderate | Low | Low |
| HgB: | | | | | | | |
| Hobbs----- | --- | --- | --- | --- | Moderate | Low | Low |
| Ho: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| HoB: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| HoC: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| HoC2: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| Hp: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| HpB: | | | | | | | |
| Holdrege----- | --- | --- | --- | --- | Moderate | Low | Low |
| Hr: | | | | | | | |
| Hord----- | --- | --- | --- | --- | Moderate | High | Low |
| InB: | | | | | | | |
| Inavale----- | --- | --- | --- | --- | Low | Moderate | Low |
| Ke: | | | | | | | |
| Kenesaw----- | --- | --- | --- | --- | Moderate | Moderate | Low |
| KgB: | | | | | | | |
| Kenesaw----- | --- | --- | --- | --- | Moderate | Moderate | Low |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| Lf: | | | | | | | |
| Lex----- | --- | --- | --- | --- | High | High | Low |
| Lg: | | | | | | | |
| Lex----- | --- | --- | --- | --- | High | High | High |
| LoB: | | | | | | | |
| Libory----- | --- | --- | --- | --- | Low | Moderate | Low |
| M-W: | | | | | | | |
| Miscellaneous Water----- | --- | --- | --- | --- | --- | --- | --- |
| Ma: | | | | | | | |
| Massie----- | --- | --- | --- | --- | High | High | Low |
| Pm: | | | | | | | |
| Platte----- | --- | --- | --- | --- | Moderate | High | Moderate |
| Ru: | | | | | | | |
| Rusco----- | --- | --- | --- | --- | High | High | Low |
| Sc: | | | | | | | |
| Scott----- | --- | --- | --- | --- | High | High | Low |
| SmB: | | | | | | | |
| Simeon----- | --- | --- | --- | --- | Low | Low | Low |
| To: | | | | | | | |
| Tryon----- | --- | --- | --- | --- | Moderate | High | Low |

| Map symbol and soil name | Restrictive layer | | | | Potential for Frost action | Risk of corrosion | |
|-----------------------------|-------------------|-----------------|-----------|----------|----------------------------------|-------------------|----------|
| | Kind | Depth to top | Thickness | Hardness | | Uncoated Steel | Concrete |
| | | In | In | | | | |
| UcF: | | | | | | | |
| Uly----- | --- | --- | --- | --- | Moderate | High | Low |
| Coly----- | --- | --- | --- | --- | Moderate | High | Low |
| UsF: | | | | | | | |
| Ustorthents---- | --- | --- | --- | --- | Moderate | High | Moderate |
| VaB: | | | | | | | |
| Valentine----- | --- | --- | --- | --- | Low | Low | Low |
| VaD: | | | | | | | |
| Valentine----- | --- | --- | --- | --- | Low | Low | Low |
| VaF: | | | | | | | |
| Valentine----- | --- | --- | --- | --- | Low | Low | Low |
| VbD: | | | | | | | |
| Valentine----- | --- | --- | --- | --- | Low | Low | Low |
| Els----- | --- | --- | --- | --- | Moderate | Moderate | Low |
| W: | | | | | | | |
| Water----- | --- | --- | --- | --- | --- | --- | --- |
| Wa: | | | | | | | |
| Wann----- | --- | --- | --- | --- | High | Moderate | Low |

WATER MANAGEMENT
Kearney County, Nebraska

The soils of the survey area are rated in the Water Management table according to limitations that affect their suitability for water management. Soils are rated for pond reservoir areas, drainage, irrigation, terraces and diversions, and grassed waterways. Restrictive features that affect each soil for the specified use is also provided in the table.

The ratings in the table are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect the specified use. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Moderately limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Limited indicates that the soil has one or more features that are significant limitations for the specified use. The limitations can be overcome, but generally require special design, soil reclamation, or installation procedures that may result in additional expense. Fair performance and moderate to high maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Limitation class terms, such as very limited or limited, etc., limitation ratings, and numerical ratings are shown for each soil feature listed. As many as three soil features may be listed for each soil component if applicable. The overall limitation rating for the soil component is based on the most severe limitation.

Pond reservoir areas hold water behind a dam or embankment. Soils best suited to this use have low seepage potential in the upper 60 inches. The seepage potential is determined by the permeability of the soil and the depth to fractured bedrock or other permeable material. Excessive slope can affect the storage capacity of the reservoir area.

Embankments, dikes, and levees are raised structures of soil material, generally less than 20 feet high, constructed to impound water or to protect land against overflow. In this table, the soils are rated as a source of material for embankment fill. The ratings apply to the soil material below the surface layer to a depth of about 5 feet. It is assumed that soil layers will be uniformly mixed and compacted during construction.

The ratings do not indicate the ability of the natural soil to support an embankment. Soil properties to a depth even greater than the height of the embankment can affect performance and safety of the embankment. Generally, deeper onsite investigation is needed to determine these properties.

Soil material in embankments must be resistant to seepage, piping, and erosion and have favorable compaction characteristics. Unfavorable features include less than 5 feet of suitable material and a high content of stones or boulders, organic matter, or salts or sodium. A high water table affects the amount of usable material. It also affects traffic ability.

Aquifer-fed excavated ponds are pits or dugouts that extend to a ground-water aquifer or to a depth below a permanent water table. Excluded are ponds that are fed only by surface runoff and embankment ponds that impound water 3 feet or more above the original surface. Excavated ponds are affected by depth to a permanent water table, permeability of the aquifer, and quality of the water as inferred from the salinity of the soil. Depth to bedrock and the content of large stones affect the ease of excavation.

Drainage is the removal of excess surface and subsurface water from the soil. How easily and effectively the soil is drained depends on the depth to bedrock, to a cemented pan, or to other layers that affect the rate of water movement; permeability; depth to a high water table or depth of standing water if the soil is subject to ponding; slope; susceptibility to flooding; subsidence of organic layers; and the potential for frost action. Excavating and grading and the stability of ditch banks are affected by depth to bedrock or to a cemented pan, large stones, slope, and the hazard of cutbanks caving. The productivity of the soil after drainage is adversely affected by extreme acidity or by toxic substances in the root zone, such as salts, sodium, and sulfur. Availability of drainage outlets is not considered in the ratings.

Irrigation is the controlled application of water to supplement rainfall and support plant growth. The design and management of an irrigation system are affected by depth to the water table, the need for drainage, flooding, available water capacity, intake rate, permeability, erosion hazard, and slope. The construction of a system is affected by large stones and depth to bedrock or to a cemented pan. The performance of a system is affected by the depth of the root zone, the amount of salts or sodium, and soil reaction.

Terraces and diversions are embankments or a combination of channels and ridges constructed across a slope to control erosion and conserve moisture by intercepting runoff. Slope, wetness, large stones, and depth to bedrock or to a cemented pan affect the construction of terraces and diversions. A restricted rooting depth, a very limited hazard of wind erosion or water erosion, an excessively coarse texture, and restricted permeability adversely affect maintenance.

Grassed waterways are natural or constructed channels, generally broad and shallow, which conduct surface water to outlets at a non-erosive velocity. Large stones, wetness, slope, and depth to bedrock or to a cemented pan affect the construction of grassed waterways. A hazard of wind erosion, low available water capacity, restricted rooting depth, toxic substances such as salts and sodium, and restricted permeability adversely affect the growth and maintenance of the grass after construction.

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

| Map symbol and soil name | Features affecting-- | | | |
|--------------------------|--|---|---|---|
| | Drainage | Irrigation | Terraces and diversions | Grassed waterways |
| Ad: Alda----- | Limitation: frost action cutbanks cave | Limitation: wetness | Limitation: too sandy wetness | Favorable |
| Bo: Boel----- | Limitation: cutbanks cave | Limitation: wetness droughty | Limitation: too sandy wetness soil blowing | Limitation: rooting depth droughty |
| Bu: Butler----- | Limitation: frost action percs slowly | Limitation: erodes easily percs slowly wetness | Limitation: erodes easily wetness | Limitation: erodes easily percs slowly wetness |
| CaC: Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily | Limitation: erodes easily |
| CaD: Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| CaF: Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| CkB: Coly----- | Limitation: deep to water | Limitation: erodes easily | Limitation: erodes easily | Limitation: erodes easily |
| Kenesaw----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| CoD2: Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| Uly----- | Limitation: deep to water | Limitation: slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| CoF2: Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| Uly----- | Limitation: deep to water | Limitation: slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| De: Detroit----- | Limitation: deep to water | Limitation: erodes easily percs slowly | Limitation: erodes easily percs slowly | Limitation: erodes easily percs slowly |
| Fm: Fillmore----- | Limitation: frost action percs slowly ponding | Limitation: erodes easily percs slowly ponding | Limitation: erodes easily percs slowly ponding | Limitation: erodes easily percs slowly wetness |
| Gb: Gibbon----- | Limitation: frost action | Limitation: wetness | Limitation: wetness | Favorable |
| Gc: Gibbon----- | Limitation: excess salt frost action | Limitation: excess salt wetness | Limitation: erodes easily wetness | Limitation: erodes easily excess salt |
| Go: Gothenburg----- | Limitation: flooding cutbanks cave | Limitation: fast intake wetness droughty | Limitation: too sandy wetness | Limitation: wetness droughty |
| GP: Pits----- | Limitation: deep to water | Limitation: fast intake slope droughty | Limitation: slope too sandy | Limitation: rooting depth slope droughty |
| HeB: Hersh----- | Limitation: deep to water | Limitation: soil blowing | Limitation: soil blowing | Favorable |
| HeC: Hersh----- | Limitation: deep to water | Limitation: slope soil blowing | Limitation: soil blowing | Favorable |
| HeD: Hersh----- | Limitation: deep to water | Limitation: slope soil blowing | Limitation: slope soil blowing | Limitation: slope |
| Hf: Hobbs----- | Limitation: deep to water | Limitation: flooding | Limitation: erodes easily | Limitation: erodes easily |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

| Map symbol and soil name | Features affecting-- | | | |
|-------------------------------------|---|---|---|---|
| | Drainage | Irrigation | Terraces and diversions | Grassed waterways |
| HgB: Hobbs----- | Limitation: deep to water | Limitation: flooding | Limitation: erodes easily | Limitation: erodes easily |
| Ho: Holdrege----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| HoB: Holdrege----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| HoC: Holdrege----- | Limitation: deep to water | Limitation: slope | Limitation: erodes easily | Limitation: erodes easily |
| HoC2: Holdrege----- | Limitation: deep to water | Limitation: slope | Limitation: erodes easily | Limitation: erodes easily |
| Hp: Holdrege----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| HpB: Holdrege----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| Hr: Hord----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| InB: Inavale----- | Limitation: deep to water | Limitation: fast intake droughty | Limitation: too sandy soil blowing | Limitation: droughty |
| Ke: Kenesaw----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| KgB: Kenesaw----- | Limitation: deep to water | Favorable | Limitation: erodes easily | Limitation: erodes easily |
| Coly----- | Limitation: deep to water | Limitation: erodes easily | Limitation: erodes easily | Limitation: erodes easily |
| Lf: Lex----- | Limitation: frost action cutbanks cave | Limitation: wetness | Limitation: erodes easily too sandy wetness | Limitation: erodes easily wetness |
| Lg: Lex----- | Limitation: excess salt frost action cutbanks cave | Limitation: excess salt wetness | Limitation: erodes easily too sandy wetness | Limitation: erodes easily excess salt |
| LoB: Libory----- | Favorable | Limitation: wetness droughty | Limitation: erodes easily wetness soil blowing | Limitation: erodes easily droughty |
| M-W: Miscellaneous Water----- | --- | --- | --- | --- |
| Ma: Massie----- | Limitation: frost action percs slowly ponding | Limitation: ponding | Limitation: erodes easily percs slowly ponding | Limitation: erodes easily wetness |
| Pm: Platte----- | Limitation: flooding cutbanks cave | Limitation: wetness droughty | Limitation: too sandy wetness | Limitation: rooting depth wetness droughty |
| Ru: Rusco----- | Limitation: frost action ponding | Limitation: ponding | Limitation: erodes easily ponding | Limitation: erodes easily wetness |
| Sc: Scott----- | Limitation: frost action percs slowly ponding | Limitation: erodes easily percs slowly ponding | Limitation: erodes easily percs slowly ponding | Limitation: erodes easily percs slowly wetness |
| SmB: Simeon----- | Limitation: deep to water | Limitation: droughty | Limitation: too sandy soil blowing | Limitation: droughty |
| To: Tryon----- | Limitation: cutbanks cave | Limitation: fast intake wetness droughty | Limitation: too sandy wetness | Limitation: wetness droughty |
| UcF: Uly----- | Limitation: deep to water | Limitation: slope | Limitation: erodes easily slope | Limitation: erodes easily slope |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this report indicates the dominant soil condition but does not eliminate the need for onsite investigation)

| Map symbol and soil name | Features affecting-- | | | |
|--------------------------|------------------------------|---|---|---|
| | Drainage | Irrigation | Terraces and diversions | Grassed waterways |
| Coly----- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| UsF: Ustorthents---- | Limitation: deep to water | Limitation: erodes easily slope | Limitation: erodes easily slope | Limitation: erodes easily slope |
| VaB: Valentine----- | Limitation: deep to water | Limitation: fast intake droughty | Limitation: too sandy soil blowing | Limitation: rooting depth droughty |
| VaD: Valentine----- | Limitation: deep to water | Limitation: fast intake slope droughty | Limitation: too sandy soil blowing | Limitation: rooting depth droughty |
| VaF: Valentine----- | Limitation: deep to water | Limitation: fast intake slope droughty | Limitation: slope too sandy soil blowing | Limitation: rooting depth slope droughty |
| VbD: Valentine----- | Limitation: deep to water | Limitation: fast intake slope droughty | Limitation: too sandy soil blowing | Limitation: rooting depth droughty |
| Els----- | Limitation: cutbanks cave | Limitation: fast intake wetness droughty | Limitation: too sandy wetness soil blowing | Limitation: droughty |
| W: Water----- | --- | --- | --- | --- |
| Wa: Wann----- | Limitation: frost action | Limitation: wetness soil blowing | Limitation: wetness soil blowing | Favorable |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Pond Reservoir Area | | Embankments, Dikes, and Levees | | Excavated Ponds (Aquifer- fed) | |
|-----------------------------|--------------------------|---------------------------------------|--------------|--|----------------------|--|------------------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Seepage | 1.00 | Somewhat limited Seepage Depth to saturated zone Piping | 0.98 0.95 0.02 | Very limited Cutbanks cave Deep to water | 1.00 0.02 |
| Bo: Boel----- | 100 | Very limited Seepage | 1.00 | Very limited Seepage Depth to saturated zone | 1.00 0.95 | Very limited Cutbanks cave Deep to water | 1.00 0.02 |
| Bu: Butler----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Depth to saturated zone | 1.00 | Very limited Deep to water | 1.00 |
| CaC: Coly----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| CaD: Coly----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| CaF: Coly----- | 100 | Somewhat limited Seepage Slope | 0.70 0.04 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| CkB: Coly----- | 60 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Kenesaw----- | 40 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| CoD2: Coly----- | 60 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Uly----- | 40 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| CoF2: Coly----- | 60 | Somewhat limited Seepage Slope | 0.70 0.04 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Uly----- | 40 | Somewhat limited Seepage Slope | 0.70 0.04 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| De: Detroit----- | 100 | Somewhat limited Seepage | 0.05 | Not limited | | Very limited Deep to water | 1.00 |
| Fm: Fillmore----- | 100 | Somewhat limited Seepage | 0.57 | Very limited Depth to saturated zone Hard to pack | 1.00 0.68 | Very limited Deep to water | 1.00 |
| Gb: Gibbon----- | 100 | Very limited Seepage | 1.00 | Very limited Piping Depth to saturated zone | 1.00 0.95 | Somewhat limited Cutbanks cave Deep to water | 0.10 0.02 |
| Gc: Gibbon----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Depth to saturated zone Salinity Piping | 0.95 0.12 0.07 | Somewhat limited Slow refill Cutbanks cave Salty water Deep to water | 0.30 0.10 0.06 0.02 |
| Go: Gothenburg----- | 100 | Very limited | | Very limited | | Very limited | |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Pond Reservoir Area | | Embankments, Dikes, and Levees | | Excavated Ponds (Aquifer- fed) | |
|-----------------------------|--------------------------|---------------------------------------|-------|--|--------------|---------------------------------------|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| GP: Pits----- | 100 | Seepage | 1.00 | Depth to saturated zone Seepage | 1.00 1.00 | Cutbanks cave | 1.00 |
| | | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Very limited Seepage | 1.00 | Very limited Piping Seepage | 1.00 0.09 | Very limited Deep to water | 1.00 |
| HeC: Hersh----- | 100 | Very limited Seepage | 1.00 | Very limited Piping Seepage | 1.00 0.09 | Very limited Deep to water | 1.00 |
| HeD: Hersh----- | 100 | Very limited Seepage | 1.00 | Very limited Piping Seepage | 1.00 0.09 | Very limited Deep to water | 1.00 |
| Hf: Hobbs----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.71 | Very limited Deep to water | 1.00 |
| HgB: Hobbs----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.71 | Very limited Deep to water | 1.00 |
| Ho: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.67 | Very limited Deep to water | 1.00 |
| HoB: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.67 | Very limited Deep to water | 1.00 |
| HoC: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.67 | Very limited Deep to water | 1.00 |
| HoC2: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.67 | Very limited Deep to water | 1.00 |
| Hp: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.44 | Very limited Deep to water | 1.00 |
| HpB: Holdrege----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.44 | Very limited Deep to water | 1.00 |
| Hr: Hord----- | 100 | Somewhat limited Seepage | 0.70 | Somewhat limited Piping | 0.59 | Very limited Deep to water | 1.00 |
| InB: Inavale----- | 100 | Very limited Seepage | 1.00 | Somewhat limited Seepage | 0.98 | Very limited Deep to water | 1.00 |
| Ke: Kenesaw----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| KgB: Kenesaw----- | 55 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Coly----- | 45 | Somewhat limited Seepage | 0.70 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Lf: Lex----- | 100 | Very limited Seepage | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Cutbanks cave | 1.00 |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Pond Reservoir Area | | Embankments, Dikes, and Levees | | Excavated Ponds (Aquifer-fed) | |
|------------------------------|-----------------|--------------------------------------|--------------|---|-------|------------------------------------|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Lg: Lex----- | 100 | Very limited Seepage | 1.00 | Seepage | 0.91 | Deep to water | 0.00 |
| | | | | Somewhat limited Depth to saturated zone Seepage | 0.95 | Very limited Cutbanks cave | 1.00 |
| | | | | Piping | 0.91 | Salty water | 0.50 |
| | | | | Salinity | 0.22 | Deep to water | 0.02 |
| | | | | | 0.12 | | |
| LoB: Libory----- | 100 | Very limited Seepage | 1.00 | Somewhat limited Piping | 0.99 | Very limited Cutbanks cave | 1.00 |
| | | | | Depth to saturated zone Seepage | 0.95 | Slow refill | 0.30 |
| | | | | | 0.62 | Deep to water | 0.02 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Not limited | | Very limited Depth to saturated zone Hard to pack | 1.00 | Very limited Deep to water | 1.00 |
| | | | | | 0.72 | | |
| Pm: Platte----- | 100 | Very limited Seepage | 1.00 | Very limited Depth to saturated zone Seepage | 1.00 | Very limited Cutbanks cave | 1.00 |
| | | | | | 0.97 | Deep to water | 0.00 |
| Ru: Rusco----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Depth to saturated zone Piping | 1.00 | Somewhat limited Slow refill | 0.30 |
| | | | | | 0.61 | Cutbanks cave | 0.10 |
| Sc: Scott----- | 100 | Somewhat limited Seepage | 0.70 | Very limited Depth to saturated zone Hard to pack | 1.00 | Very limited Deep to water | 1.00 |
| | | | | | 0.32 | | |
| SmB: Simeon----- | 100 | Very limited Seepage | 1.00 | Somewhat limited Seepage | 0.91 | Very limited Deep to water | 1.00 |
| To: Tryon----- | 100 | Very limited Seepage | 1.00 | Very limited Depth to saturated zone Seepage | 1.00 | Very limited Cutbanks cave | 1.00 |
| | | | | | 1.00 | | |
| UcF: Uly----- | 60 | Somewhat limited Seepage Slope | 0.70 0.04 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| Coly----- | 40 | Somewhat limited Seepage Slope | 0.70 0.04 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| UsF: Ustorthents----- | 100 | Somewhat limited Seepage Slope | 0.70 0.24 | Very limited Piping | 1.00 | Very limited Deep to water | 1.00 |
| VaB: Valentine----- | 100 | Very limited Seepage | 1.00 | Very limited Seepage | 1.00 | Very limited Deep to water | 1.00 |
| VaD: Valentine----- | 100 | Very limited Seepage | 1.00 | Very limited Seepage | 1.00 | Very limited Deep to water | 1.00 |
| VaF: Valentine----- | 100 | Very limited | | Very limited | | Very limited | |

WATER MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Pond Reservoir Area | | Embankments, Dikes, and Levees | | Excavated Ponds (Aquifer-fed) | |
|--------------------------|-----------------|------------------------------------|--------------|--|----------------------|---|--------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| | | Seepage Slope | 1.00 0.01 | Seepage | 1.00 | Deep to water | 1.00 |
| VbD: Valentine----- | 65 | Very limited Seepage | 1.00 | Very limited Seepage | 1.00 | Very limited Deep to water | 1.00 |
| Els----- | 35 | Very limited Seepage | 1.00 | Very limited Seepage Depth to saturated zone | 1.00 0.95 | Very limited Cutbanks cave Deep to water | 1.00 0.02 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Very limited Seepage | 1.00 | Somewhat limited Depth to saturated zone Seepage Piping | 0.84 0.10 0.02 | Very limited Cutbanks cave Deep to water | 1.00 0.07 |

SANITARY FACILITIES
Kearney County, Nebraska

Sanitary Facilities

The following tables show the degree and kind of soil limitations that affect septic tank absorption fields, sewage lagoons, sanitary landfills, and daily cover for landfill. The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect these uses. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Septic tank absorption fields are areas in which effluent from a septic tank is distributed into the soil through subsurface tiles or perforated pipe. Only that part of the soil between depths of 24 and 60 inches is evaluated. The ratings are based on the soil properties that affect absorption of the effluent, construction and maintenance of the system, and public health. Permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, and flooding affect absorption of the effluent. Stones and boulders, ice, and bedrock or a cemented pan interfere with installation. Subsidence interferes with installation and maintenance. Excessive slope may cause lateral seepage and surfacing of the effluent in downslope areas.

Some soils are underlain by loose sand and gravel or fractured bedrock at a depth of less than 4 feet below the distribution lines. In these soils the absorption field may not adequately filter the effluent, particularly when the system is new. As a result, the ground water may become contaminated.

Sewage lagoons are shallow ponds constructed to hold sewage while aerobic bacteria decompose the solid and liquid wastes. Lagoons should have a nearly level floor surrounded by cut slopes or embankments of compacted soil. Nearly impervious soil material for the lagoon floor and sides is required to minimize seepage and contamination of ground water. Considered in the ratings are slope, permeability, depth to a water table, ponding, depth to bedrock or a cemented pan, flooding, large stones, and content of organic matter.

Soil permeability is a critical property affecting the suitability for sewage lagoons. Most porous soils eventually become sealed when they are used as sites for sewage lagoons. Until sealing occurs, however, the hazard of pollution is severe. Soils that have a permeability rate of more than 2 inches per hour are too porous for the proper functioning of sewage lagoons. In these soils, seepage of the effluent can result in contamination of the ground water. Ground-water contamination is also a hazard if fractured bedrock is within a depth of 40 inches, if the water table is high enough to raise the level of sewage in the lagoon, or if floodwater overtops the lagoon.

A high content of organic matter is detrimental to proper functioning of the lagoon because it inhibits aerobic activity. Slope, bedrock, and cemented pans can cause construction problems, and large stones can hinder compaction of the lagoon floor. If the lagoon is to be uniformly deep throughout, the slope must be gentle enough and the soil material must be thick enough over bedrock or a cemented pan to make land smoothing practical.

A trench sanitary landfill is an area where solid waste is placed in successive layers in an excavated trench. The waste is spread, compacted, and covered daily with a thin layer of soil excavated at the site. When the trench is full, a final cover of soil material at least 2 feet thick is placed over the landfill. The ratings in the table are based on the soil properties that affect the risk of pollution, the ease of excavation, trafficability, and revegetation. These properties include permeability, depth to bedrock or a cemented pan, depth to a water table, ponding, slope, flooding, texture, stones and boulders, highly organic layers, soil reaction, and content of salts and sodium. Unless otherwise stated, the ratings apply only to that part of the soil within a depth of about 6 feet. For deeper trenches, onsite investigation may be needed.

Hard, nonrippable bedrock, creviced bedrock, or highly permeable strata in or directly below the proposed trench bottom can affect the ease of excavation and the hazard of ground-water pollution. Slope affects construction of the trenches and the movement of surface water around the landfill. It also affects the construction and performance of roads in areas of the landfill.

Soil texture and consistence affect the ease with which the trench is dug and the ease with which the soil can be used as daily or final cover. They determine the workability of the soil when dry and when wet. Soils that are plastic and sticky when wet are difficult to excavate, grade, or compact and are difficult to place as a uniformly thick cover over a layer of refuse.

The soil material used as the final cover for a trench landfill should be suitable for plants. It should not have excess sodium or salts and should not be too acid. The surface layer generally has the best workability, the highest content of organic matter, and the best potential for plants. Material from the surface layer should be stockpiled for use as the final cover.

SANITARY FACILITIES
Kearney County, Nebraska

In an area sanitary landfill, solid waste is placed in successive layers on the surface of the soil. The waste is spread, compacted, and covered daily with a thin layer of soil from a source away from the site. A final cover of soil material at least 2 feet thick is placed over the completed landfill. The ratings in the table are based on the soil properties that affect trafficability and the risk of pollution. These properties include flooding, permeability, depth to a water table, ponding, slope, and depth to bedrock or a cemented pan.

Flooding is a serious problem because it can result in pollution in areas downstream from the landfill. If permeability is too rapid or if fractured bedrock, a fractured cemented pan, or the water table is close to the surface, the leachate can contaminate the water supply. Slope is a consideration because of the extra grading required to maintain roads in the steeper areas of the landfill. Also, leachate may flow along the surface of the soils in the steeper areas and cause difficult seepage problems.

Daily cover for landfill is the soil material that is used to cover compacted solid waste in an area sanitary landfill. The soil material is obtained offsite, transported to the landfill, and spread over the waste. The ratings in the table also apply to the final cover for a landfill. They are based on the soil properties that affect workability, the ease of digging, and the ease of moving and spreading the material over the refuse daily during wet and dry periods. These properties include soil texture, depth to a water table, ponding, rock fragments, slope, depth to bedrock or a cemented pan, reaction, and content of salts, sodium, or lime.

Loamy or silty soils that are free of large stones and excess gravel are the best cover for a landfill. Clayey soils may be sticky and difficult to spread; sandy soils are subject to wind erosion.

Slope affects the ease of excavation and of moving the cover material. Also, it can influence runoff, erosion, and reclamation of the borrow area.

After soil material has been removed, the soil material remaining in the borrow area must be thick enough over bedrock, a cemented pan, or the water table to permit revegetation. The soil material used as the final cover for a landfill should be suitable for plants. It should not have excess sodium, salts, or lime and should not be too acid.

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Septic tank absorption fields | | Sewage lagoons | |
|--------------------------|-----------------|---|-------|---|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 |
| Bo: Boel----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 |
| Bu: Butler----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| | | Depth to saturated zone | 1.00 | | |
| CaC: Coly----- | 100 | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Slope | 0.67 |
| | | | | Seepage | 0.50 |
| CaD: Coly----- | 100 | Somewhat limited Restricted permeability | 0.50 | Very limited Slope | 1.00 |
| | | Slope | 0.04 | Seepage | 0.50 |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| CkB: Coly----- | 60 | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| Kenesaw----- | 40 | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| CoD2: Coly----- | 60 | Somewhat limited Restricted permeability | 0.50 | Very limited Slope | 1.00 |
| | | Slope | 0.04 | Seepage | 0.50 |
| Uly----- | 40 | Somewhat limited Restricted permeability | 0.50 | Very limited Slope | 1.00 |
| | | Slope | 0.04 | Seepage | 0.50 |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| De: Detroit----- | 100 | Very limited Restricted permeability | 1.00 | Not limited | |
| Fm: Fillmore----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.32 |
| | | Depth to saturated zone | 1.00 | | |
| Gb: Gibbon----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Septic tank absorption fields | | Sewage lagoons | |
|-----------------------------|--------------------------|--|-------|--|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Gc: Gibbon----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Restricted permeability | 1.00 | Seepage | 0.50 |
| | | Flooding | 0.40 | Flooding | 0.40 |
| Go: Gothenburg----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| | | Depth to saturated zone | 1.00 | Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| GP: Pits----- | 100 | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Not limited | | Very limited Seepage | 1.00 |
| | | | | Slope | 0.00 |
| HeC: Hersh----- | 100 | Not limited | | Very limited Seepage | 1.00 |
| | | | | Slope | 0.67 |
| HeD: Hersh----- | 100 | Somewhat limited Slope | 0.04 | Very limited Seepage | 1.00 |
| | | | | Slope | 1.00 |
| Hf: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| | | | | Slope | 0.00 |
| Ho: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| HoB: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| HoC: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Slope | 0.67 |
| | | | | Seepage | 0.50 |
| HoC2: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Slope | 0.67 |
| | | | | Seepage | 0.50 |
| Hp: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| HpB: Holdrege----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| Hr: Hord----- | 100 | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Seepage | 0.50 |
| InB: Inavale----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Seepage | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 |
| Ke: Kenesaw----- | 100 | Somewhat limited | | Slope | 0.00 |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Septic tank absorption fields | | Sewage lagoons | |
|------------------------------|-----------------|--|-------|--------------------------------------|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| KgB: Kenesaw----- | 55 | Restricted permeability | 0.50 | Seepage | 0.50 |
| | | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| Coly----- | 45 | Somewhat limited Restricted permeability | 0.50 | Somewhat limited Seepage | 0.50 |
| | | | | Slope | 0.00 |
| Lf: Lex----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| | | Restricted permeability | 0.68 | Flooding | 0.40 |
| | | Flooding | 0.40 | | |
| Lg: Lex----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| | | Restricted permeability | 0.68 | Flooding | 0.40 |
| | | Flooding | 0.40 | | |
| LoB: Libory----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 0.56 |
| | | Restricted permeability | 0.50 | Slope | 0.00 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Restricted permeability | 1.00 | Not limited | |
| | | Depth to saturated zone | 1.00 | | |
| Pm: Platte----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| | | Depth to saturated zone | 1.00 | Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Restricted permeability | 1.00 | Seepage | 0.50 |
| | | | | | |
| Sc: Scott----- | 100 | Very limited Restricted permeability | 1.00 | Somewhat limited Seepage | 0.50 |
| | | Depth to saturated zone | 1.00 | | |
| SmB: Simeon----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Seepage | 1.00 |
| | | | | Slope | 0.00 |
| To: Tryon----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 |
| UcF: Uly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Septic tank absorption fields | | Sewage lagoons | |
|--------------------------|-----------------|--------------------------------------|-------|------------------------------------|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Coly----- | 40 | Restricted permeability | 0.50 | Seepage | 0.50 |
| | | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| | | Restricted permeability | 0.50 | Seepage | 0.50 |
| UsF: Ustorthents----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Restricted permeability | 0.50 | | |
| | | Very limited Filtering capacity | 1.00 | Very limited Seepage | 1.00 |
| VaD: Valentine----- | 100 | | | Slope | 0.00 |
| | | Very limited Filtering capacity | 1.00 | Very limited Seepage | 1.00 |
| VaF: Valentine----- | 100 | | | Slope | 0.91 |
| | | Very limited Filtering capacity | 1.00 | Very limited Slope | 1.00 |
| VbD: Valentine----- | 65 | Slope | 0.84 | Seepage | 1.00 |
| | | Very limited Filtering capacity | 1.00 | Very limited Seepage | 1.00 |
| Els----- | 35 | Very limited Slope | 0.91 | Slope | 0.91 |
| | | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Filtering capacity | 1.00 | Depth to saturated zone | 1.00 |
| W: Water----- | 100 | Not rated | | Slope | 0.00 |
| Wa: Wann----- | 100 | Not rated | | Not rated | |
| | | Very limited Depth to saturated zone | 1.00 | Very limited Seepage | 1.00 |
| | | Flooding | 0.40 | Depth to saturated zone | 1.00 |
| | | | | Flooding | 0.40 |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Trench sanitary landfill | | Area sanitary landfill | | Daily cover for landfill | |
|--------------------------|-----------------|---|------------------------------|--|----------------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Depth to saturated zone Seepage Too Sandy Flooding | 1.00 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.68 |
| Bo: Boel----- | 100 | Very limited Depth to saturated zone Seepage Too Sandy Flooding | 1.00 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.68 |
| Bu: Butler----- | 100 | Very limited Depth to saturated zone Too clayey | 1.00 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Too clayey Hard to compact Depth to saturated zone | 1.00 1.00 |
| CaC: Coly----- | 100 | Not limited | | Not limited | | Not limited | |
| CaD: Coly----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| CkB: Coly----- | 60 | Not limited | | Not limited | | Not limited | |
| Kenesaw----- | 40 | Not limited | | Not limited | | Not limited | |
| Cod2: Coly----- | 60 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 |
| Uly----- | 40 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| De: Detroit----- | 100 | Somewhat limited Too clayey | 0.50 | Not limited | | Very limited Hard to compact Too clayey | 1.00 0.50 |
| Fm: Fillmore----- | 100 | Very limited Depth to saturated zone Too clayey | 1.00 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Too clayey Hard to compact Depth to saturated zone | 1.00 1.00 |
| Gb: Gibbon----- | 100 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Somewhat limited Depth to saturated zone Seepage | 0.68 0.21 |
| Gc: Gibbon----- | 100 | Very limited Depth to saturated zone Flooding | 1.00 0.40 | Very limited Depth to saturated zone Flooding | 1.00 0.40 | Somewhat limited Depth to saturated zone | 0.68 |
| Go: Gothenburg----- | 100 | Very limited Flooding Depth to saturated zone Seepage Too Sandy | 1.00 1.00 1.00 1.00 | Very limited Flooding Depth to saturated zone Seepage | 1.00 1.00 1.00 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 1.00 |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Very limited | | Very limited | | Somewhat limited | |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Trench sanitary landfill | | Area sanitary landfill | | Daily cover for landfill | |
|------------------------------|-----------------|---|------------------------------|---|----------------------|---|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HeC: Hersh----- | 100 | Seepage | 1.00 | Seepage | 1.00 | Seepage | 0.50 |
| HeD: Hersh----- | 100 | Very limited Seepage | 1.00 | Very limited Seepage | 1.00 | Somewhat limited Seepage | 0.50 |
| Hf: Hobbs----- | 100 | Very limited Seepage Slope | 1.00 0.04 | Very limited Seepage Slope | 1.00 0.04 | Somewhat limited Seepage Slope | 0.50 0.04 |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 | Very limited Hard to compact | 1.00 |
| Ho: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HoB: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HoC: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HoC2: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| Hp: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| HpB: Holdrege----- | 100 | Not limited | | Not limited | | Not limited | |
| Hr: Hord----- | 100 | Not limited | | Not limited | | Not limited | |
| InB: Inavale----- | 100 | Very limited Seepage Too Sandy Flooding | 1.00 1.00 0.40 | Very limited Seepage Flooding | 1.00 0.40 | Very limited Too Sandy Seepage | 1.00 1.00 |
| Ke: Kenesaw----- | 100 | Not limited | | Not limited | | Not limited | |
| KgB: Kenesaw----- | 55 | Not limited | | Not limited | | Not limited | |
| Coly----- | 45 | Not limited | | Not limited | | Not limited | |
| Lf: Lex----- | 100 | Very limited Depth to saturated zone Seepage Too Sandy Flooding | 1.00 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.86 |
| Lg: Lex----- | 100 | Very limited Depth to saturated zone Seepage Too Sandy Flooding | 1.00 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.68 |
| LoB: Libory----- | 100 | Somewhat limited Depth to saturated zone | 0.95 | Very limited Seepage Depth to saturated zone | 1.00 0.44 | Somewhat limited Depth to saturated zone | 0.68 |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Depth to saturated zone Too clayey | 1.00 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone Too clayey Hard to compact | 1.00 1.00 |
| Pm: Platte----- | 100 | Very limited Flooding Depth to saturated zone Seepage Too Sandy | 1.00 1.00 1.00 | Very limited Flooding Depth to saturated zone Seepage | 1.00 1.00 1.00 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.86 |
| Ru: Rusco----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |

SANITARY FACILITIES--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.01 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Trench sanitary landfill | | Area sanitary landfill | | Daily cover for landfill | |
|--------------------------|-----------------|---|------------------------------|--|----------------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Sc: Scott----- | 100 | Very limited Depth to saturated zone Too clayey | 1.00 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone Too clayey Hard to compact | 1.00 1.00 1.00 |
| SmB: Simeon----- | 100 | Very limited Seepage Too Sandy | 1.00 1.00 | Very limited Seepage | 1.00 | Very limited Too Sandy Seepage | 1.00 1.00 |
| To: Tryon----- | 100 | Very limited Depth to saturated zone Seepage Too Sandy Flooding | 1.00 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 1.00 |
| UcF: Uly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| Coly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| UsF: Ustorthents----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Slope | 1.00 |
| VaB: Valentine----- | 100 | Very limited Seepage Too Sandy | 1.00 1.00 | Very limited Seepage | 1.00 | Very limited Too Sandy Seepage | 1.00 1.00 |
| VaD: Valentine----- | 100 | Very limited Seepage Too Sandy | 1.00 1.00 | Very limited Seepage | 1.00 | Very limited Too Sandy Seepage | 1.00 1.00 |
| VaF: Valentine----- | 100 | Very limited Seepage Too Sandy Slope | 1.00 1.00 0.84 | Very limited Seepage Slope | 1.00 0.84 | Very limited Too Sandy Seepage Slope | 1.00 1.00 0.84 |
| VbD: Valentine----- | 65 | Very limited Seepage Too Sandy | 1.00 1.00 | Very limited Seepage | 1.00 | Very limited Too Sandy Seepage | 1.00 1.00 |
| Els----- | 35 | Very limited Depth to saturated zone Seepage Too Sandy | 1.00 1.00 1.00 | Very limited Depth to saturated zone Seepage | 1.00 1.00 | Very limited Too Sandy Seepage Depth to saturated zone | 1.00 1.00 0.68 |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Very limited Depth to saturated zone Seepage Flooding | 1.00 1.00 0.40 | Somewhat limited Seepage Depth to saturated zone | 0.50 0.44 |

AGRICULTURAL WASTE MANAGEMENT
Kearney County, Nebraska

The nature of the soil is also important in the application of organic wastes and wastewater to land as fertilizers and irrigation; it is also important when the soil is used as a medium for treatment and disposal of these wastes. Favorable soil properties are required to prevent environmental damage.

The use of organic wastes and wastewater as production resources will result in energy conservation, prevent the waste of these important resources, and prevent problems associated with their disposal. Where disposal is the goal, and a maximum amount is disposed in a minimum area to hold costs to a minimum, risk of environmental damage is the principal constraint. Where the reuse goal is pursued, and a minimum amount is applied to a maximum area to obtain the greatest benefit, environmental damage is unlikely.

Interpretations developed for waste management may include ratings for (1) manure and food processing wastes; (2) municipal sewage sludge; (3) irrigation use of wastewater; or (4) treatment of wastewater by the slow rate process, overland flow process, or rapid infiltration process. If available, these should be located in this subsection.

Soil properties are important considerations in areas where soils are used as sites for the treatment and disposal of organic waste and wastewater. Selection of soils with properties that favor waste management can help to prevent environmental damage.

The Ag-Waste tables show the degree and kind of soil limitations affecting the treatment of agricultural waste, including municipal and food-processing wastewater and effluent from lagoons or storage ponds. Municipal wastewater is the waste stream from a municipality. It contains domestic waste and may contain industrial waste. It may have received primary or secondary treatment. It is rarely untreated sewage. Food-processing wastewater results from the preparation of fruits, vegetables, milk, cheese, and meats for public consumption. In places it is high in content of sodium and chloride. In the context of these tables, the effluent in lagoons and storage ponds is from facilities used to treat or store food-processing wastewater or domestic or animal waste. Domestic and food-processing wastewater is very dilute, and the effluent from the facilities that treat or store it commonly is very low in content of carbonaceous and nitrogenous material; the content of nitrogen commonly ranges from 10 to 30 milligrams per liter. The wastewater from animal waste treatment lagoons or storage ponds, however, has much higher concentrations of these materials, mainly because the manure has not been diluted as much as the domestic waste. The content of nitrogen in this wastewater generally ranges from 50 to 2,000 milligrams per liter. When wastewater is applied, checks should be made to ensure that nitrogen, phosphorus, heavy metals, and salts are not added in excessive amounts.

The ratings in the tables are for waste management systems that not only dispose of and treat organic waste or wastewater but also are beneficial to crops (application of manure and food-processing waste, application of sewage sludge, and disposal of wastewater by irrigation) and for waste management systems that are designed only for the purpose of wastewater disposal and treatment (overland flow of wastewater, rapid infiltration of wastewater, and slow rate treatment of wastewater).

The ratings are both verbal and numerical. Rating class terms indicate the extent to which the soils are limited by all of the soil features that affect agricultural waste management. Not limited indicates that the soil has features that are very favorable for the specified use. Good performance and very low maintenance can be expected. Slightly limited indicates that the soil has features that are generally favorable for the specified use. The limitations are minor and can be easily overcome. Good performance and low maintenance can be expected. Somewhat limited indicates that the soil has features that are moderately favorable for the specified use. The limitations can be overcome or minimized by special planning, design, or installation. Fair performance and moderate maintenance can be expected. Very limited indicates that the soil has one or more features that are unfavorable for the specified use. The limitations generally cannot be overcome without major soil reclamation, special design, or expensive installation procedures. Poor performance and high maintenance can be expected.

Numerical ratings in the tables indicate the severity of individual limitations. The ratings are shown as decimal fractions ranging from 0.00 to 1.00. They indicate gradations between the point at which a soil feature has the greatest negative impact on the use (1.00) and the point at which the soil feature is not a limitation (0.00).

Application of manure and food-processing waste not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. Manure is the excrement of livestock and poultry, and food-processing waste is damaged fruit and vegetables and the peelings, stems, leaves, pits, and soil particles removed in food preparation. The manure and food-processing waste are either solid, slurry, or liquid. Their nitrogen content varies. A high content of nitrogen limits the application rate. Toxic or otherwise dangerous wastes, such as those mixed with the lye used in food processing, are not considered in the ratings.

The ratings are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the waste is applied, and the method by which the waste is applied. The properties that affect absorption include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, and available water capacity. The properties that affect plant growth and microbial activity include reaction, the sodium adsorption ratio, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of waste. Permanently frozen soils are unsuitable for waste treatment.

Application of sewage sludge not only disposes of waste material but also can improve crop production by increasing the supply of nutrients in the soils where the material is applied. In the context of this table, sewage sludge is the residual product of the treatment of municipal sewage. The solid component consists mainly of cell mass, primarily bacteria cells that developed during secondary treatment and have incorporated soluble organics into their own bodies. The sludge has small amounts of sand, silt, and other solid debris. The content of nitrogen varies. Some sludge has constituents that are toxic to plants or hazardous to the food chain, such as heavy metals and exotic organic compounds, and should be analyzed chemically prior to use.

AGRICULTURAL WASTE MANAGEMENT
Kearney County, Nebraska

The content of water in the sludge ranges from about 98 percent to less than 40 percent. The sludge is considered liquid if it is more than about 90 percent water, slurry if it is about 50 to 90 percent water, and solid if it is less than about 50 percent water.

The ratings in the table are based on the soil properties that affect absorption, plant growth, microbial activity, erodibility, the rate at which the sludge is applied, and the method by which the sludge is applied. The properties that affect absorption, plant growth, and microbial activity include permeability, depth to a water table, ponding, the sodium adsorption ratio, depth to bedrock or a cemented pan, available water capacity, reaction, salinity, and bulk density. The wind erodibility group, the soil erodibility factor K, and slope are considered in estimating the likelihood that wind erosion or water erosion will transport the waste material from the application site. Stones, cobbles, a water table, ponding, and flooding can hinder the application of sludge. Permanently frozen soils are unsuitable for waste treatment.

Disposal of wastewater by irrigation not only disposes of municipal wastewater and wastewater from food-processing plants, lagoons, and storage ponds but also can improve crop production by increasing the amount of water available to crops. The ratings in the table are based on the soil properties that affect the design, construction, management, and performance of the irrigation system. The properties that affect design and management include the sodium adsorption ratio, depth to a water table, ponding, available water capacity, permeability, slope, and flooding. The properties that affect construction include stones, cobbles, depth to bedrock or a cemented pan, depth to a water table, and ponding.

The properties that affect performance include depth to bedrock or a cemented pan, bulk density, the sodium adsorption ratio, salinity, reaction, and the cation-exchange capacity, which is used to estimate the capacity of a soil to adsorb heavy metals. Permanently frozen soils are not suitable for disposal of wastewater by irrigation.

See the National Soil Handbook, September 1992, Part 620, for criteria used in rating soils for sanitary facilities and waste management.

AGRICULTURAL WASTE MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Application of manure and food-processing waste | | Application of sewage sludge | | Disposal of wastewater by irrigation | |
|--------------------------|-----------------|---|------------------------------|---|------------------------------|--|----------------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ad: Alda----- | 100 | Very limited Filtering capacity Depth to saturated zone Sodium content | 1.00 0.95 0.08 | Very limited Filtering capacity Depth to saturated zone Flooding Sodium content | 1.00 0.95 0.40 0.08 | Very limited Filtering capacity Depth to saturated zone Sodium content | 1.00 0.95 0.08 |
| Bo: Boel----- | 100 | Very limited Filtering capacity Depth to saturated zone Leaching limitation | 1.00 0.95 0.45 | Very limited Filtering capacity Depth to saturated zone Flooding | 1.00 0.95 0.40 | Very limited Filtering capacity Depth to saturated zone | 1.00 0.95 |
| Bu: Butler----- | 100 | Very limited Restricted permeability Depth to saturated zone Runoff limitation Too acid | 1.00 1.00 0.40 0.11 | Very limited Restricted permeability Depth to saturated zone Too acid | 1.00 1.00 0.42 | Very limited Restricted permeability Depth to saturated zone Too acid | 1.00 1.00 0.42 |
| CaC: Coly----- | 100 | Not limited | | Not limited | | Somewhat limited Too steep for surface application | 0.31 |
| CaD: Coly----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 0.22 |
| CaF: Coly----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 1.00 |
| CkB: Coly----- | 60 | Not limited | | Not limited | | Not limited | |
| Kenesaw----- | 40 | Not limited | | Not limited | | Not limited | |
| CoD2: Coly----- | 60 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 0.22 |
| Uly----- | 40 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 0.22 |
| CoF2: Coly----- | 60 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 1.00 |
| Uly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Too steep for surface application Too steep for sprinkler application | 1.00 1.00 |

AGRICULTURAL WASTE MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Application of manure and food-processing waste | | Application of sewage sludge | | Disposal of wastewater by irrigation | |
|--------------------------|-----------------|--|--------------|--|--------------|--|--------------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| De: Detroit----- | 100 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 |
| Fm: Fillmore----- | 100 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 |
| | | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 |
| | | Runoff limitation | 0.40 | Too acid | 0.14 | Too acid | 0.14 |
| | | Too acid | 0.03 | | | | |
| Gb: Gibbon----- | 100 | Somewhat limited Depth to saturated zone | 0.95 | Somewhat limited Depth to saturated zone Flooding | 0.95 0.40 | Somewhat limited Depth to saturated zone | 0.95 |
| Gc: Gibbon----- | 100 | Somewhat limited Depth to saturated zone Salinity | 0.95 0.50 | Very limited Salinity | 1.00 | Very limited Salinity | 1.00 |
| | | Restricted permeability | 0.30 | Depth to saturated zone Flooding | 0.95 0.40 | Depth to saturated zone | 0.95 |
| | | Sodium content | 0.08 | Restricted permeability Sodium content | 0.22 0.08 | Restricted permeability Sodium content | 0.22 0.08 |
| Go: Gothenburg----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Droughty | 1.00 | Very limited Droughty | 1.00 |
| | | Depth to saturated zone | 1.00 | Filtering capacity | 1.00 | Filtering capacity | 1.00 |
| | | Flooding | 1.00 | Depth to saturated zone Flooding | 1.00 1.00 | Depth to saturated zone Flooding | 1.00 1.00 |
| | | Droughty | 1.00 | | | | |
| | | Runoff limitation | 0.40 | | | | |
| GP: Pits----- | 100 | Not rated | | Not rated | | Not rated | |
| HeB: Hersh----- | 100 | Somewhat limited Filtering capacity | 0.00 | Somewhat limited Filtering capacity | 0.00 | Somewhat limited Filtering capacity | 0.00 |
| HeC: Hersh----- | 100 | Somewhat limited Filtering capacity | 0.00 | Somewhat limited Filtering capacity | 0.00 | Somewhat limited Too steep for surface application Filtering capacity | 0.31 0.00 |
| HeD: Hersh----- | 100 | Somewhat limited Slope | 0.04 | Somewhat limited Slope | 0.04 | Very limited Too steep for surface application | 1.00 |
| | | Filtering capacity | 0.00 | Filtering capacity | 0.00 | Too steep for sprinkler application Filtering capacity | 0.22 0.00 |
| Hf: Hobbs----- | 100 | Somewhat limited Flooding | 0.60 | Very limited Flooding | 1.00 | Somewhat limited Flooding | 0.60 |
| HgB: Hobbs----- | 100 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 | Very limited Flooding | 1.00 |
| Ho: Holdrege----- | 100 | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Restricted permeability | 0.31 |
| HoB: Holdrege----- | 100 | Somewhat limited | | Somewhat limited | | Somewhat limited | |

AGRICULTURAL WASTE MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Application of manure and food-processing waste | | Application of sewage sludge | | Disposal of wastewater by irrigation | |
|------------------------------|-----------------|---|-------|--|-------|--|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| HoC: Holdrege----- | 100 | Restricted permeability | 0.41 | Restricted permeability | 0.31 | Restricted permeability | 0.31 |
| | | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Too steep for surface application Restricted permeability | 0.31 |
| HoC2: Holdrege----- | 100 | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Too steep for surface application Restricted permeability | 0.31 |
| | | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Restricted permeability | 0.31 |
| Hp: Holdrege----- | 100 | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Restricted permeability | 0.31 |
| HpB: Holdrege----- | 100 | Somewhat limited Restricted permeability | 0.41 | Somewhat limited Restricted permeability | 0.31 | Somewhat limited Restricted permeability | 0.31 |
| Hr: Hord----- | 100 | Not limited | | Not limited | | Not limited | |
| InB: Inavale----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Leaching limitation | 0.45 | Flooding | 0.40 | Droughty | 0.19 |
| | | Droughty | 0.19 | Droughty | 0.19 | | |
| Ke: Kenesaw----- | 100 | Not limited | | Not limited | | Not limited | |
| KgB: Kenesaw----- | 55 | Not limited | | Not limited | | Not limited | |
| Coly----- | 45 | Not limited | | Not limited | | Not limited | |
| Lf: Lex----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 |
| | | Flooding | 0.40 | Flooding | 0.40 | | |
| Lg: Lex----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Depth to saturated zone | 0.95 | Salinity | 1.00 | Salinity | 1.00 |
| | | Salinity | 0.50 | Depth to saturated zone | 0.95 | Depth to saturated zone | 0.95 |
| | | Sodium content | 0.32 | Flooding Sodium content | 0.40 | Sodium content | 0.32 |
| LoB: Libory----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Depth to saturated zone | 0.95 | Depth to saturated zone | 0.95 | Depth to saturated zone | 0.95 |
| | | Leaching limitation | 0.45 | | | | |
| M-W: Miscellaneous Water- | 100 | Not rated | | Not rated | | Not rated | |
| Ma: Massie----- | 100 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 |
| | | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 |
| | | Runoff limitation | 0.40 | Too acid | 0.42 | Too acid | 0.42 |
| Pm: Platte----- | 100 | Very limited | 0.11 | Very limited | | Very limited | |

AGRICULTURAL WASTE MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Application of manure and food-processing waste | | Application of sewage sludge | | Disposal of wastewater by irrigation | |
|--------------------------|-----------------|---|-------|--------------------------------------|-------|--|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| Ru: Rusco----- | 100 | Filtering capacity | 1.00 | Filtering capacity | 1.00 | Filtering capacity | 1.00 |
| | | Depth to dense layer | 1.00 | Flooding | 1.00 | Depth to saturated zone | 1.00 |
| | | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 | Droughty | 0.86 |
| | | Droughty | 0.86 | Droughty | 0.86 | Flooding | 0.60 |
| | | Flooding | 0.60 | | | | |
| Sc: Scott----- | 100 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Restricted permeability | 0.30 | Restricted permeability | 0.22 | Restricted permeability | 0.22 |
| | | | | | | | |
| SmB: Simeon----- | 100 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 | Very limited Restricted permeability | 1.00 |
| | | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 | Depth to saturated zone | 1.00 |
| | | Runoff limitation | 0.40 | Too acid | 0.14 | Too acid | 0.14 |
| | | Too acid | 0.03 | | | | |
| To: Tryon----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Leaching limitation | 0.45 | Droughty | 0.13 | Droughty | 0.13 |
| | | Droughty | 0.13 | | | | |
| | | | | | | | |
| UcF: Uly----- | 60 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 | Very limited Depth to saturated zone | 1.00 |
| | | Filtering capacity | 1.00 | Filtering capacity | 1.00 | Filtering capacity | 1.00 |
| | | Droughty | 0.55 | Droughty | 0.55 | Droughty | 0.55 |
| | | Runoff limitation | 0.40 | Flooding | 0.40 | | |
| Coly----- | 40 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Too steep for surface application | 1.00 |
| | | | | | | Too steep for sprinkler application | 1.00 |
| UsF: Ustorthents----- | 100 | Very limited Slope | 1.00 | Very limited Slope | 1.00 | Very limited Too steep for surface application | 1.00 |
| | | | | | | Too steep for sprinkler application | 1.00 |
| | | | | | | | |
| VaB: Valentine----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Droughty | 0.77 | Droughty | 0.77 | Droughty | 0.77 |
| | | Leaching limitation | 0.45 | | | | |
| VaD: Valentine----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Droughty | 0.77 | Droughty | 0.77 | Droughty | 0.77 |
| | | Leaching limitation | 0.45 | | | Too steep for surface application | 0.66 |
| | | | | | | Too steep for sprinkler application | 0.00 |

AGRICULTURAL WASTE MANAGEMENT--Continued
Kearney County, Nebraska

(The information in this table indicates the dominant soil condition but does not eliminate the need for onsite investigation. The numbers in the value columns range from 0.00 to 1.00. The larger the value, the greater the potential limitation. See text for further explanation of ratings in this table.)

| Map symbol and soil name | Pct of map unit | Application of manure and food-processing waste | | Application of sewage sludge | | Disposal of wastewater by irrigation | |
|--------------------------|-----------------|---|-------|--|-------|--|-------|
| | | Rating class and limiting features | Value | Rating class and limiting features | Value | Rating class and limiting features | Value |
| VaF: Valentine----- | 100 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Too steep for surface application | 1.00 |
| | | Slope | 0.84 | Slope | 0.84 | Filtering capacity | 1.00 |
| | | Droughty | 0.77 | Droughty | 0.77 | Too steep for sprinkler application | 0.89 |
| | | Leaching limitation | 0.45 | | | Droughty | 0.77 |
| VbD: Valentine----- | 65 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Droughty | 0.77 | Droughty | 0.77 | Droughty | 0.77 |
| | | Leaching limitation | 0.45 | | | Too steep for surface application | 0.66 |
| | | | | | | Too steep for sprinkler application | 0.00 |
| Els----- | 35 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 | Very limited Filtering capacity | 1.00 |
| | | Depth to saturated zone | 0.95 | Depth to saturated zone | 0.95 | Depth to saturated zone | 0.95 |
| | | Droughty | 0.78 | Droughty | 0.78 | Droughty | 0.78 |
| | | Leaching limitation | 0.45 | | | | |
| W: Water----- | 100 | Not rated | | Not rated | | Not rated | |
| Wa: Wann----- | 100 | Somewhat limited Depth to saturated zone | 0.84 | Somewhat limited Depth to saturated zone | 0.84 | Somewhat limited Depth to saturated zone | 0.84 |
| | | Sodium content | 0.08 | Flooding | 0.40 | Sodium content | 0.08 |
| | | Filtering capacity | 0.00 | Sodium content | 0.08 | Filtering capacity | 0.00 |
| | | | | Filtering capacity | 0.00 | | |

In this section, hydric soils are defined and described and the hydric soils in the survey area are listed. The three essential characteristics of wetlands are hydrophytic vegetation, hydric soils, and wetland hydrology (Cowardin and others, 1979; U.S. Army Corps of Engineers, 1987; National Research Council, 1995; Tiner, 1985). Criteria for each of the characteristics must be met for areas to be identified as wetlands. Undrained hydric soils that have natural vegetation should support a dominant population of ecological wetland plant species. Hydric soils that have been converted to other uses should be capable of being restored to wetlands.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part (Federal Register, 1994). These soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

The NTCHS definition identifies general soil properties that are associated with wetness. In order to determine whether a specific soil is a hydric soil or nonhydric soil, however, more specific information, such as information about the depth and duration of the water table, is needed. Thus, criteria that identify those estimated soil properties unique to hydric soils have been established (Federal Register, 1995). These criteria are used to identify a phase of a soil series that normally is associated with wetlands. The criteria used are selected estimated soil properties that are described in "Soil Taxonomy" (USDA, 1999) and "Keys to Soil Taxonomy" (USDA, 1998) and in the "Soil Survey Manual" (USDA, 1993).

If soils are wet enough for a long enough period to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils in this survey area are specified in "Field Indicators of Hydric Soils in the United States" (Hurt and others, 1996).

Hydric soils are identified by examining and describing the soil to a depth of about 20 inches. This depth may be greater if determination of an appropriate indicator so requires. It is always recommended that soils be excavated and described to the depth necessary for an understanding of the redoximorphic processes. Then, using the completed soil descriptions, soil scientists can compare the soil features required by each indicator and specify which indicators have been matched with the conditions observed in the soil. The soil can be identified as a hydric soil if at least one of the approved indicators is present.

Map units in the Hydric Soil Interpretations table meet the definition of hydric soils and, in addition, have at least one of the hydric soil indicators. This list can help in planning land uses; however, onsite investigation is recommended to determine the hydric soils on a specific site (National Research Council, 1995; Hurt and others, 1996).

Map units that are made up of hydric soils may have small areas, or inclusions, of nonhydric soils in the higher positions on the landform, and map units made up of nonhydric soils may have inclusions of hydric soils in the lower positions on the landform.

These map units, in general, do not meet the definition of hydric soils because they do not have one of the hydric soil indicators. A portion of these map units, however, may include hydric soils. Onsite investigation is recommended to determine whether hydric soils occur and the location of the included hydric soils.

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Kearney County, Nebraska

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

| Map symbol and map unit name | Component | Hydric | Local landform | Hydric soils criteria | | | |
|---|---------------------|------------|-----------------------------|-----------------------|---------------------------|-------------------------|------------------------|
| | | | | Hydric criteria code | Meets saturation criteria | Meets flooding criteria | Meets ponding criteria |
| Ad: ALDA LOAM, 0 TO 1 PERCENT SLOPES | ALDA | No | flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| Bo: BOEL FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES | BOEL | No | flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| Bu: BUTLER SILT LOAM, 0 TO 1 PERCENT SLOPES | BUTLER | No | playa | --- | --- | --- | --- |
| | FILLMORE SCOTT | Yes Yes | playa playa | 2A 2B3,3 | YES YES | NO NO | NO YES |
| CaC: COLY SILT LOAM, 3 TO 6 PERCENT SLOPES | COLY | No | hillslope | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | flood plain | 2B3 | YES | NO | NO |
| CaD: COLY SILT LOAM, 6 TO 11 PERCENT SLOPES | COLY | No | hillslope | --- | --- | --- | --- |
| CaF: COLY SILT LOAM, 11 TO 20 PERCENT SLOPES | COLY | No | hillslope | --- | --- | --- | --- |
| CkB: COLY-KENESAW SILT LOAMS, 0 TO 3 PERCENT SLOPES | COLY | No | hillslope | --- | --- | --- | --- |
| | KENESAW FILLMORE | No Yes | hillslope playa | --- | --- | --- | --- |
| CoD2: COLY-ULY SILT LOAMS, 6 TO 11 PERCENT SLOPES, ERODED | COLY | No | hillslope | --- | --- | --- | --- |
| | ULY | No | hillslope | --- | --- | --- | --- |
| CoF2: COLY-ULY SILT LOAMS, 11 TO 20 PERCENT SLOPES, ERODED | COLY | No | hillslope | --- | --- | --- | --- |
| De: DETROIT SILT LOAM, 0 TO 1 PERCENT SLOPES | DETROIT | No | flat | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| Fm: FILLMORE SILT LOAM, 0 TO 1 PERCENT SLOPES | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| | SCOTT | Yes | playa | 2B3,3 | YES | NO | YES |
| Gb: GIBBON LOAM, 0 TO 1 PERCENT SLOPES | GIBBON | No | flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| Gc: GIBBON LOAM, SALINE, 0 TO 1 PERCENT SLOPES | GIBBON | No | flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| Go: GOTHENBURG LOAMY SAND, 0 TO 2 PERCENT SLOPES | GOTHENBURG | Yes | flood plain | 2B2 | YES | NO | NO |
| GP: GRAVEL PIT | PITS | Unranked | --- | --- | --- | --- | --- |
| HeB: HERSH FINE SANDY LOAM, 0 TO 3 PERCENT SLOPES | HERSH | No | hillslope | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| HeC: HERSH FINE SANDY LOAM, 3 TO 6 PERCENT SLOPES | HERSH | No | hillslope | --- | --- | --- | --- |
| HeD: HERSH FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES | HERSH | No | hillslope | --- | --- | --- | --- |
| Hf: HOBBS SILT LOAM, 0 TO 2 PERCENT SLOPES | HOBBS | No | drainageway, flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| HgB: HOBBS SILT LOAM, CHANNELED | HOBBS | No | drainageway, flood plain | --- | --- | --- | --- |

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Kearney County, Nebraska

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

| Map symbol and map unit name | Component | Hydric | Local landform | Hydric soils criteria | | | |
|--|----------------------|------------|---------------------------|-----------------------|---------------------------|-------------------------|------------------------|
| | | | | Hydric criteria code | Meets saturation criteria | Meets flooding criteria | Meets ponding criteria |
| Ho: HOLDREGE SILT LOAM, 0 TO 1 PERCENT SLOPES | HOLDREGE | No | broad interstream divide | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| HoB: HOLDREGE SILT LOAM, 1 TO 3 PERCENT SLOPES | HOLDREGE | No | hillslope | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| HoC: HOLDREGE SILT LOAM, 3 TO 6 PERCENT SLOPES | HOLDREGE | No | hillslope | --- | --- | --- | --- |
| HoC2: HOLDREGE SILT LOAM, 3 TO 6 PERCENT SLOPES, ERODED | HOLDREGE | No | hillslope | --- | --- | --- | --- |
| Hp: HOLDREGE SILT LOAM, OVERBLOWN, 0 TO 1 PERCENT SLOPES | HOLDREGE | No | interfluve | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| HpB: HOLDREGE SILT LOAM, OVERBLOWN, 1 TO 3 PERCENT SLOPES | HOLDREGE | No | hillslope | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| Hr: HORD SILT LOAM, 0 TO 1 PERCENT SLOPES | HORD | No | broad interstream divide | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| InB: INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES | INAVALE | No | flood plain | --- | --- | --- | --- |
| | TRYON | Yes | swale | 2B2 | YES | NO | NO |
| Ke: KENESAW SILT LOAM, 0 TO 1 PERCENT SLOPES | KENESAW | No | interfluve | --- | --- | --- | --- |
| | FILLMORE | Yes | playa | 2A | YES | NO | NO |
| | KENESAW | No | hillslope | --- | --- | --- | --- |
| KgB: KENESAW-COLY SILT LOAMS, 1 TO 3 PERCENT SLOPES | COLY | No | hillslope | --- | --- | --- | --- |
| | LEX | No | flood plain | --- | --- | --- | --- |
| Lf: LEX LOAM, 0 TO 1 PERCENT SLOPES | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| | LEX | No | flood plain | --- | --- | --- | --- |
| Lg: LEX LOAM, SALINE, 0 TO 1 PERCENT SLOPES | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |
| | LIBORY | No | hillslope, stream terrace | --- | --- | --- | --- |
| | MARSH WT AT 0-1 FOOT | Yes Yes | swale swale | 2B1,3 2B3 | YES YES | NO NO | YES NO |
| M-W: MISCELLANEOUS WATER (SEWAGE LAGOONS) | MISCELLANEOUS WATER | --- | --- | --- | --- | --- | --- |
| Ma: MASSIE SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES | MASSIE | Yes | playa | 2B3,3 | YES | NO | YES |
| Pm: PLATTE LOAM, 0 TO 1 PERCENT SLOPES | PLATTE | No | flood plain | --- | --- | --- | --- |
| | GOTHENBURG | Yes | flood plain | 2B2 | YES | NO | NO |
| Ru: RUSCO SILT LOAM, 0 TO 1 PERCENT SLOPES | RUSCO | No | depression | --- | --- | --- | --- |
| | PERCHED WT | Yes | playa | 2A | YES | NO | NO |
| Sc: SCOTT SILT LOAM, 0 TO 1 PERCENT SLOPES | SCOTT | Yes | playa | 2B3,3 | YES | NO | YES |
| Smb: SIMEON SANDY LOAM, 0 TO 3 PERCENT SLOPES | SIMEON | No | stream terrace | --- | --- | --- | --- |

HYDRIC SOIL INTERPRETATIONS
HYDRIC SOILS LIST
Kearney County, Nebraska

All mapunits are displayed regardless of hydric status and are listed in alpha-numeric order by mapunit symbol. The "Hydric Soils Criteria" columns indicate the conditions that caused the mapunit component to be classified as "Hydric" or "Non-Hydric". These criteria are defined in "Hydric Soils of the United States" (USDA Miscellaneous Publication No. 1491, June, 1991). See the "Criteria for Hydric Soils" endnote to determine the meaning of these columns. Spot symbols are footnoted at the end of the table.

| Map symbol and map unit name | Component | Hydric | Local landform | Hydric soils criteria | | | |
|---|-------------------|----------|--------------------------|-----------------------|---------------------------|-------------------------|------------------------|
| | | | | Hydric criteria code | Meets saturation criteria | Meets flooding criteria | Meets ponding criteria |
| To: TRYON LOAMY FINE SAND, 0 TO 1 PERCENT SLOPES | TRYON | Yes | flood plain, swale | 2B2 | YES | NO | NO |
| UcF: ULY-COLY SILT LOAMS, 11 TO 20 PERCENT SLOPES | ULY | No | hillslope | --- | --- | --- | --- |
| | COLY | No | hillslope | --- | --- | --- | --- |
| UsF: USTORTHENTS, STEEP | USTORTHENTS | Unranked | --- | --- | --- | --- | --- |
| VaB: VALENTINE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES | VALENTINE | No | hummock, sand sheet | --- | --- | --- | --- |
| | TRYON | Yes | swale | 2B2 | YES | NO | NO |
| VaD: VALENTINE LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES | VALENTINE | No | hummock, sand sheet | --- | --- | --- | --- |
| | TRYON | Yes | swale | 2B2 | YES | NO | NO |
| VaF: VALENTINE LOAMY FINE SAND, ROLLING | VALENTINE | No | hillslope, sand sheet | --- | --- | --- | --- |
| | TRYON | Yes | swale | 2B2 | YES | NO | NO |
| VbD: VALENTINE-ELS LOAMY FINE SANDS, 0 TO 9 PERCENT SLOPES | VALENTINE | No | hummock, sand sheet | --- | --- | --- | --- |
| | ELS | No | sand sheet, swale | --- | --- | --- | --- |
| | TRYON | Yes | swale | 2B2 | YES | NO | NO |
| W: WATER | WATER | Unranked | --- | --- | --- | --- | --- |
| Wa: WANN FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES | WANN | No | flood plain | --- | --- | --- | --- |
| | WT AT 0-1 FOOT | Yes | swale | 2B3 | YES | NO | NO |

FOOTNOTE: There may be small areas of included soils or miscellaneous areas that are significant to use and management of the soil; yet are too small to delineate on the soil map at the map's original scale. These may be designated as spot symbols and are defined in the published Soil Survey Report or the USDA-NRCS Technical Guide, Part II. Areas mapped as water or any map unit that contains one of the following conventional symbols is considered a hydric soil map unit: marshes or swamps; wet spots; depressions; streams, lakes and ponds.

1. All Histosols except Folists, or
2. Soils in Aquic suborders, great groups, or subgroups, Albolls suborder, Aquisalids, Pachic subgroups, or Cumulic subgroups that are:
 - a. Somewhat poorly drained with a water table equal to 0.0 foot (ft) from the surface during the growing season, or
 - b. poorly drained or very poorly drained and have either:
 - (1) water table equal to 0.0 ft during the growing season if textures are coarse sand, sand, or fine sand in all layers within 20 inches (in), or for other soils
 - (2) water table at less than or equal to 0.5 ft from the surface during the growing season if permeability is equal to or greater than 6.0 in/hour (h) in all layers within 20 in, or
 - (3) water table at less than or equal to 1.0 ft from the surface during the growing season if permeability is less than 6.0 in/h in any layer within 20 in, or
3. Soils that are frequently ponded for long duration or very long duration during the growing season, or
4. Soils that are frequently flooded for long duration or very long duration during the growing season.

HIGHLY ERODIBLE LANDS REPORT

Survey Area- KEARNEY COUNTY, NEBRASKA

| Map Symbol | Soil Mapunit Name | HEL Classifications | | |
|---------------|--|---------------------|-------|------------|
| | | C=30 | R=125 | wnd wat mu |
| Ad | ALDA LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Bo | BOEL FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Bu | BUTLER SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| CaC | COLY SILT LOAM, 3 TO 6 PERCENT SLOPES | 3 | 2 | 2 |
| CaD | COLY SILT LOAM, 6 TO 11 PERCENT SLOPES | 3 | 1 | 1 |
| CaF | COLY SILT LOAM, 11 TO 20 PERCENT SLOPES | 3 | 1 | 1 |
| CkB | COLY-KENESAW SILT LOAMS, 0 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| CoD2 | COLY-ULY SILT LOAMS, 6 TO 11 PERCENT SLOPES, ERODED | 3 | 2 | 2 |
| CoF2 | COLY-ULY SILT LOAMS, 11 TO 20 PERCENT SLOPES, ERODED | 3 | 1 | 1 |
| De | DETROIT SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Fm | FILLMORE SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Gb | GIBBON LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Gc | GIBBON LOAM, SALINE, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Go | GOTHENBURG LOAMY SAND, 0 TO 2 PERCENT SLOPES | 1 | 3 | 1 |
| HeB | HERSH FINE SANDY LOAM, 0 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| HeC | HERSH FINE SANDY LOAM, 3 TO 6 PERCENT SLOPES | 3 | 2 | 2 |
| HeD | HERSH FINE SANDY LOAM, 6 TO 11 PERCENT SLOPES | 3 | 2 | 2 |
| Hf | HOBBS SILT LOAM, 0 TO 2 PERCENT SLOPES | 3 | 3 | 3 |
| HgB | HOBBS SILT LOAM, CHANNELED | 3 | 3 | 3 |
| Ho | HOLDREGE SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| HoB | HOLDREGE SILT LOAM, 1 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| HoC | HOLDREGE SILT LOAM, 3 TO 6 PERCENT SLOPES | 3 | 2 | 2 |
| HoC2 | HOLDREGE SILT LOAM, 3 TO 6 PERCENT SLOPES, ERODED | 3 | 2 | 2 |
| Hp | HOLDREGE SILT LOAM, OVERBLOWN, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| HpB | HOLDREGE SILT LOAM, OVERBLOWN, 1 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| Hr | HORD SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| InB | INAVALE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES | 1 | 3 | 1 |
| Ke | KENESAW SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| KgB | KENESAW-COLY SILT LOAMS, 1 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| Lf | LEX LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Lg | LEX LOAM, SALINE, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| LoB | LIBORY LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES | 1 | 3 | 1 |
| Ma | MASSIE SILTY CLAY LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Pg | PITS, SAND AND GRAVEL | | 2 | 2 |
| Pm | PLATTE LOAM, 0 TO 1 PERCENT SLOPES | 1 | 3 | 1 |
| Ru | RUSCO SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| Sc | SCOTT SILT LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| SmB | SIMEON SANDY LOAM, 0 TO 3 PERCENT SLOPES | 3 | 3 | 3 |
| To | TRYON LOAMY FINE SAND, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |
| UcF | ULY-COLY SILT LOAMS, 11 TO 20 PERCENT SLOPES | 3 | 1 | 1 |
| UsF | USTORTHENTS, STEEP | 3 | 1 | 1 |
| VaB | VALENTINE LOAMY FINE SAND, 0 TO 3 PERCENT SLOPES | 1 | 3 | 1 |
| VaD | VALENTINE LOAMY FINE SAND, 3 TO 9 PERCENT SLOPES | 1 | 2 | 1 |
| VaF | VALENTINE LOAMY FINE SAND, ROLLING | 1 | 2 | 1 |
| VbD | VALENTINE-ELS LOAMY FINE SANDS, 0 TO 9 PERCENT SLOPES | 1 | 2 | 1 |
| Wa | WANN FINE SANDY LOAM, 0 TO 1 PERCENT SLOPES | 3 | 3 | 3 |