

Nontechnical Descriptions

* Lowndes County, Alabama

10 - Riverview Silt Loam, 0 To 2 Percent Slopes, Occasionally Flooded

CAPABILITY UNIT IIw-12 These deep, moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands and stream terraces. They have loamy surface layers and subsoils. The root zone can be penetrated by plant roots. Where tilled, plow pans often form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to most hay crops and pasture. The erosion hazard is slight. These soils are seasonally wet during winter and early spring months. This wetness results in restricted growth of some cool season plants and often delays spring tillage. This hazard can be partially overcome by subsurface and/or surface drainage systems. These soils can be used for row crops each year if an adequate drainage system is installed and maintained.

WOODLAND SUITABILITY GROUP-1w8. Soils in this group have very high site indexes. The site class for loblolly pine and sweetgum is 100. the soils are loamy and are moderately well to somewhat poorly drained. They occur primarily on flood plains with slopes ranging from 0 to 2 percent and have occasional flooding. This creates moderate equipment limitations and seedling mortality. These soils are suitable for growing either pines or hardwood. Species suitable to plant are loblolly pine, sweetgum, sycamore, water oak, and slash pine.

11 - Mantachie, Bibb, And Iuka Soils, 0 To 1 Percent Slopes, Frequently Flooded

CAPABILITY UNIT IIw-12a These deep, moderately well drained and somewhat poorly drained, soils (0 to 2 percent slopes) are on flood plains. They have loamy surface layers and subsoils. The root zone can be penetrated by plant roots.. Where tilled, plow pans tend to form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to most hay crops and pasture. The erosion hazard is slight. These soils are seasonally wet during winter and early spring months, and some are subject to occasional flooding. This wetness and flooding results in restricted growth of some cool season plants and often delays spring tillage. This hazard can be partially overcome by surface drainage systems and flood control measures. These soils can be used for row crops each year if an adequate subsurface and/or surface drainage system is installed and maintained.

CAPABILITY UNIT IIw-13 These deep, moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils that are low in clay. The root zone can be penetrated by plant roots. When tilled, plow pans often form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to most hay crops and pasture. The erosion hazard is slight. These soils are seasonally wet during winter and early spring months. This wetness results in restricted growth of some cool season plants and often delays spring tillage. This hazard can be partially overcome by subsurface and/or surface drainage systems. These soils can be used for row crops each year if an adequate drainage system is installed and maintained.

CAPABILITY UNIT Vw-13. These deep, poorly drained, nearly level soils (0 to 2 percent slopes) are on flood plains. They have loamy and sandy surface layers and subsoils that are loamy and sandy. The root zone is often restricted by a seasonally high water table. these soils are not suited to row crops, small grains and most hay crops and pasture because of wetness and the hazard of flooding. they are moderately well suited to water tolerant pasture plants. these soils are subject to frequent flooding in the winter and spring. This hazard can be overcome only by major flood control measures. Soil wetness can be partially overcome by extensive subsurface and/or surface drainage systems. The erosion hazard is slight.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-2w9. Soils in this group are poorly drained to very poorly drained and are either loamy throughout or have a loamy surface layer with a clayey subsoil. They are usually found on stream terraces, floodplains, and low-lying areas with slopes ranging from 0 to 2 percent. Soils in this group may have frequent flooding. This may cause severe equipment limitations and seedling mortality. The site class for these soils is 90 for loblolly pine and water oak. These soils are suitable for either pines or hardwoods. Species suitable to plant are loblolly pine, slash pine, sweetgum, and water oak.

WOODLAND SUITABILITY GROUP-1w8. Soils in this group have very high site indexes. The site class for loblolly pine and sweetgum is 100. The soils are loamy and are moderately well to somewhat poorly drained. They occur primarily on flood plains with slopes ranging from 0 to 2 percent and have occasional flooding. This creates moderate equipment limitations and seedling mortality. These soils are suitable for growing either pines or hardwood. Species suitable to plant are loblolly pine, sweetgum, sycamore, water oak, and slash pine.

WOODLAND SUITABILITY GROUP-1w9. Soils occurring in this group have a very high site index. The site class for loblolly pine is 100. The soils are loamy and are typically poorly drained. They occur primarily on flood plains with slopes of 0 to 2 percent and have frequent flooding. This causes severe equipment limitations and seedling mortality. Species suitable to plant are sweetgum, loblolly pine, green ash, water oak, and slash pine.

12 - Houlika Silty Clay Loam, 0 To 1 Percent Slopes, Frequently Flooded

CAPABILITY UNIT IVw-28. These deep, moderately well drained and somewhat poorly drained, nearly level, alkaline soils (0 to 2 percent slopes) are on flood plains. They have clayey surface layers and clayey subsoils that are sticky and plastic. The root zone is often restricted by a seasonally high water table. Spring planting of some crops is usually delayed because of flooding and wetness. These soils are moderately well suited to soybean and most hay crops and pasture. These soils are subject to frequent flooding in the winter and early spring. This hazard can be overcome only by major flood control measures. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-1w6. Soils occurring in this group have a very high site index. The site class for sweetgum is 100. The soils are loamy and are typically poorly drained. They occur primarily on flood plains with 0 to 3 percent slopes and have frequent flooding. This causes severe equipment limitations and seedling mortality. These soils are best suited for growing hardwood trees. Species suitable to plant are sweetgum, water oak, and green ash.

15 - Una Clay, Ponded

CAPABILITY UNIT IVw-28. These deep, moderately well drained and somewhat poorly drained, nearly level, alkaline soils (0 to 2 percent slopes) are on flood plains. They have clayey surface layers and clayey subsoils that are sticky and plastic. The root zone is often restricted by a seasonally high water table. Spring planting of some crops is usually delayed because of flooding and wetness. These soils are moderately well suited to soybean and most hay crops and pasture. These soils are subject to frequent flooding in the winter and early spring. This hazard can be overcome only by major flood control measures. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-1w9. Soils occurring in this group have a very high site index. The site class for loblolly pine is 100. The soils are loamy and are typically poorly drained. They occur primarily on flood plains with slopes of 0 to 2 percent and have frequent flooding. This causes severe equipment limitations and seedling mortality. Species suitable to plant are sweetgum, loblolly pine, green ash, water oak, and slash pine.

16 - Pits, Sand And Gravel

CAPABILITY UNIT VIIIs-571. These miscellaneous areas are on uplands and terraces throughout Alabama. Slopes range from 0 to more than 45 percent. Included are borrow pits, gravel pits, sand pits, and areas mined for different types of rock and ore. They are open excavations from which soil and part of the underlying material have been moved for use at another location. Some areas are reopened each year and the adjacent soil material removed. These areas are not suited to row crops, small grains, hay crops and pasture.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-4s3. This group consists of remnants from open excavations from which soil and part of the underlying material have been moved for use at another location. Included are borrow pits, gravel pits, sand pits, and areas mined for different types of rock and ore. Slopes range from 0 to more than 45 percent. Some areas are reopened each year and the adjacent soil material removed. These areas have severe ratings for erosion hazards, equipment limitations, and seedling mortality due to the different types of material. The site index ranges from 60 to 80 for loblolly pine.

17B - Troup Loamy Fine Sand, 0 To 5 Percent Slopes

CAPABILITY UNIT IIIIs-14c. These deep, well and moderately well drained, nearly level to gently sloping soils are on uplands. They have sandy surface layers about 40 to 80 inches thick over a loamy subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are well suited to bahiagrass and bermudagrass. They have low available water capacity and crops suffer from drought in most years. Young plants may be damaged by wind blown soil particles. Plant nutrients are readily leached from the root zone and frequent, light applications of fertilizers are required for maximum yields. The erosion hazard is moderate and the soils are subject to gully erosion in areas where water flow is concentrated. Conservation practices are needed to help control erosion and reduce runoff. Cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-3s2. Soils in this group are well drained to somewhat excessively drained. They are found on upland slopes ranging from 0 to 25 percent. The site class for soils in this group is 80 for loblolly pine and slash pine and 70 for longleaf pine. These soils generally have a sandy surface layer with a sandy to loamy subsoil. The sandy nature of these soils causes moderate management problems. These soils are best suited for growing pines. Species suitable for planting are slash pine, loblolly pine and longleaf pine.

17C - Troup Loamy Fine Sand, 5 To 8 Percent Slopes

CAPABILITY UNIT IVs-14d. These deep, well and moderately well drained, sloping soils are on uplands. They have sandy surface layers about 40 to 80 inches thick over a loamy subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are poorly suited to row crops and small grains. They are moderately well suited to bahiagrass and bermudagrass. They have low available water capacity and crops suffer from drought in most years. Young plants may be damaged by wind blown soil particles. Plant nutrients are readily leached from the root zone and frequent, light applications of fertilizer are required for maximum yields. The erosion hazard is severe and the soils are subject to gully erosion in areas where water is concentrated. Conservation practices are needed to help control erosion and reduce runoff. A cropping system that includes sod and close-growing crops should be used if cultivated crops are grown.

WOODLAND SUITABILITY GROUP-3s2. Soils in this group are well drained to somewhat excessively drained. They are found on upland slopes ranging from 0 to 25 percent. The site class for soils in this group is 80 for loblolly pine and slash pine and 70 for longleaf pine. These soils generally have a sandy surface layer with a sandy to loamy subsoil. The sandy nature of these soils causes moderate management problems. These soils are best suited for growing pines. Species suitable for planting are slash pine, loblolly pine and longleaf pine.

18E - Luverne-Saffell Complex, 8 To 20 Percent Slopes

CAPABILITY UNIT VIIe-13. These deep, well drained, steep soils (more than 17 percent slopes) are on uplands. They have loamy surface layers and subsoils. The root zone can be easily penetrated by plant roots. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

CAPABILITY UNIT VIIe-15. These moderately deep, well drained, steep soils (more than 12 percent or more than 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils. The root zone can be penetrated by plant roots. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

WOODLAND SUITABILITY GROUP-4f3. Soils in this group are somewhat excessively drained with a cherty silt loam surface layer and a cherty silt loam to a cherty silty clay loam subsoil. These soils occur on slopes ranging from 35 to 60 percent. The site class for loblolly pine is 70. These soils have severe management problems for erosion hazards and equipment limitations while seedling mortality is moderate. These soils are best suited for growing pines, especially loblolly pines.

21A - Bassfield Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT IIs-14a. These deep, well drained, nearly level to gently sloping soils are on uplands. They have sandy surface layers about 20 to 40 inches thick over a loamy or clayey subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are well suited to row crops such as cotton, peanuts, and sorghum; and small grains, hay crops, and pasture. They are moderately well suited to corn and soybeans. They have low available water capacity and crops suffer from drought in most years. The erosion hazard is slight to moderate. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-2o7. Soils in this group are well drained with a loamy surface layer and a loamy or clayey subsoil. These soils occur primarily on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine and 100 for yellow-poplar. These soils are suitable for growing either pines or hardwoods. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, sweetgum, yellow-poplar, black walnut, and sycamore.

24A - Heidel Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT I-13 These deep well drained and moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands. They have loamy and sandy surface layers and loamy subsoils that are low in clay. The root zone can be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight, but these soils tend to be somewhat droughty during periods of low rainfall and crop damage may be caused by wind blown soil particles during the spring. These soils can be used for cultivated crops each year with a minimum amount of conservation practices.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

24B - Heidel Sandy Loam, 2 To 5 Percent Slopes

CAPABILITY UNIT IIe-13 These deep, well drained, gently sloping soils (2 to 5 and 2 to 6 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils that are low in clay. The root zone can be easily penetrated by plant roots. When tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. They tend to be somewhat droughty during periods of low rainfall and crop damage may be caused by wind blown soil particles during the spring. The erosion hazard is slight to moderate. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

28 - Bigbee Loamy Sand, 0 To 5 Percent Slopes, Occasionally Flooded

CAPABILITY UNIT IIIIs-14. These deep, excessively drained, nearly level to gently sloping soils are on flood plains, stream terraces, and uplands. They are sandy throughout. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are moderately well suited to small grains and to crops such as cotton, peanuts, and sorghum. They are well suited to bahiagrass and bermudagrass. They have low available water capacity and crops suffer from drought in most years. Young plants may be damaged by wind blown soil particles. Plant nutrients are readily leached from the root zone and frequent, light applications of fertilizers are required for maximum yields. The erosion hazard is moderate. Conservation practices are needed to help control erosion and reduce runoff. Cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-2s2. Soils in this group are deep, excessively drained sandy soils which occur on low terraces along stream flood plains. They occur on slopes ranging from 0 to 5 percent. The site class for these soils is 90 for loblolly and slash pine. Management problems are moderate for equipment limitations and seedling mortality. These soils are best suited for pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

29 - Urbo Silty Clay Loam, 0 To 1 Percent Slopes, Frequently Flooded

CAPABILITY UNIT Vw-11. These deep, poorly drained, nearly level soils (0 to 2 percent slopes) are on low stream terraces and uplands. They have loamy surface layers and clayey subsoils. they root zone is deep, but root growth is often restricted by a seasonally high water table and the clayey subsoil. These soils are not suited to row crops, small grains, and most hay crops and pasture because of wetness and the hazard of flooding. They are moderately well suited to water tolerant pasture plants. Soil wetness can be partially overcome by extensive surface drainage systems. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-1w9. Soils occurring in this group have a very high site index. The site class for loblolly pine is 100. The soils are loamy and are typically poorly drained. They occur primarily on flood plains with slopes of 0 to 2 percent and have frequent flooding. This causes severe equipment limitations and seedling mortality. Species suitable to plant are sweetgum, loblolly pine, green ash, water oak, and slash pine.

30A - Lucedale Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT I-12 These deep, well drained and moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands and high stream terraces. They have a loamy or sandy surface layer and a loamy subsoil. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight, but some crop damage may be caused by wind blown soil particles during the spring. These soils can be used for cultivated crops each year with a minimum of conservation practices.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

30B - Lucedale Fine Sandy Loam, 2 To 5 Percent Slopes

CAPABILITY UNIT IIe-12 These deep, well drained and moderately well drained, gently sloping soils (2 to 5 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight to moderate, and some crop damage may be caused by wind blown soil particles during the spring. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

30C - Smithdale Fine Sandy Loam, 5 To 8 Percent Slopes

CAPABILITY UNIT IIIe-12. These deep, well drained and moderately well drained sloping soils (5 to 8 percent and 6 to 10 percent slopes) are on stream terraces and uplands. They have loamy and sandy surface layers and sandy subsoils. The root zone can be easily penetrated by plant roots. When tilled, plow pans form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to hay crops and pasture. The erosion hazard is moderate to severe and some crop damage may be caused by wind blown soil particles during the spring. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

30D - Smithdale Fine Sandy Loam, 8 To 12 Percent Slopes

CAPABILITY UNIT IVe-12. These deep, well drained and moderately well drained, strongly sloping soils (8 to 12 percent slopes) are on uplands. They have loamy surface layers and subsoils. The root zone can be easily penetrated by plant roots. When tilled, plow pans form and restrict root growth of some annual crops. These soils are poorly suited to row crops and moderately well to poorly suited to small grains. They are moderately well suited to hay crops and pasture. The erosion hazard is severe. A good system of conservation practices is essential when these soils are used for cultivated row crops. No-till or cropping systems that include sod and close growing crops must be used in combination if cultivated crops are grown.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

31A - Bama Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT I-12 These deep, well drained and moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands and high stream terraces. They have a loamy or sandy surface layer and a loamy subsoil. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight, but some crop damage may be caused by wind blown soil particles during the spring. These soils can be used for cultivated crops each year with a minimum of conservation practices.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

31B - Bama Fine Sandy Loam, 2 To 5 Percent Slopes

CAPABILITY UNIT IIe-12 These deep, well drained and moderately well drained, gently sloping soils (2 to 5 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight to moderate, and some crop damage may be caused by wind blown soil particles during the spring. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

34A - Minter Loam, 0 To 1 Percent Slopes, Occasionally Flooded

CAPABILITY UNIT IVw-11. These deep, poorly drained, nearly level soils (0 to 2 percent slopes) are on uplands, stream terraces, and flood plains. They have loamy surface layers and clayey subsoils. The root zone is often restricted by a seasonally high water table. These soils are poorly suited to row crops, small grains and most hay crops and pasture because of wetness. They are moderately well suited to Argentine bahiagrass in the southern part of the state. Some of these soils are subject to flooding in the winter and early spring. Soil wetness can be partially overcome by extensive surface drainage systems. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-2w9. Soils in this group are poorly drained to very poorly drained and are either loamy throughout or have a loamy surface layer with a clayey subsoil. They are usually found on stream terraces, floodplains, and low-lying areas with slopes ranging from 0 to 2 percent. Soils in this group may have frequent flooding. This may cause severe equipment limitations and seedling mortality. The site class for these soils is 90 for loblolly pine and water oak. These soils are suitable for either pines or hardwoods. Species suitable to plant are loblolly pine, slash pine, sweetgum, and water oak.

35A - Cahaba Fine Sandy Loam, 0 To 2 Percent Slopes, Rarely Flooded

CAPABILITY UNIT I-36. These deep, well drained, nearly level soils (0 to 2 percent slopes) are on uplands and stream terraces. They have loamy surface layers and loamy subsoils. The root zone is easily penetrated by plant roots. These soils are well suited to row crops, small grains, hay crops and pasture. The erosion hazard is slight. These soils can be used for cultivated crops each year with a minimum of conservation practices.

WOODLAND SUITABILITY GROUP-2o7. Soils in this group are well drained with a loamy surface layer and a loamy or clayey subsoil. These soils occur primarily on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine and 100 for yellow-poplar. These soils are suitable for growing either pines or hardwoods. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, sweetgum, yellow-poplar, black walnut, and sycamore.

35B - Cahaba Fine Sandy Loam, 2 To 5 Percent Slopes, Rarely Flooded

CAPABILITY UNIT IIe-36. These deep and moderately deep, well drained, gently sloping soils (2 to 6 percent slopes) are on uplands and stream terraces. They have loamy surface layers and loamy subsoils. The root zone can be easily penetrated by plant roots. These soils are well suited to row crops, small grains, hay crops and pasture. The erosion hazard is slight to moderate. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated crops each year if a good system of conservation practices is established and maintained.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-2o7. Soils in this group are well drained with a loamy surface layer and a loamy or clayey subsoil. These soils occur primarily on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine and 100 for yellow-poplar. These soils are suitable for growing either pines or hardwoods. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, sweetgum, yellow-poplar, black walnut, and sycamore.

36A - Mcqueen Fine Sandy Loam, 0 To 2 Percent Slopes, rarely Flooded

CAPABILITY UNIT I-15 These deep, well drained, nearly level soils (0 to 2 percent slopes) are on stream terraces and uplands. They have loamy and sandy surface layers and clayey subsoils. The root zone can usually be penetrated by plant roots. Where tilled, plow pans tend to form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. Canton Bend and McQueen soils are subject to occasional brief flooding mostly during the winter. The erosion hazard is slight. These soils can be used for cultivated crops each year with a minimum amount of conservation practices.

WOODLAND SUITABILITY GROUP-3o7. Soils in this group are usually well drained with a loamy surface layer and either a loamy or clayey subsoil. These soils occur primarily on uplands with slopes ranging from 0 to 15 percent. The site class is 80 for loblolly pine and 90 for yellow poplar. These soils are suitable for growing either pines or hardwoods. There are no significant management problems associate with these soils. Species suitable to plant are loblolly pine, slash pine, sweetgum, yellow-poplar, water oak, and sycamore.

36B - Mcqueen Fine Sandy Loam, 2 To 5 Percent Slopes, Rarely Flooded

CAPABILITY UNIT IIe-13 These deep, well drained, gently sloping soils (2 to 5 and 2 to 6 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils that are low in clay. The root zone can be easily penetrated by plant roots. When tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. They tend to be somewhat droughty during periods of low rainfall and crop damage may be caused by wind blown soil particles during the spring. The erosion hazard is slight to moderate. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-3o7. Soils in this group are usually well drained with a loamy surface layer and either a loamy or clayey subsoil. These soils occur primarily on uplands with slopes ranging from 0 to 15 percent. The site class is 80 for loblolly pine and 90 for yellow poplar. These soils are suitable for growing either pines or hardwoods. There are no significant management problems associate with these soils. Species suitable to plant are loblolly pine, slash pine, sweetgum, yellow-poplar, water oak, and sycamore.

40B - Luverne Fine Sandy Loam, 2 To 5 Percent Slopes

CAPABILITY UNIT IIIe-15. These deep, moderately well drained and well drained, gently sloping soils (1 to 4 or 2 to 5 percent slopes) are on stream terraces and uplands. They have loamy surface layers and clayey subsoils. The root zone can usually be penetrated by plant roots. When tilled, plow pans tend to form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to hay crops and pasture. The erosion hazard is moderate to severe. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

40D - Luverne Fine Sandy Loam, 5 To 15 Percent Slopes

CAPABILITY UNIT VIe15. These moderately deep to deep, well drained, strongly sloping soils (8 to 12 and 10 to 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils. The root zone is often restricted by the clayey subsoil. These soils are not suited to row crops, hay crops and small grains. They are moderately well to poorly suited to pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

40E - Luverne Fine Sandy Loam, 15 To 35 Percent Slopes

CAPABILITY UNIT VIIe-15. These moderately deep, well drained, steep soils (more than 12 percent or more than 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils. The root zone can be penetrated by plant roots. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

41A - Pheba Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT IIIw-19. These deep, somewhat poorly drained, nearly level soils (0 to 2 percent slopes) with fragipans are on uplands and stream terraces. They have loamy surface layers and subsoils. When tilled, plow pans may form and restrict root growth of some annual crops. A compact, brittle layer in the subsoil restricts root growth of most annual plants and perches water during winter, early spring, and after other periods of high rainfall. These soils are moderately well suited to crops such as soybeans, sorghum and potatoes and poorly suited to hay crops and pasture. Wetness results in restricted growth of many plants and delays spring tillage. This can be partially overcome by extensive subsurface and/or surface drainage systems. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-2w9. Soils in this group are poorly drained to very poorly drained and are either loamy throughout or have a loamy surface layer with a clayey subsoil. They are usually found on stream terraces, floodplains, and low-lying areas with slopes ranging from 0 to 2 percent. Soils in this group may have frequent flooding. This may cause severe equipment limitations and seedling mortality. The site class for these soils is 90 for loblolly pine and water oak. These soils are suitable for either pines or hardwoods. Species suitable to plant are loblolly pine, slash pine, sweetgum, and water oak.

44A - Benndale Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT I-13 These deep well drained and moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands. They have loamy and sandy surface layers and loamy subsoils that are low in clay. The root zone can be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight, but these soils tend to be somewhat droughty during periods of low rainfall and crop damage may be caused by wind blown soil particles during the spring. These soils can be used for cultivated crops each year with a minimum amount of conservation practices.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

49 - Marietta Fine Sandy Loam, 0 To 1 Percent Slopes, Frequently Flooded

CAPABILITY UNIT IVw-22. These deep, moderately well drained, nearly level, alkaline soils (5 to 8 percent slopes) are on flood plains. They have loamy surface layers and subsoils. The root zone can be penetrated by plant roots. When tilled, plow pans tend to form and restrict root growth of some annual crops. These soils are moderately well suited to soybean and moderately well suited to most hay crops and pasture. The erosion hazard is slight. These soils are seasonally wet during winter and early spring months because of a high water table and frequent flooding. This wetness and flooding results in restricted growth of some cool season plants and often delays spring tillage. This hazard can be partially overcome by surface and/or subsurface drainage systems and flood control measures. These soils can be used for row crops each year if selection in row crops is practiced and an adequate drainage system and flood control measures are installed and maintained.

WOODLAND SUITABILITY GROUP-1w6. Soils occurring in this group have a very high site index. The site class for sweetgum is 100. The soils are loamy and are typically poorly drained. They occur primarily on flood plains with 0 to 3 percent slopes and have frequent flooding. This causes severe equipment limitations and seedling mortality. These soils are best suited for growing hardwood trees. Species suitable to plant are sweetgum, water oak, and green ash.

50C - Conecuh Fine Sandy Loam, 1 To 5 Percent Slopes

CAPABILITY UNIT IIIe-18. These deep, moderately well drained to poorly drained, gently sloping soils (2 to 5 percent or 2 to 6 percent slopes) are on uplands. They have loamy or sandy surface layers and sticky, plastic, clayey subsoils. The root zone is often restricted because of the clayey subsoil. These soils are difficult to till because of the high clay content in the upper subsoil. These soils are moderately well suited to row crops and small grains and well suited to hay crops and pasture. The erosion hazard is moderate to severe. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

50E - Conecuh Fine Sandy Loam, 5 To 15 Percent Slopes

CAPABILITY UNIT VIe-18. These deep and moderately deep, moderately well drained, strongly sloping soils (8 to 12 and 10 to 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils that are sticky and plastic. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are not suited to row crops, hay crops, and small grains. They are moderately well to poorly suited to pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

52C - Oktibbeha Clay, 1 To 5 Percent Slopes

CAPABILITY UNIT IIIe-28. These deep, well drained, gently sloping, alkaline soils (3 to 5 percent slopes) are on uplands. They have clayey surface layers and sticky and plastic clayey subsoils. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are moderately well suited to soybeans and okra and well suited to hay crops and pasture. The erosion hazard is moderate. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-3c8. Soils in this group are moderately well drained and clayey throughout. They occur on uplands with slopes ranging from 2 to 17 percent. The site class is 80 for loblolly pine, sweetgum, and water oak. Because of the clayey nature of these soils, moderate management problems occur for equipment limitations and seedling mortality. These soils are suitable for either pines or hardwoods. Species suitable for planting include loblolly pine, Eastern redcedar, sweetgum and water oak.

52E - Oktibbeha Clay, 5 To 15 Percent Slopes, Eroded

CAPABILITY UNIT VIe-28a. These deep, moderately well and somewhat poorly drained, strongly sloping to moderately steep, acid soils (8 to 17 percent slopes) are on uplands. They have clayey surface layers and clayey subsoils that are sticky and plastic. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are not suited to row crops, hay crops, and small grains. They are moderately well to poorly suited for pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-3c8. Soils in this group are moderately well drained and clayey throughout. They occur on uplands with slopes ranging from 2 to 17 percent. The site class is 80 for loblolly pine, sweetgum, and water oak. Because of the clayey nature of these soils, moderate management problems occur for equipment limitations and seedling mortality. These soils are suitable for either pines or hardwoods. Species suitable for planting include loblolly pine, Eastern redcedar, sweetgum and water oak.

52F - Oktibbeha-Brantley Complex, 15 To 35 Percent Slopes, Eroded

CAPABILITY UNIT VIIe-15. These moderately deep, well drained, steep soils (more than 12 percent or more than 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils. The root zone can be penetrated by plant roots. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

CAPABILITY UNIT VIIe-22. These moderately deep, well drained, moderately steep to steep, alkaline soils (more than 8 percent) are on uplands. They have a clay surface layer and a clayey subsoil. In many areas the present surface layer is a mixture of the original loamy surface layer and the upper part of the clayey subsoil. The root zone can be penetrated by plant roots. The erosion hazard is very severe and these soils are not suited to row crops, hay crops, or small grains. They are poorly suited for pasture.

WOODLAND SUITABILITY GROUP-3o1. Soils in this group are well drained to excessively drained and are primarily loamy and occur on uplands with slopes ranging from 0 to 15 percent. The site class is 90 for loblolly pine. These soils have no significant management problems. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

WOODLAND SUITABILITY GROUP-3c8. Soils in this group are moderately well drained and clayey throughout. They occur on uplands with slopes ranging from 2 to 17 percent. The site class is 80 for loblolly pine, sweetgum, and water oak. Because of the clayey nature of these soils, moderate management problems occur for equipment limitations and seedling mortality. These soils are suitable for either pines or hardwoods. Species suitable for planting include loblolly pine, Eastern redcedar, sweetgum and water oak.

53C - Sumter Silty Clay, 1 To 5 Percent Slopes, Eroded

CAPABILITY UNIT IIIe-22. These moderately deep, well drained, gently sloping alkaline soils (3 to 5 percent slopes) are on uplands. They have clayey surface layers and subsoils. The root zone can usually be penetrated by plant roots. These soils are moderately well suited to soybeans, hay crops and pasture. The erosion hazard is moderate. Conservation practices are needed to help control erosion and reduce runoff. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

53F - Sumter Silty Clay, 5 To 17 Percent Slopes, Eroded

CAPABILITY UNIT VIIe-22. These moderately deep, well drained, moderately steep to steep, alkaline soils (more than 8 percent) are on uplands. They have a clay surface layer and a clayey subsoil. In many areas the present surface layer is a mixture of the original loamy surface layer and the upper part of the clayey subsoil. The root zone can be penetrated by plant roots. The erosion hazard is very severe and these soils are not suited to row crops, hay crops, or small grains. They are poorly suited for pasture.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

54B - Okolona Clay, 1 To 3 Percent Slopes

CAPABILITY UNIT IIe-28 These deep, well drained, gently sloping alkaline soils (1 to 3 percent slopes) are on uplands. They have clayey surface layers and sticky and plastic, clayey subsoils. The root zone can usually be penetrated by plant roots. These soils are well suited to soybeans, okra, hay crops, and pasture. They are moderately well suited to small grains. The erosion hazard is moderate. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

54C - Okolona Clay, 3 To 5 Percent Slopes

CAPABILITY UNIT IIIe-28. These deep, well drained, gently sloping, alkaline soils (3 to 5 percent slopes) are on uplands. They have clayey surface layers and sticky and plastic clayey subsoils. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are moderately well suited to soybeans and okra and well suited to hay crops and pasture. The erosion hazard is moderate. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

56A - Catalpa Clay, 0 To 1 Percent Slopes, Occasionally Flooded

CAPABILITY UNIT IIw-28. These deep, moderately well drained and somewhat poorly drained, nearly level, alkaline soils (0 to 2 percent slopes) are on flood plains. They have clayey surface layers and clayey subsoils that are sticky and plastic. The root zone is often restricted by a seasonally high water table. Spring planting of some crops is usually delayed because of flooding and wetness. These soils are moderately well suited to soybeans and most hay crops and pasture. These soils are subject to occasional flooding in the winter and early spring. This hazard can be overcome only by major flood control measures. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-1w5. Soils occurring in this group have a very high site index. The site class for sweetgum is 100. The soils are loamy and are moderately well to somewhat poorly drained and occur primarily on flood plains with slopes ranging from 0 to 3 percent with occasional flooding. This causes moderate equipment limitations and seedling mortality. These soils are best suited for growing hardwoods. Species suitable to plant are cottonwood, sweetgum, sycamore, yellow-poplar, green ash, and water oak.

56B - Faunsdale Clay Loam, 1 To 3 Percent Slopes

CAPABILITY UNIT IIe-28 These deep, well drained, gently sloping alkaline soils (1 to 3 percent slopes) are on uplands. They have clayey surface layers and sticky and plastic, clayey subsoils. The root zone can usually be penetrated by plant roots. These soils are well suited to soybeans, okra, hay crops, and pasture. They are moderately well suited to small grains. The erosion hazard is moderate. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

57C - Sumter-Oktibbeha Complex, 1 To 5 Percent Slopes

CAPABILITY UNIT IIIe-22. These moderately deep, well drained, gently sloping alkaline soils (3 to 5 percent slopes) are on uplands. They have clayey surface layers and subsoils. The root zone can usually be penetrated by plant roots. These soils are moderately well suited to soybeans, hay crops and pasture. The erosion hazard is moderate. Conservation practices are needed to help control erosion and reduce runoff. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

57F - Sumter-Oktibbeha Complex, 5 To 17 Percent Slopes, Eroded

CAPABILITY UNIT VIe-22. These moderately deep, well drained, strongly sloping to moderately steep, alkaline soils (more than 8 percent slopes) are on uplands. They have clayey surface layers and clayey subsoils. The root zone can be penetrated by plant roots. These soils are not suited to row crops, hay crops, or small grains. They are moderately well to poorly suited for pasture. The erosion hazard is very severe.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

CAPABILITY UNIT VIe-28a. These deep, moderately well and somewhat poorly drained, strongly sloping to moderately steep, acid soils (8 to 17 percent slopes) are on uplands. They have clayey surface layers and clayey subsoils that are sticky and plastic. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are not suited to row crops, hay crops, and small grains. They are moderately well to poorly suited for pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.

58A - Macon Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT I-12 These deep, well drained and moderately well drained, nearly level soils (0 to 2 percent slopes) are on uplands and high stream terraces. They have a loamy or sandy surface layer and a loamy subsoil. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight, but some crop damage may be caused by wind blown soil particles during the spring. These soils can be used for cultivated crops each year with a minimum of conservation practices.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

58B - Macon Fine Sandy Loam, 2 To 5 Percent Slopes

CAPABILITY UNIT IIe-12 These deep, well drained and moderately well drained, gently sloping soils (2 to 5 percent slopes) are on uplands and stream terraces. They have loamy and sandy surface layers and loamy subsoils. The root zone can easily be penetrated by plant roots. Where tilled, plow pans form and restrict root growth of some annual crops. These soils are well suited to row crops, small grains, hay crops, and pasture. The erosion hazard is slight to moderate, and some crop damage may be caused by wind blown soil particles during the spring. Conservation practices are needed to help control erosion and reduce runoff. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

58C - Macon Fine Sandy Loam, 5 To 8 Percent Slopes

CAPABILITY UNIT IIIe-12. These deep, well drained and moderately well drained sloping soils (5 to 8 percent and 6 to 10 percent slopes) are on stream terraces and uplands. They have loamy and sandy surface layers and sandy subsoils. The root zone can be easily penetrated by plant roots. When tilled, plow pans form and restrict root growth of some annual crops. These soils are moderately well suited to row crops and small grains and well suited to hay crops and pasture. The erosion hazard is moderate to severe and some crop damage may be caused by wind blown soil particles during the spring. A combination of several conservation practices is needed on cultivated fields to control erosion and provide for proper water disposal. No-till or cropping systems that include sod and close growing crops are usually needed in rotation with cultivated crops.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

WOODLAND SUITABILITY GROUP-2o1. Soils in this group are well drained and are primarily loamy. These soils occur on uplands with slopes ranging from 0 to 15 percent. The site class is high and is 90 for loblolly pine. These soils are best suited for growing pines. There are no significant management problems associated with these soils. Species suitable to plant are loblolly pine, slash pine, and longleaf pine.

60F - Luverne-Conecuh Complex, 8 To 25 Percent Slopes

CAPABILITY UNIT VIIe-15. These moderately deep, well drained, steep soils (more than 12 percent or more than 15 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils. The root zone can be penetrated by plant roots. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

CAPABILITY UNIT VIIe-18. These deep and moderately deep, moderately well drained and somewhat poorly drained, moderately steep and steep soils (more than 12 percent slopes) are on uplands. They have loamy surface layers and clayey subsoils that are sticky and plastic. The root zone is deep, but plant roots may be restricted by the clayey subsoil. These soils are not suited to row crops, small grains, hay crops or pasture. The erosion hazard is very severe.

WOODLAND SUITABILITY GROUP-3c2. Soils in this group are well drained to poorly drained and occur on slopes ranging from 1 to 35 percent. These soils have a site class of 80 for loblolly pine. These soils have moderate erosion hazards, equipment limitations, and seedling mortality due to the steep slopes and clay content. They are best suited for growing pines. Species suitable to plant are slash pine, loblolly pine, and longleaf pine.

70B - Lucy Loamy Fine Sand, 0 To 5 Percent Slopes

CAPABILITY UNIT IIIs-14a. These deep, well drained, nearly level to gently sloping soils are on uplands. They have sandy surface layers about 20 to 40 inches thick over a loamy or clayey subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are well suited to row crops such as cotton, peanuts, and sorghum; and small grains, hay crops, and pasture. They are moderately well suited to corn and soybeans. They have low available water capacity and crops suffer from drought in most years. The erosion hazard is slight to moderate. These soils can be used for cultivated row crops each year if a good system of conservation practices is established and maintained.

WOODLAND SUITABILITY GROUP-3s2. Soils in this group are well drained to somewhat excessively drained. They are found on upland slopes ranging from 0 to 25 percent. The site class for soils in this group is 80 for loblolly pine and slash pine and 70 for longleaf pine. These soils generally have a sandy surface layer with a sandy to loamy subsoil. The sandy nature of these soils causes moderate management problems. These soils are best suited for growing pines. Species suitable for planting are slash pine, loblolly pine and longleaf pine.

70C - Lucy Loamy Fine Sand, 5 To 8 Percent Slopes

CAPABILITY UNIT IIIIs-14b. These deep, well drained, sloping soils are on uplands. They have sandy surface layers about 20 to 40 inches thick over a loamy or clayey subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are moderately well suited to small grains and to crops such as cotton, peanuts, and sorghum. They are well suited to bahiagrass and bermudagrass. They have low available water capacity and crops suffer from drought in most years. Young plants may be damaged by wind blown soil particles. The erosion hazard is moderate to severe and the soils are subject to gully erosion in areas where water is concentrated. Conservation practices are needed to help control erosion and reduce runoff. Cropping systems that include sod and close-growing crops are usually needed in rotation with cultivated crops.

WOODLAND SUITABILITY GROUP-3s2. Soils in this group are well drained to somewhat excessively drained. They are found on upland slopes ranging from 0 to 25 percent. The site class for soils in this group is 80 for loblolly pine and slash pine and 70 for longleaf pine. These soils generally have a sandy surface layer with a sandy to loamy subsoil. The sandy nature of these soils causes moderate management problems. These soils are best suited for growing pines. Species suitable for planting are slash pine, loblolly pine and longleaf pine.

Nontechnical Descriptions (cont.)

* Lowndes County, Alabama

70D - Lucy Loamy Fine Sand, 8 To 12 Percent Slopes

CAPABILITY UNIT IVs-14b. These deep, well drained, strongly sloping soils are on uplands. They have sandy surface layers about 20 to 40 inches thick over a loamy or clayey subsoil. The root zone is easily penetrated by plant roots. Plow pans form easily and may restrict root growth of some crops. These soils are poorly suited to crops and small grains. They are moderately well suited to bahiagrass and bermudagrass. They have low available water capacity and crops suffer from drought in most years. Young plants may be damaged by wind blown soil particles. The erosion hazard is severe and the soils are subject to gully erosion in areas where water is concentrated. Conservation practices are needed to help control erosion and reduce runoff. A cropping system that includes sod and close growing crops should be used if cultivated crops are grown.

WOODLAND SUITABILITY GROUP-3s2. Soils in this group are well drained to somewhat excessively drained. They are found on upland slopes ranging from 0 to 25 percent. The site class for soils in this group is 80 for loblolly pine and slash pine and 70 for longleaf pine. These soils generally have a sandy surface layer with a sandy to loamy subsoil. The sandy nature of these soils causes moderate management problems. These soils are best suited for growing pines. Species suitable for planting are slash pine, loblolly pine and longleaf pine.

83A - Rains Fine Sandy Loam, 0 To 2 Percent Slopes

CAPABILITY UNIT IVw-12. These deep, poorly drained, nearly level soils (0 to 2 percent slopes) are on uplands and stream terraces. They have loamy surface layers and subsoils. When tilled, plow pans may form and restrict root growth of some annual crops. Also, the root zone is often restricted by a seasonally high water table. These soils are poorly suited to row crops, small grains and most hay crops and pasture because of wetness. They are moderately well suited to Argentine bahiagrass in the southern part of the state. Soil wetness can be partially overcome by extensive subsurface and/or surface drainage systems. The erosion hazard is slight.

WOODLAND SUITABILITY GROUP-2w9. Soils in this group are poorly drained to very poorly drained and are either loamy throughout or have a loamy surface layer with a clayey subsoil. They are usually found on stream terraces, floodplains, and low-lying areas with slopes ranging from 0 to 2 percent. Soils in this group may have frequent flooding. This may cause severe equipment limitations and seedling mortality. The site class for these soils is 90 for loblolly pine and water oak. These soils are suitable for either pines or hardwoods. Species suitable to plant are loblolly pine, slash pine, sweetgum, and water oak.

99 - Gullied Land, Alkaline

CAPABILITY UNIT VIIe-151. These moderately deep to deep, well drained, strongly sloping to steep soils (more than 12 percent and more than 15 percent slopes) are on uplands. Most of the original surface layers have been lost from erosion. The present surface layer is a mixture of the original loamy surface layer and the upper part of the clayey subsoil. This mixture results in poor tilth and increases runoff. The root zone can be penetrated by plant roots. The erosion hazard is very severe. The soils are not suited to row crops, small grains, hay crops or pasture.

WOODLAND SUITABILITY GROUP-4c2. Soils in this group are well drained to moderately well drained and generally have a thin sandy loam surface layer with a clayey subsoil. Severely eroded soils may be clayey throughout as well as some of the soils that occur in the blackbelt area of the state. These soils occur on slopes ranging from 1 to 35 percent. Management problems are moderate because of the clayey nature of these soils and steep slopes. The site class is 70 for loblolly pine and 40 for eastern redcedar. These soils are best suited for pines on acid soils and cedar on calcareous soils. Species suitable to plant are loblolly pine and eastern redcedar.
