

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT  
 Wicomico County, Maryland: Detailed Soil Map Legend

Map Symbol	Soil Mapunit Name	HEL Classification	
		R=	C=
Ba	BAYBORO LOAM	not highly erodible	
Bb	BAYBORO SILT LOAM	not highly erodible	
Be	BEACHES	not highly erodible	
Bo	BORROW PITS	not highly erodible	
DoA	DOWNER LOAMY SAND, 0 TO 2 PERCENT SLOPES	not highly erodible	
DoB2	DOWNER LOAMY SAND, 2 TO 5 PERCENT SLOPES, MODERATELY ERODED	potentially highly erodible	
DoC	DOWNER LOAMY SAND, 5 TO 10 PERCENT SLOPES	potentially highly erodible	
Ea	ELKTON LOAM	not highly erodible	
Ek	ELKTON SANDY LOAM	not highly erodible	
Em	ELKTON SILT LOAM	not highly erodible	
En	ELKTON SILTY CLAY LOAM	not highly erodible	
EoD	EVESBORO LOAMY SAND, 5 TO 15 PERCENT SLOPES	potentially highly erodible	
EpB	EVESBORO LOAMY SAND, CLAYEY SUBSTRATUM, 0 TO 5 PERCENT SLOPES	not highly erodible	
ErD	EVESBORO SAND, 5 TO 15 PERCENT SLOPES	potentially highly erodible	
EsB	EVESBORO SAND, CLAYEY SUBSTRATUM, 0 TO 5 PERCENT SLOPES	not highly erodible	
EtF	EVESBORO SOILS, 15 TO 40 PERCENT SLOPES	highly erodible	
EvD	EVESBORO-GALESTOWN SANDS, 5 TO 15 PERCENT SLOPES	potentially highly erodible	
EwB	EVESBORO-GALESTOWN SANDS, CLAYEY SUBSTRATUM, 0 TO 5 PERCENT SLOPES	not highly erodible	
EyC	EVESBORO-GALESTOWN-DOWNER LOAMY SANDS, 0 TO 10 PERCENT SLOPES	potentially highly erodible	
Fa	FALLSINGTON FINE SANDY LOAM	not highly erodible	
Fg	FALLSINGTON LOAM	not highly erodible	
Fs	FALLSINGTON SANDY LOAM	not highly erodible	
GaD	GALESTOWN LOAMY SAND, 5 TO 15 PERCENT SLOPES	potentially highly erodible	
GcB	GALESTOWN LOAMY SAND, CLAYEY SUBSTRATUM, 0 TO 5 PERCENT SLOPES	not highly erodible	
KeA	KEYPORT SILT LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
KeB	KEYPORT SILT LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
KsA	KLEJ LOAMY SAND, 0 TO 2 PERCENT SLOPES	not highly erodible	
KsB	KLEJ LOAMY SAND, 2 TO 5 PERCENT SLOPES	not highly erodible	
Le	LEON LOAMY SAND	not highly erodible	
Ma	MADE LAND	not highly erodible	
MdA	MATAPEAKE FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
MdB2	MATAPEAKE FINE SANDY LOAM, 2 TO 5 PERCENT SLOPES MODERATELY ERODED	potentially highly erodible	
MeA	MATAPEAKE SILT LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
MeB2	MATAPEAKE SILT LOAM, 2 TO 5 PERCENT SLOPES, MODERATELY ERODED	potentially highly erodible	
MeC	MATAPEAKE SILT LOAM, 5 TO 10 PERCENT SLOPES	potentially highly erodible	
MfA	MATAWAN FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	potentially highly erodible	
MfB	MATAWAN FINE SANDY LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
MmA	MATAWAN LOAMY SAND, 0 TO 2 PERCENT SLOPES	potentially highly erodible	
MmB	MATAWAN LOAMY SAND, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
MmC	MATAWAN LOAMY SAND, 5 TO 10 PERCENT SLOPES	highly erodible	
MmE	MATAWAN LOAMY SAND, 10 TO 30 PERCENT SLOPES	highly erodible	

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued  
 Wicomico County, Maryland: Detailed Soil Map Legend

Map Symbol	Soil Mapunit Name	HEL Classification	
		R=	C=
MnA	MATAWAN SANDY LOAM, 0 TO 2 PERCENT SLOPES	potentially highly erodible	
MnB	MATAWAN SANDY LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
MpA	MATTAPEX LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
MpB	MATTAPEX LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
MtA	MATTAPEX SILT LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
MtB	MATTAPEX SILT LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
Mu	MUCK	not highly erodible	
Mv	MIXED ALLUVIAL LAND	not highly erodible	
NoA	NORFOLK LOAMY SAND, 0 TO 2 PERCENT SLOPES	not highly erodible	
NoB	NORFOLK LOAMY SAND, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
NoC	NORFOLK LOAMY SAND, 5 TO 10 PERCENT SLOPES	potentially highly erodible	
NsD	NORFOLK AND SASSAFRAS SOILS, 10 TO 15 PERCENT SLOPES	potentially highly erodible	
NsE	NORFOLK AND SASSAFRAS SOILS, 15 TO 30 PERCENT SLOPES	highly erodible	
Ot	OTHELLO SILT LOAM	not highly erodible	
Ow	OTHELLO SILT LOAM, LOW	not highly erodible	
Pe	PLUMMER LOAMY SAND	not highly erodible	
Pk	POCOMOKE LOAM	not highly erodible	
Po	POCOMOKE SANDY LOAM	not highly erodible	
Pr	PORTSMOUTH SANDY LOAM	not highly erodible	
Pt	PORTSMOUTH SILT LOAM	not highly erodible	
Ru	RUTLEGE LOAMY SAND	not highly erodible	
SaA	SASSAFRAS FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
SaB	SASSAFRAS FINE SANDY LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
SsA	SASSAFRAS SANDY LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
SsB2	SASSAFRAS SANDY LOAM, 2 TO 5 PERCENT SLOPES, MODERATELY ERODED	potentially highly erodible	
SsC2	SASSAFRAS SANDY LOAM, 5 TO 10 PERCENT SLOPES, MODERATELY ERODED	potentially highly erodible	
St	ST. JOHNS LOAMY SAND	not highly erodible	
Su	ST. JOHNS MUCKY LOAMY SAND	not highly erodible	
Sw	SWAMP	not highly erodible	
Tm	TIDAL MARSH	not highly erodible	
WfA	WOODSTOWN FINE SANDY LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
WfB	WOODSTOWN FINE SANDY LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	
WoA	WOODSTOWN LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
WsA	WOODSTOWN SANDY LOAM, 0 TO 2 PERCENT SLOPES	not highly erodible	
WsB	WOODSTOWN SANDY LOAM, 2 TO 5 PERCENT SLOPES	potentially highly erodible	

