

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Loamy (Ly) 10-14” Precipitation Zone,

Site ID: 058BY122WY

Major Land Resource Area: 58B – Northern Rolling High Plains

Physiographic Features

This site occurs on gently undulating rolling land.

Landform: Hill sides, alluvial fans, ridges & stream terraces

Aspect: N/A

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	3800	5100
Slope (percent):	0	30
Water Table Depth (inches):	None within 60 inches	
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):	0	0
Frequency:	None	None
Duration:	None	None
Runoff Class:	negligible	high

Climatic features

Annual precipitation ranges from 10-14 inches per year. Wide fluctuations may occur in yearly precipitation and result in more drought years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Wind speed averages about 8 mph, ranging from 10 mph during the spring to 7 mph during late summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 75 mph.

Growth of native cool season plants begins about April 1 and continues to about July 1. Native warm season plants begin growth about May 15 and continue to about August 15. Green up of cool season plants may occur in September and October of most years.

The following information is from the “Clearmont 5 SW” climate station:

Site Type: Rangeland
MLRA: 58B – Northern Rolling High Plains

**Loamy 10-14" P.Z.
R058BY122WY**

Frost-free period (32 °F): 76 - 132 days; (5 yrs. out of 10, these days will occur between May 30 – September 11)

Freeze-free period 28 °F): 110 - 145 days; (5 yrs. out of 10, these days will occur between May 16 – September 21)

Mean annual precipitation: 12.4 inches

Mean annual air temperature: 43.2 °F (28.4°F Avg. Min. – 57.9°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include: "Dull Center"

Influencing Water Features

Wetland Description:	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
None	None	None	None	None

Stream Type: None

Representative Soil Features

The soils of this site are deep to moderately deep (greater than 20" to bedrock), well drained & moderately permeable. Layers of the soil most influential to the plant community varies from 3 to 6 inches thick. These layers consist of the A horizon with very fine sandy loam, loam, or silt loam texture and may also include the upper few inches of the B horizon with sandy clay loam, silty clay loam or clay loam texture.

Major Soil Series correlated to this site includes: Bidman, Cambria, Cushman, Forkwood, Kishona, Parmleed, Theedle and Zigweid.

Other Soil Series correlated to this site in MLRA 58B include: Absted, Arvada, Ascalon, Big Horn, Bowbac, Briggsdale, Cambria Variant, Cedak Dry, Clarkelen, Connerton, Docpar, El Rancho, Emigha, Emigrant, Forkwood Variant, Fort Collins, Garrett, Glendo, Harlan, Harlan Dry, Haverdad, Hiland, Jonpol, Kadoka, Keota, Keyner, Kim, Kirtley, Larim, Larimer, Lawver, Lohsman, Maysdorf, Neville, Noden, Nuncho, Platmak, Platmak Dry, Pugsley, Recluse, Recluse Dry, Redbow, Reddale, Renohill, Roughlock, Senlar, Spearman, Stoneham, Teckla, Thirtynine, Ulm, Ulm Dry, Wages, Wolf, Wolf Variant, Wolf Dry, and Wyotite.

Parent Material Kind: alluvium and residuum

Parent Material Origin: sandstone, shale

Surface Texture: loam, silt loam, very fine sandy loam

Surface Texture Modifier: none is most common but gravelly or cobbly may occur

Subsurface Texture Group: loam

Surface Fragments ≤ 3" (% Cover): 0

Surface Fragments > 3" (%Cover): typically 0, occasionally up to 10

Subsurface Fragments ≤ 3" (% Volume): typically 0, occasionally up to 15

Subsurface Fragments > 3" (% Volume): typically 0, occasionally up to 10

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	moderately well drained	well drained
Permeability Class:	moderately slow	moderate
Depth (inches):	20	>60
Electrical Conductivity (mmhos/cm) ≤20":	0	4
Sodium Absorption Ratio ≤20":	0	5

Site Type: Rangeland
MLRA: 58B – Northern Rolling High Plains

**Loamy 10-14" P.Z.
R058BY122WY**

Soil Reaction (1:1 Water) $\leq 20''$:	6.6	8.4
Soil Reaction (0.1M CaCl₂) $\leq 20''$:	NA	NA
Available Water Capacity (inches) $\leq 30''$:	3.0	6.3
Calcium Carbonate Equivalent (percent) $\leq 20''$:	0	10

Plant Communities

Ecological Dynamics of the Site:


As this site deteriorates because of a combination of frequent and severe grazing, species such as blue grama and big sagebrush will increase. Cool-season grasses such as green needlegrass, needleandthread, and rhizomatous wheatgrasses will decrease in frequency and production.

Big sagebrush may become dominant on some areas with an absence of fire. Wildfires are actively controlled in recent times so chemical control using herbicides has replaced the historic role of fire on this site. Recently, prescribed burning has regained some popularity.

Due to the amount and pattern of the precipitation, the big sagebrush component typically is not resilient once it has been removed if a healthy and vigorous stand of grass exists and is maintained. The exception to this is where the herbaceous component is severely degraded at the time of treatment, growing conditions are unfavorable after treatment, and/or recovery periods are inadequate.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



- BM** - Brush Management (fire, chemical, mechanical)
- Freq. & Severe Grazing** - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season
- GLMT** - Grazing Land Mechanical Treatment
- LTPG** - Long-term Prescribed Grazing
- MCSLG** - Moderate, Continuous Season-long Grazing
- NU, NF** - No Use and No Fire
- PG** - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)
- VLTPG** - Very Long-term Prescribed Grazing (could possibly take generations)
- Na** - found adjacent to a saline site

PLANT COMMUNITY DYNAMICS
REFERENCE PLANT COMMUNITY

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Grp	Allowable Annual Production			% Comp (MAX.)
				lbs./acre			
				below normal	normal	above normal	
				700	1200	1500	
GRASSES/GRASSLIKES							
RHIZOMATOUS WHEATGRASSES:							
thickspike wheatgrass	Elymus lanceolatus	ELLAL	1	175	300	375	25%
western wheatgrass	Pascopyrum smithii	PASM	1	175	300	375	25%
OTHER GRASSES							
green needlegrass	Nassella viridula	NAV14	2	105	180	225	15%
needleandthread	Hesperostipa comata	HECO26	3	175	300	375	25%
Cusick's bluegrass	Poa cusickii	POCU3	4	70	120	150	10%
blue grama	Bouteloua gracilis	BOGR2	5	105	180	225	15%
MISCELLANEOUS GRASSES/GRASSLIKES*							
bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6	6	35	60	75	5%
hairy grama	Bouteloua hirsuta	BOHI2	6	35	60	75	5%
Indian ricegrass	Achnatherum hymenoides	ACHY	6	35	60	75	5%
plains reedgrass	Calamagrostis montanensis	CAMO	6	35	60	75	5%
needleleaf sedge	Carex duriuscula	CADU6	6	35	60	75	5%
prairie junegrass	Koeleria macrantha	KOMA	6	35	60	75	5%
Sandberg bluegrass	Poa secunda	POSE	6	35	60	75	5%
threadleaf sedge	Carex filifolia	CAFI	6	35	60	75	5%
FORBS							
MISCELLANEOUS FORBS*							
American vetch	Vicia americana	VIAM	7	105	180	225	15%
prairie coneflower	Ratibida columnifera	RACO3	7	35	60	75	5%
asters	Asters	ASTER	7	35	60	75	5%
biscuitroots	Lomatium spp.	LOMAT	7	35	60	75	5%
breadroot scurfpea	Pediomelum esculentum	PEES	7	35	60	75	5%
western yarrow	Achillea lanulosa	ACHIL	7	35	60	75	5%
rosy pussytoes	Antennaria rosea	ANRO2	7	35	60	75	5%
milkvetches	Astragalus	ASTRA	7	35	60	75	5%
stemless goldenweed	Haplopappus acaulis	HAAC	7	35	60	75	5%
sulphur flower buckwheat	Eriogonum umbellatum	ERUM	7	35	60	75	5%
scarlet gaura	Gaura coccinea	GACO5	7	35	60	75	5%
purple prairie clover	Dalea purpurea	DAPU5	7	35	60	75	5%
white prairie clover	Dalea candida	DACA7	7	35	60	75	5%
bluebells	Mertensia	MERTE	7	35	60	75	5%
wild onion	Allium textile	ALTE	7	35	60	75	5%
hawksbeard	Crepis acuminata	CRAC2	7	35	60	75	5%
TREES, SHRUBS & HALF-SHRUBS							
big sagebrush	Artemisia tridentata	ARTR2	8	70	120	150	10%
winterfat	Krascheninnikovia lanata	KRLA2	9	35	60	75	5%

* Common native perennials are listed. Other native perennials may also be counted but no species in the group may be counted for more than 5%.

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

Rhizomatous Wheatgrasses, Needleandthread, Blue Grama Plant Community

This plant community is the interpretive plant community for this site and is considered to be the Historic Climax Plant Community (HCPC). This plant community evolved with grazing by large herbivores and is well suited for grazing by domestic livestock. This plant community can be found on areas that are properly managed with grazing and/or prescribed burning, and sometimes on areas receiving occasional short periods of rest. The potential vegetation is about 75% grasses or grass-like plants, 15% forbs, and 10% woody plants. This state is dominated by cool season mid-grasses.

The major grasses include western wheatgrass, needleandthread, and green needlegrass. Other grasses occurring in this state include Cusick’s and Sandberg’s bluegrass, bluebunch wheatgrass, and blue grama. A variety of forbs and half-shrubs also occur, as shown in the preceding table. Big sagebrush is a conspicuous element of this state, occurs in a mosaic pattern, and makes up 5 to 10% of the annual production. Plant diversity is high.

The total annual production (air-dry weight) of this state is about 1,200 lbs./acre, but it can range from about 700 lbs./acre in unfavorable years to about 1,500 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This plant community is extremely stable and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- No use and no fire for 20 years or more will convert this plant community to the *Heavy Sagebrush Plant Community*.
- Moderate, continuous season-long grazing will convert the plant community to the *Mixed Sagebrush/Grass Plant Community*.

- Moderate continuous season-long grazing, where greasewood occurs adjacent to the site, will convert the plant community to the *Greasewood Plant Community*.
- When cropped annually and then abandoned without reseeding, the site is converted to the *Go-back Land Plant Community*.

Mixed Sagebrush/Grass Plant Community

Historically, this plant community evolved under grazing by bison and a low fire frequency. Currently, it is found under moderate, season-long grazing by livestock in the absence of fire or brush management. Wyoming big sagebrush is a significant component of this plant community. Cool-season grasses make up the majority of the understory with the balance made up of short warm-season grasses, annual cool-season grasses, and miscellaneous forbs.

Dominant grasses include needleandthread, western wheatgrass, and green needlegrass. Grasses of secondary importance include blue grama, prairie junegrass, and Sandberg bluegrass. Forbs commonly found in this plant community include plains wallflower, hairy goldaster, slimflower scurfpea, and scarlet globemallow. Sagebrush canopy ranges from 20% to 30%. Fringed sagewort is commonly found. Plains pricklypear can also occur.

When compared to the Historic Climax Plant Community, sagebrush and blue grama have increased. Production of cool-season grasses, particularly green needlegrass, has been reduced. The sagebrush canopy protects the cool-season mid-grasses, but this protection makes them unavailable for grazing. Cheatgrass (downy brome) has invaded the site. The overstory of sagebrush and understory of grass and forbs provide a diverse plant community that will support domestic livestock and wildlife such as mule deer and antelope.

The total annual production (air-dry weight) of this state is about 900 pounds per acre, but it can range from about 700 lbs./acre in unfavorable years to about 1,200 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change. A significant reduction of big sagebrush can only be accomplished through fire or brush management. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term.

Transitions or pathways leading to other plant communities are as follows:

- Brush management (chemical, fire, or mechanical), followed by prescribed grazing, will convert this plant community to the *Rhizomatous wheatgrasses, Needleandthread, Blue grama Plant Community*. The probability of this occurring is high. When prescribed fire is used, sufficient fine fuels will need to be present. This may require deferment from grazing prior to treatment. Post management is critical to ensure success. This can range from two or more years of rest to partial growing season deferment, depending on the condition of the understory at the time

of treatment and the growing conditions following treatment. In the case of an intense wildfire that occurs when desirable plants are not completely dormant, the length of time required to reach the *Rhizomatous wheatgrasses, Needleandthread, Blue grama Plant Community* may be increased.

- Brush management, followed by frequent and severe grazing, will convert the plant community to the *Western Wheatgrass/Cheatgrass Plant Community*. The probability of this occurring is high. If bare areas exist after treatment, along with no recovery periods from grazing, cheatgrass will invade and plants not as resistant to grazing as western wheatgrass will be reduced.
- Moderate continuous season-long grazing, where greasewood occurs adjacent to this state, will convert the plant community to the *Greasewood Plant Community*.

Heavy Sagebrush Plant Community

This plant community is the result of long-term protection from grazing and fire. Sagebrush eventually dominates this plant community with canopy cover often exceeding 60%. At first, excessive litter builds up, shading out some of the grasses and forbs. Other plants become decadent with low vigor. Bunch grasses often develop dead centers. Eventually, the interspaces between plants increase in size leaving more soil surface exposed. Organic matter oxidizes in the air rather than being incorporated into the soil.

The dominant plants tend to be somewhat similar to those found in the Historic Climax Plant Community. Weedy species, cool-season grasses, and sedges have increased. Blue grama has decreased. Rodent activity has resulted in an increase in soil disturbance. Cactus and sageworts often increase. Noxious weeds such as Dalmatian toadflax, leafy spurge, or Canada thistle may invade the site if a seed source is present. Plant diversity is moderate to high.

The total annual production (air-dry weight) of this state is about 800 pounds per acre, but it can range from about 600 lbs./acre in unfavorable years to about 1,000 lbs./acre in above average years.

The following is the growth curve of the plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This plant community is not resistant to change and is more vulnerable to severe disturbance than the HCPC. The introduction of grazing or fire quickly changes the plant community.

Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestaling are obvious. Infiltration is reduced and runoff is increased.

Transitions or pathways leading to other plant communities are as follows:

- Brush management, followed by prescribed grazing, will return this plant community to at or near the *Rhizomatous Wheatgrasses, Needleandthread, Blue Grama Plant Community*.

- Brush management, followed by frequent and severe grazing, will convert the plant community to the *Western Wheatgrass/Cheatgrass Plant Community*. The probability of this occurring is high because of the amount of bare ground exposed to cheatgrass invasion.

Western Wheatgrass/Cheatgrass Plant Community

This plant community is created when the Mixed Sagebrush/Grass Plant Community or the Heavy Sagebrush Plant Community is subjected to fire or brush management not followed by prescribed grazing. Rhizomatous wheatgrasses and annuals will eventually dominate the site.

Compared to the HCPC, cheatgrass has invaded with western wheatgrass and thickspike wheatgrass maintaining at a similar or slightly higher level. Virtually all other cool-season mid-grasses are severely decreased. Blue grama is the same or slightly less than found in the HCPC. Plant diversity is low.

The total annual production (air-dry weight) of this state is about 600 pounds per acre, but it can range from about 450 lbs./acre in unfavorable years to about 750 lbs./acre in above average years.

The following is the growth curve of the plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This plant community is relatively stable with the rhizomatous wheatgrasses being somewhat resistant to overgrazing and the cheatgrass effectively competing against the establishment of perennial cool-season grasses.

An increase in bare ground reduces water infiltration and increases soil erosion. The watershed is usually functioning. The biotic integrity is reduced by the lack of diversity in the plant community.

Transitions or pathways leading to other plant communities are as follows:

- Moderate continuous season-long grazing will eventually return this plant community to the *Mixed Sagebrush/Grass Plant Community*.
- Frequent and severe grazing will convert this plant community to *Blue Grama Sod Plant Community*.
- Frequent and severe yearlong grazing will convert this plant community to *Blue grama, Plains Pricklypear, Bare Ground Plant Community*.
- Long-term, prescribed grazing will eventually return this plant community to at or near the *Rhizomatous Wheatgrasses, Needleandthread, Blue Grama Plant Community*.

Blue Grama Sod Plant Community

This plant community is the result of frequent and severe grazing during the growing season of the cool-season mid-grasses. A dense sod of blue grama dominates it. Pricklypear cactus can become dense enough so that livestock cannot graze forage growing within the cactus clumps.

Site Type: Rangeland
MLRA: 58B – Northern Rolling High Plains

**Loamy 10-14” P.Z.
R058BY122WY**

When compared to the Historic Climax Plant Community, blue grama and threadleaf sedge have increased. All cool-season mid-grasses and forbs have been greatly reduced. Plant diversity is extremely low.

The total annual production (air-dry weight) of this state is about 600 pounds per acre, but it can range from about 450 lbs./acre in unfavorable years to about 750 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	15	30	25	15	5	0	0	0

(Monthly percentages of total annual growth)

This sod bound plant community is very resistant to water infiltration. While this sod protects the site itself, off-site areas are affected by excessive runoff that can cause gully erosion. This sod is very resistant to change and may require a grazing land mechanical treatment, such as chiseling, to return the cool-season grass component.

Transitions or pathways leading to other plant communities are as follows:

- Grazing land mechanical treatment (chiseling, etc.) and pricklypear cactus control (if needed), followed by prescribed grazing, will return this plant community to near *Historic Climax Plant Community* condition.
- Grazing land mechanical treatment, followed by frequent and severe grazing, will convert this plant community to the *Western Wheatgrass/Cheatgrass Plant Community*.
- Frequent and severe yearlong grazing will eventually convert this state to the *Blue Grama, Plains Pricklypear, Bare Ground Plant Community*.

Greasewood Plant Community

This plant community can occur where states are subjected to continuous season-long grazing at moderate stocking rates and where greasewood occurs adjacent to the site. It is dominated by an overstory of greasewood and possibly big sagebrush. Rhizomatous wheatgrasses, cheatgrass, and inland saltgrass make up the understory. Salts in the surface will increase due to the shedding of the salt-filled leaves of the greasewood. Plant diversity is high.

The total annual production (air-dry weight) of this state is about 700 pounds per acre, but it can range from about 525 lbs./acre in unfavorable years to about 875 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change. A significant reduction of greasewood can only be accomplished through repeated brush control treatments. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term.

The site is protected from erosion as long as ground cover is maintained. The biotic integrity of this state is somewhat intact because of the woody overstory and perennial grass understory. The watershed is functioning as long as a grass cover is maintained.

- Recovery to near *Historic Climax Plant Community* condition is difficult due to the resistance of greasewood to herbicides and accumulated effects of salts on the soil.

Blue Grama/ Plains Pricklypear/ Bare Ground Plant Community

This plant community is the result of frequent and severe yearlong grazing over the long-term. Perennial plants are decreased. Cheatgrass, annual weeds, and bare ground are increased. Plains pricklypear may have increased, rendering much of the forage unusable by livestock.

This plant community is highly variable depending on the severity, frequency, and duration of the grazing and also the condition of the plant community when this level of grazing began. Virtually all plants not resistant to overgrazing may have been eliminated. Dominant plants may include blue grama, threeawns, annuals, and, to a lesser degree, rhizomatous wheatgrasses. Perennial plant diversity is low.

The total annual production (air-dry weight) of this state is about 500 pounds per acre, but it can range from about 375 lbs./acre in unfavorable years to about 625 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:
 Growth curve name:
 Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	15	30	25	15	5	0	0	0

(Monthly percentages of total annual growth)

This state is unhealthy and subject to increased erosion. Runoff is high on this state due to the sod nature of blue grama and bare ground.

Transitions or pathways leading to other plant communities are as follows:

- Long-term prescribed grazing will convert this plant community initially to the *Blue Grama Sod Plant Community*, when this state is dominated by blue grama sod at the time of treatment.
- Long-term prescribed grazing will convert this plant community to the *Western Wheatgrass /Cheatgrass Plant Community*, when this state has large amounts of cheatgrass, annual weeds, and bare ground at the time of treatment. Control of plains pricklypear cactus may be necessary.

Reseeding areas with native plant species and proper grazing management may be necessary to accelerate recovery where few desirable plants remain.

Go-back Land Plant Community

This plant community occurs on land that has been cropped annually in the past and then abandoned without reseeding. Natural succession has resulted in a plant community dominated by varying combinations of red threeawn, cheatgrass, blue grama, Sandberg bluegrass, and some rhizomatous wheatgrasses. Forage production is low and grasses such as red threeawn and cheatgrass are not used efficiently by livestock.

The total annual production (air-dry weight) of this state is about 600 pounds per acre, but it can range from about 500 lbs./acre in unfavorable years to about 900 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	0	0	0	0	0	0	0	0

(Monthly percentages of total annual growth)

The potential for accelerated erosion can be highly variable depending on amount of bare ground present. Biological diversity is low.

Transitions or pathways leading to other plant communities are as follows:

- Prescribed grazing may be used to increase desirable native cool-season grass production. It is usually difficult to return to near *Historic Climax Plant Community* condition in a timely manner because of past soil loss.
- Grazing land mechanical treatment (i.e., chiseling) may improve forage production where significant rhizomatous wheatgrass is present to respond.

Where there is a lack of perennial grasses, reseeding to tame or native species may be necessary to return these lands to production in the form of pastureland. These pastures are normally seeded to crested wheatgrass, pubescent wheatgrass, or Russian wildrye. They require considerable investment to establish and have a variable life expectancy. They do produce up to 50% more than native range, but their value as forage is somewhat limited due to the single species usually seeded. In some cases, the single species or certain groups of species (e.g., wheatgrasses) may be more vulnerable to infestation by associated insects and/or diseases (e.g., black grass bugs).

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

Rhizomatous Wheatgrasses, Needleandthread, Blue Grama Plant Community (HCPC): The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant

community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

Mixed Sagebrush/Grass Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants, and hosts of other nesting birds utilize stands in the 20-30% cover range.

Heavy Sagebrush Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15% protein and 40-60% digestibility during that time. This community provides excellent escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse.

Western Wheatgrass/Cheatgrass Plant Community: This plant community may be useful for the same large grazers that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for sage grouse when it occurs proximal to woody cover. Good grasshopper habitat equals good foraging for birds.

Blue Grama Sod and Go-back Land Plant Communities: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Western Wheatgrass/Cheatgrass Plant Community is limiting. Generally, these are not target plant communities for wildlife habitat management.

Greasewood Plant Community: This plant community exhibits a low level of plant species diversity due to the accumulation of salts in the soil. It may provide some thermal and escape cover for deer and antelope if no other woody community is nearby, but in most cases it is not a desirable plant community to select as a wildlife habitat management objective.

Blue Grama, Plains Pricklypear, Bare Ground Plant Community: Benefits to other wildlife are largely due to the subterranean structure created by the prairie dogs, not the sparse vegetation found on this plant community.

Introduced Pasture: These communities are highly variable depending on the species planted. Refer to Forage Suitability Groups for more information.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 58B, 10-14 inch Northern Plains

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope
GRASSES/GRASSLIKES							
alkali bluegrass	<i>Poa secunda ssp. juncifolia</i>	POSEJ	DDDD	PPPP	DDDD	PPPP	PPPP
alkali cordgrass	<i>Spartina gracilis</i>	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU
alkali sacaton	<i>Sporobolus airoides</i>	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD
Baltic rush	<i>Juncus balticus</i>	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU
basin wildrye	<i>Leymus cinereus</i>	LEC14	PPPP	PPPP	PPPP	DDDD	DDDD
bearded wheatgrass	<i>Elymus caninus</i>	ELCA	PPPP	DDDD	PPPP	DDDD	DDDD
big bluestem	<i>Andropogon gerardii</i>	ANGE	PPPP	PPPP	PPPP	DDDD	DDDD
blue grama	<i>Bouteloua gracilis</i>	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD
bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD
bluejoint reedgrass	<i>Calamagrostis canadensis</i>	CACA4	PPPP	DDDD	PPPP	UUUU	UUUU
bottlebrush squirreltail	<i>Elymus elymoides</i>	ELELE	DDDD	DDDD	DDDD	UUUU	UUUU
buffalograss	<i>Buchloe dactyloides</i>	BUDA	DDDD	DDDD	DDDD	DDDD	DDDD
Canada wildrye	<i>Elymus canadensis</i>	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD
Canby bluegrass	<i>Poa canbyi (syn. to Poa secunda)</i>	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
Cusick's bluegrass	<i>Poa cusickii</i>	POCU3	PPPP	PPPP	PPPP	PPPP	PPPP
Fendler threeawn	<i>Aristida purpurea</i>	ARPUL	UUUU	UUUU	UUUU	UUUU	UUUU
green needlegrass	<i>Nassella viridula</i>	NAVI4	PPPP	PPPP	PPPP	PPPP	PPPP
hairy grama	<i>Bouteloua hirsuta</i>	BOH12	DDDD	DDDD	DDDD	DDDD	DDDD
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP
inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU
inland sedge	<i>Carex interior</i>	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU
little bluestem	<i>Schizachyrium scoparium</i>	SCSC	PPPP	PPPP	PPPP	DDDD	DDDD
mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	<i>Carex nebraskensis</i>	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD
needleandthread	<i>Hesperostipa comata</i>	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP
needleleaf sedge	<i>Carex durivuscula</i>	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU
northern reedgrass	<i>Calamagrostis stricta</i>	CAST13	PPPP	DDDD	PPPP	UUUU	UUUU
Nuttall's alkaligrass	<i>Puccinellia nuttaliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP
plains muhly	<i>Muhlenbergia cuspidata</i>	MUCU3	DDDD	DDDD	DDDD	UUUU	UUUU
plains reedgrass	<i>Calamagrostis montanensis</i>	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD
prairie cordgrass	<i>Spartina pectinata</i>	SPPE	PPPP	DDDD	PPPP	UUUU	UUUU
prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD
prairie sandreed	<i>Calamovilfa longifolia</i>	CALO	PPPP	DDDD	PPPP	UUUU	UUUU
sand bluestem	<i>Andropogon halli</i>	ANHA	PPPP	DDDD	PPPP	UUUU	UUUU
sand dropseed	<i>Sporobolus cryptandrus</i>	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU
Sandberg bluegrass	<i>Poa secunda</i>	POSE	DDDD	DDDD	DDDD	DDDD	DDDD
sideoats grama	<i>Bouteloua curtipendula</i>	BOCU	PPPP	PPPP	PPPP	DDDD	UUUU
slender wheatgrass	<i>Elymus trachycaulus</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD
spike sedge	<i>Carex nardina</i>	CANA2	DDDD	DDDD	DDDD	UUUU	UUUU
thickspike wheatgrass	<i>Elymus lanceolatus</i>	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD
threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP
tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD
western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DDDD	DDDD	DDDD	DDDD	DDDD
FORBS							
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU
American vetch	<i>Vicia americana</i>	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP
arrowgrass	<i>Triglochin spp.</i>	TRIGL	T	T	T	T	T
asters	Asters	ASTER	UUUU	UUUU	UUUU	UUUU	UUUU
biscuitroots	<i>Lomatium spp.</i>	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD
bluebells	<i>Mertensia</i>	MERTE	DDDD	PPPP	DDDD	DDDD	DDDD
blue-eyed grass	<i>Sisyrinchium spp.</i>	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD
breadroot scurfpea	<i>Pediomelum esculentum</i>	PEES	DDDD	DDDD	DDDD	DDDD	DDDD
cattail, broad-leaf	<i>Typha latifolia</i>	TYLA	DDDD	UUUU	DDDD	UUUU	UUUU
cattail, narrow-leaf	<i>Typha angustifolia</i>	TYAN	DDDD	UUUU	DDDD	UUUU	UUUU
fringed sagewort	<i>Artemisia frigida</i>	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU
green sagewort	<i>Artemisia dracunculul</i>	ARDR4	UUUU	UUUU	UUUU	UUUU	UUUU
hawkbeard	<i>Crepis acuminata</i>	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD
horsetails	<i>Equisetum spp.</i>	EQUIS	UUUU	UUUU	UUUU	UUUU	UUUU
iris	<i>Iris spp.</i>	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU
milkvetches	<i>Astragalus</i>	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD
poison hemlock	<i>Conium maculatum</i>	COMA2	T	T	T	T	T
prairie coneflower	<i>Ratibida columnifera</i>	RACO3	DDDD	PPPP	DDDD	PPPP	PPPP
prairie thermopsis	<i>Thermopsis rhombifolia</i>	THRHA	UUUU	UUUU	UUUU	UUUU	UUUU
purple prairie clover	<i>Dalea purpurea</i>	DAPU5	PPPP	PPPP	PPPP	PPPP	PPPP
Pursh seepweed	<i>Suaeda calceoliformis</i>	SUCA2	UUUU	UUUU	UUUU	UUUU	UUUU
rosy pussytoes	<i>Antennaria rosea</i>	ANRO2	UUUU	UUUU	UUUU	UUUU	UUUU
scarlet gaura	<i>Gaura coccinea</i>	GACO5	UUUU	UUUU	UUUU	UUUU	UUUU
stemless goldenweed	<i>Haplopappus acaulis</i>	HAAC	UUUU	UUUU	UUUU	UUUU	UUUU
sulphur flower buckwheat	<i>Eriogonum umbellatum</i>	ERUM	UUUU	UUUU	UUUU	UUUU	UUUU
twogrooved milkvetch	<i>Astragalus bisulcatus</i>	ASB12	T	T	T	T	T
water hemlocks	<i>Cicuta spp.</i>	CICUT	T	T	T	T	T
western yarrow	<i>Achillea lanulosa</i>	ACHIL	UUUU	UUUU	UUUU	UUUU	UUUU
white prairie clover	<i>Dalea candida</i>	DACA7	PPPP	PPPP	PPPP	PPPP	PPPP
wild onion	<i>Allium textile</i>	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD
TREES, SHRUBS & HALF-SHRUBS							
big sagebrush	<i>Artemisia tridentata</i>	ARTR2	UUUU	DDDD	UUUU	DDDD	DDDD
birdfoot sagebrush	<i>Artemisia pedatifida</i>	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU
black greasewood	<i>Sarcobatus vermiculatus</i>	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD
fourwing saltbush	<i>Atriplex canescens</i>	ATCA2	PPPP	PPPP	PPPP	PPPP	PPPP
Gardners saltbush	<i>Atriplex gardneri</i>	ATGA	PPPP	PPPP	DDDD	PPPP	PPPP
green rabbitbrush	<i>Chrysothamnus viscidiflorous</i>	CHV18	DDDD	DDDD	DDDD	DDDD	DDDD
junipers	<i>Juniperus scopulorum</i>	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU
plains cottonwood (sprouts)	<i>Populus deltoides</i>	PODEM	DDDD	DDDD	DDDD	DDDD	DDDD
ponderosa pine (abortion in cattle)	<i>Pinus ponderosa</i>	PIPO	UUUU	UUUU	UUUU	UUUU	UUUU
rubber rabbitbrush	<i>Ericameria nauseosa</i>	ERNA10	UUUU	DDDD	UUUU	DDDD	DDDD
silver sagebrush	<i>Artemisia cana</i>	ARCA5	DDDD	DDDD	DDDD	PPPP	PPPP
silverberry	<i>Eleagnus commutata</i>	ELCO	UUUU	UUUU	UUUU	DDDD	UUUU
skunkbush sumac	<i>Rhus trilobata</i>	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD
western snowberry	<i>Symphoricarpos occidentalis</i>	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU
wildrose	<i>Rosa woodsii var. woodsii</i>	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD
willows	<i>Salix L.</i>	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU
winterfat	<i>Krascheninnikovia lanata</i>	KRLA2	PPPP	PPPP	PPPP	PPPP	PPPP
yucca	<i>Yucca glauca</i>	YUGL	DDDD	DDDD	DDDD	DDDD	DDDD

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (lb./ac)	Carrying Capacity* (AUM/ac)
Rhizomatous WG, Needleandthread, Blue Grama	700-1500	.4
Heavy Sagebrush	800-1400	.3
Blue Grama Sod	400-1000	.2
Mixed Sagebrush/Grass	700-1200	.33
Western Wheatgrass/Cheatgrass	600-1200	.2
Blue grama, Plains Pricklypear, Bare ground	300-800	.1
Greasewood	525-875	.3
Go-back Land	500-900	.2

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderately slow to moderate. Runoff potential for this site varies from low to moderate depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Site Type: Rangeland
MLRA: 58B – Northern Rolling High Plains

**Loamy 10-14” P.Z.
R058BY122WY**

Recreational Uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood Products

No appreciable wood products are present on the site.

Other Products

None noted.

Supporting Information

Associated Sites

Shallow Loamy	058BY162WY
Sandy	058BY150WY
Clayey	058BY104WY
Overflow	058BY130WY
Lowland	058BY128WY

Similar Sites

() – Loamy 15-17” Northern Plains P.Z. 058BY222WY
[higher production]

Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel was also used. Those involved in developing this site include: Glen Mitchell, Range Management Specialist, NRCS; Chuck Ring, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417	12	1971-1994	WY	Campbell & others
Ocular estimates	5	1990-1999	WY	Campbell & others

Site Correlation

This site has been correlated with Montana in MLRA 58B.

Type Locality

Field Offices

Buffalo, Douglas, Gillette, Lusk, Newcastle, Sheridan

Site Type: Rangeland
MLRA: 58B – Northern Rolling High Plains

**Loamy 10-14” P.Z.
R058BY122WY**

Relationship to Other Established Classifications

Other References

Site Description Approval

State Range Management Specialist

Date

State Range Management Specialist

Date