

## United States Department of Agriculture Natural Resources Conservation Service

### Ecological Site Description

**Site Type:** Rangeland

**Site Name:** Saline Lowland (SL) 10-14” Precipitation Zone,

**Site ID:** 058BY138WY

**Major Land Resource Area:** 58B – Northern Rolling High Plains

### Physiographic Features

This site normally occurs on land that receives overflow from intermittent streams or runoff from adjacent slopes.

**Landform:** alluvial fans, drainage ways & stream terraces      **Aspect:** N/A

	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	3800	5100
<b>Slope (percent):</b>	0	6
<b>Water Table Depth (inches):</b>	None within 60 inches	
<b>Flooding:</b>		
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Ponding:</b>		
<b>Depth (inches):</b>	0	0
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Runoff Class:</b>	negligible	low

### Climatic features

Annual precipitation ranges from 10-14 inches per year. Wide fluctuations may occur in yearly precipitation and result in more drought years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Wind speed averages about 8 mph, ranging from 10 mph during the spring to 7 mph during late summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 75 mph.

Growth of native cool season plants begins about April 1 and continues to about July 1. Native warm season plants begin growth about May 15 and continue to about August 15. Green up of cool season plants may occur in September and October of most years.

Site Type: Rangeland  
MLRA: 58B – Northern Rolling High Plains

**Saline Lowland 10-14” P.Z.  
R058BY138WY**

The following information is from the “Clearmont 5 SW” climate station:

Frost-free period (32 °F): 76 - 132 days; (5 yrs. out of 10, these days will occur between May 30 – September 11)

Freeze-free period 28 °F): 110 - 145 days; (5 yrs. out of 10, these days will occur between May 16 – September 21)

Mean annual precipitation: 12.4 inches

Mean annual air temperature: 43.2 °F (28.4°F Avg. Min. – 57.9°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include: “Dull Center”

## Influencing Water Features

<b>Wetland Description:</b>	<b><u>System</u></b>	<b><u>Subsystem</u></b>	<b><u>Class</u></b>	<b><u>Sub-class</u></b>
None	None	None	None	None

**Stream Type:** None

## Representative Soil Features

The soils of this site are deep and very deep well-drained soils formed in alluvium. Layers of the soil most influential to the plant community vary from 3 to 6 inches thick. These soils have moderate to slow permeability and are moderately to strongly saline and/or alkaline. Higher soluble salt concentrations may be found in the subsoils. The surface soil will be highly variable and vary from 2 to 8 inches in thickness. The surface soil will be one or more of the following textures: very fine sandy loam, fine sandy loam, sandy loam, loam, silt loam, clay loam, clay, or silty clay. A fluctuating water table occurs in these areas and ranges from 1 to 5 feet. These areas are subject to occasional overflow.

Major Soil Series correlated to this site include: Boruff, Clarkelen, Draknab, Haverdad, Lohmiller,

Other Soil Series correlated in MLRA 58B to this site include: Bankard, Barnum, Bigwinder, Coaliams, Colombo, Connerton, Docpar, Emigha, Glenberg, Haverson, Havertel, Livan, Redbank, and Worthenton

**Parent Material Kind:** alluvium

**Parent Material Origin:** sandstone, shale

**Surface Texture:** loam, clay loam, silt loam, clay

**Surface Texture Modifier:** none

**Subsurface Texture Group:** loam

**Surface Fragments ≤ 3” (% Cover):** 0

**Surface Fragments > 3” (%Cover):** 0

**Subsurface Fragments ≤ 3” (% Volume):** 0

**Subsurface Fragments > 3” (% Volume):** 0

	<b><u>Minimum</u></b>	<b><u>Maximum</u></b>
<b>Drainage Class:</b>	poorly	excessive
<b>Permeability Class:</b>	moderate	rapid
<b>Depth (inches):</b>	20	>60
<b>Electrical Conductivity (mmhos/cm) ≤20”:</b>	4	16
<b>Sodium Absorption Ratio ≤20”:</b>	5	13

Site Type: Rangeland  
MLRA: 58B – Northern Rolling High Plains

**Saline Lowland 10-14" P.Z.  
R058BY138WY**

<b>Soil Reaction (1:1 Water) <math>\leq 20''</math>:</b>	6.6	9.0
<b>Soil Reaction (0.1M CaCl<sub>2</sub>) <math>\leq 20''</math>:</b>	NA	NA
<b>Available Water Capacity (inches) <math>\leq 30''</math>:</b>	1	6.2
<b>Calcium Carbonate Equivalent (percent) <math>\leq 20''</math>:</b>	0	10


## **Plant Communities**

### **Ecological Dynamics of the Site:**

As this site deteriorates, species such as inland saltgrass and greasewood increase, and cheatgrass invades the site. Grasses such as alkali sacaton, rhizomatous wheatgrasses and Nuttall's alkaligrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



**BM** - Brush Management (fire, chemical, mechanical)

**Freq. & Severe Grazing** - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season

**GLMT** - Grazing Land Mechanical Treatment

**LTPG** - Long-term Prescribed Grazing

**MCSLG** - Moderate, Continuous Season-long Grazing

**NU, NF** - No Use and No Fire

**PG** - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)

**VLTPG** - Very Long-term Prescribed Grazing (could possibly take generations)

**Na** - Moderate Sodium in Soil

PLANT COMMUNITY DYNAMICS  
REFERENCE PLANT COMMUNITY

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Grp	Allowable Annual Production			% Comp (MAX.)
				lbs./acre			
				below normal 1400	normal 1700	above normal 2200	
<b>GRASSES/GRASSLIKES</b>							
<b>RHIZOMATOUS WHEATGRASSES:</b>							
thickspike wheatgrass	Elymus lanceolatus	ELLAL	1	210	255	330	15%
western wheatgrass	Pascopyrum smithii	PASM	1	210	255	330	15%
<b>OTHER GRASSES</b>							
alkali sacaton	Sporobolus airoides	SPAI	2	350	425	550	25%
alkali bluegrass	Poa secunda ssp. juncifolia	POSEJ	3	210	255	330	15%
Nuttall's alkaligrass	Puccinellia nuttaliana	PUNU2	4	140	170	220	10%
bottlebrush squirreltail	Elymus elymoides	ELEL5	5	140	170	220	10%
inland saltgrass	Distichlis spicata	DISP	6	210	255	330	15%
<b>FORBS</b>							
<b>MISCELLANEOUS FORBS*</b>							
American vetch	Vicia americana	VIAM	7	140	170	220	10%
prairie coneflower	Ratibida columnifera	RACO3	7	70	85	110	5%
asters	Asters	ASTER	7	70	85	110	5%
milkvetches	Astragalus	ASTRA	7	70	85	110	5%
scarlet gaura	Gaura coccinea	GACO5	7	70	85	110	5%
purple prairie clover	Dalea purpurea	DAPU5	7	70	85	110	5%
white prairie clover	Dalea candida	DACA7	7	70	85	110	5%
American licorice	Glycyrrhiza lepidota	GLLE3	7	70	85	110	5%
wild onion	Allium textile	ALTE	7	70	85	110	5%
woodyaster	Xylorhiza spp.	XYLOR	7	70	85	110	5%
hawksbeard	Crepis acuminata	CRAC2	7	70	85	110	5%
<b>TREES, SHRUBS &amp; HALF-SHRUBS*</b>							
black greasewood	Sarcobatus vermiculatus	SAVE4	8	280	340	440	20%
Gardners saltbush	Atriplex gardneri	ATGA	9	70	85	110	5%
fourwing saltbush	Atriplex canescens	ATCA2	10	70	85	110	5%
rubber rabbitbrush	Ericameria nauseosa	ERNA10	11	70	85	110	5%
plains cottonwood	Populus deltoides	PODEM	12	70	85	110	5%
winterfat	Krascheinnikovia lanata	KRLA2	13	140	170	220	10%

\* Common native perennials are listed. Other native perennials may also be counted but no species in the group may be counted for more than 5%.

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

**Plant Community Narratives**

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

**Rhizomatous Wheatgrasses, Alkali sacaton Plant Community**

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores and is well suited for grazing by domestic livestock. Potential vegetation is about 75% grasses or grass-like plants, 5% forbs and 20% woody plants. Saline tolerant grasses dominate the state. The major grasses include rhizomatous wheatgrasses, alkali sacaton, alkali bluegrass, Nuttall’s alkaligrass and inland saltgrass. Woody plants are greasewood, four-wing saltbush and Gardners saltbush.

The total annual production (air-dry weight) of this state is about 1700 pounds per acre, but it can range from about 1400 lbs./acre in unfavorable years to about 2200 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	30	35	10	5	5	5	0	0

(Monthly percentages of total annual growth)

This state is stable and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought resistance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Moderate, continuous season-long grazing will convert this plant community to the *Rhizomatous wheatgrasses/Greasewood Vegetation State*.
- Frequent and Severe grazing and fire will convert this plant community to the *Inland saltgrass/Cheatgrass Vegetation State*.

**Rhizomatous wheatgrasses/Greasewood Plant Community**

This plant community evolved under moderate grazing by domestic livestock. Saline resistant grasses make up the majority of the understory. Greasewood and rubber rabbitbrush have increased on the site. Dominant grasses include rhizomatous wheatgrasses, inland saltgrass, alkali bluegrass, and alkali sacaton. Forbs, commonly found in this plant community, include Louisiana sagewort (cudweed), plains wallflower, hairy goldaster, and scarlet globemallow. Greasewood canopy cover may be 20-40%.

When compared to the Historical Climax Plant Community, rhizomatous wheatgrasses and alkali sacaton have decreased, greasewood and rubber rabbitbrush have increased. The overstory of greasewood and understory of grass and forbs provide a diverse plant community that will support domestic livestock and wildlife such as birds, mule deer and antelope.

The total annual production (air-dry weight) of this state is about 1100 pounds per acre, but it can range from about 800 lbs./acre in unfavorable years to about 1400 lbs./acre in above average years.

The following is the growth curve expected during a normal year:

Growth curve number:  
 Growth curve name:  
 Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	20	30	15	10	15	5	0	0

(Monthly percentages of total annual growth)

This state is stable and protected from excessive erosion. The biotic integrity of this plant community is usually intact. The watershed is usually functioning.

Transitional pathways leading to other plant communities are as follows:

- Prescribed grazing over the long-term will result in a plant community very similar to the *Historic Climax Plant Community*, except that greasewood will persist.
- Frequent and Severe grazing with Fire will convert this plant community to the *inland saltgrass/cheatgrass Vegetation State*.

**Inland Saltgrass/Cheatgrass Plant Community**

This plant community is the result of long-term improper grazing use and fire. This state is dominated by inland saltgrass, cheatgrass and alkali bluegrass. Bare ground has increased, and production has decreased.

The total annual production (air-dry weight) of this state is about 1000 pounds per acre, but it can range from about 600 lbs./acre in unfavorable years to about 1400 lbs./acre in above average years.

The following is the growth curve expected during a normal year:

Growth curve number:  
 Growth curve name:  
 Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	20	30	15	10	15	5	0	0

(Monthly percentages of total annual growth)

The soil of this state is not well protected. The biotic integrity is compromised by cheatgrass and bare ground. The watershed is functioning but may produce excessive runoff.

Transitional pathways leading to other plant communities are as follows:

- Prescribed grazing over the long-term will return this state to near *Historic Climax Plant Community*.

## **Ecological Site Interpretations**

### **Animal Community – Wildlife Interpretations**

**Historic Climax Plant Community:** The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

**Rhizomatous wheatgrasses/greasewood:** This plant community exhibits a low level of plant species diversity due to the accumulation of salts in the soil. It may provide some thermal and escape cover for deer and antelope if no other woody community is nearby, but in most cases it is not a desirable plant community to select as a wildlife habitat management objective.

**Inland saltgrass/Cheatgrass:** This plant community may be useful for the same large grazers that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for sage grouse when it occurs proximal to woody cover.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 58B, 10-14 inch Northern Plains

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope
<b>GRASSES/GRASSLIKES</b>							
alkali bluegrass	<i>Poa secunda ssp. juncifolia</i>	POSEJ	DDDD	PPPP	DDDD	PPPP	PPPP
alkali cordgrass	<i>Spartina gracilis</i>	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU
alkali sacaton	<i>Sporobolus airoides</i>	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD
Baltic rush	<i>Juncus balticus</i>	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU
basin wildrye	<i>Leymus cinereus</i>	LEC14	PPPP	PPPP	PPPP	DDDD	DDDD
bearded wheatgrass	<i>Elymus caninus</i>	ELCA	PPPP	DDDD	PPPP	DDDD	DDDD
big bluestem	<i>Andropogon gerardii</i>	ANGE	PPPP	PPPP	PPPP	DDDD	DDDD
blue grama	<i>Bouteloua gracilis</i>	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD
bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD
bluejoint reedgrass	<i>Calamagrostis canadensis</i>	CACA4	PPPP	DDDD	PPPP	UUUU	UUUU
bottlebrush squirreltail	<i>Elymus elymoides</i>	ELELE	DDDD	DDDD	DDDD	UUUU	UUUU
buffalograss	<i>Buchloe dactyloides</i>	BUDA	DDDD	DDDD	DDDD	DDDD	DDDD
Canada wildrye	<i>Elymus canadensis</i>	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD
Canby bluegrass	<i>Poa canbyi (syn. to Poa secunda)</i>	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
Cusick's bluegrass	<i>Poa cusickii</i>	POCUC3	PPPP	PPPP	PPPP	PPPP	PPPP
Fendler threeawn	<i>Aristida purpurea</i>	ARPUL	UUUU	UUUU	UUUU	UUUU	UUUU
green needlegrass	<i>Nassella viridula</i>	NAV14	PPPP	PPPP	PPPP	PPPP	PPPP
hairy grama	<i>Bouteloua hirsuta</i>	BOH12	DDDD	DDDD	DDDD	DDDD	DDDD
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP
inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU
inland sedge	<i>Carex interior</i>	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU
little bluestem	<i>Schizachyrium scoparium</i>	SCSC	PPPP	PPPP	PPPP	DDDD	DDDD
mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	<i>Carex nebraskensis</i>	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD
needleandthread	<i>Hesperostipa comata</i>	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP
needleleaf sedge	<i>Carex durivuscula</i>	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU
northern reedgrass	<i>Calamagrostis stricta</i>	CAST13	PPPP	DDDD	PPPP	UUUU	UUUU
Nuttall's alkaligrass	<i>Puccinellia nuttalliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP
plains muhly	<i>Muhlenbergia cuspidata</i>	MUCU3	DDDD	DDDD	DDDD	UUUU	UUUU
plains reedgrass	<i>Calamagrostis montanensis</i>	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD
prairie cordgrass	<i>Spartina pectinata</i>	SPPE	PPPP	DDDD	PPPP	UUUU	UUUU
prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD
prairie sandreed	<i>Calamovilfa longifolia</i>	CALO	PPPP	DDDD	PPPP	UUUU	UUUU
sand bluestem	<i>Andropogon halli</i>	ANHA	PPPP	DDDD	PPPP	UUUU	UUUU
sand dropseed	<i>Sporobolus cryptandrus</i>	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU
Sandberg bluegrass	<i>Poa secunda</i>	POSE	DDDD	DDDD	DDDD	DDDD	DDDD
sideoats grama	<i>Bouteloua curtipendula</i>	BOCU	PPPP	PPPP	PPPP	DDDD	UUUU
slender wheatgrass	<i>Elymus trachycaulus</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD
spike sedge	<i>Carex nardina</i>	CANA2	DDDD	DDDD	DDDD	UUUU	UUUU
thickspike wheatgrass	<i>Elymus lanceolatus</i>	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD
threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP
tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD
western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DDDD	DDDD	DDDD	DDDD	DDDD
<b>FORBS</b>							
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU
American vetch	<i>Vicia americana</i>	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP
arrowgrass	<i>Triglochin spp.</i>	TRIGL	T	T	T	T	T
asters	Asters	ASTER	UUUU	UUUU	UUUU	UUUU	UUUU
biscuitroots	<i>Lomatium spp.</i>	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD
bluebells	<i>Mertensia</i>	MERTE	DDDD	PPPP	DDDD	DDDD	DDDD
blue-eyed grass	<i>Sisyrinchium spp.</i>	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD
breadroot scurfpea	<i>Pediomelum esculentum</i>	PEES	DDDD	DDDD	DDDD	DDDD	DDDD
cattail, broad-leaf	<i>Typha latifolia</i>	TYLA	DDDD	UUUU	DDDD	UUUU	UUUU
cattail, narrow-leaf	<i>Typha angustifolia</i>	TYAN	DDDD	UUUU	DDDD	UUUU	UUUU
fringed sagewort	<i>Artemisia frigida</i>	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU
green sagewort	<i>Artemisia dracunculul</i>	ARDR4	UUUU	UUUU	UUUU	UUUU	UUUU
hawksbeard	<i>Crepis acuminata</i>	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD
horsetails	<i>Equisetum spp.</i>	EQUIS	UUUU	UUUU	UUUU	UUUU	UUUU
iris	<i>Iris spp.</i>	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU
milkvetches	<i>Astragalus</i>	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD
poison hemlock	<i>Conium maculatum</i>	COMA2	T	T	T	T	T
prairie coneflower	<i>Ratibida columnifera</i>	RACO3	DDDD	PPPP	DDDD	PPPP	PPPP
prairie thermopsis	<i>Thermopsis rhombifolia</i>	THRHA	UUUU	UUUU	UUUU	UUUU	UUUU
purple prairie clover	<i>Dalea purpurea</i>	DAPU5	PPPP	PPPP	PPPP	PPPP	PPPP
Pursh seepweed	<i>Suaeda calceoliformis</i>	SUCA2	UUUU	UUUU	UUUU	UUUU	UUUU
rosy pussytoes	<i>Antennaria rosea</i>	ANRO2	UUUU	UUUU	UUUU	UUUU	UUUU
scarlet gaura	<i>Gaura coccinea</i>	GACO5	UUUU	UUUU	UUUU	UUUU	UUUU
stemless goldenweed	<i>Haplopappus acaulis</i>	HAAC	UUUU	UUUU	UUUU	UUUU	UUUU
sulphur flower buckwheat	<i>Eriogonum umbellatum</i>	ERUM	UUUU	UUUU	UUUU	UUUU	UUUU
twogrooved milkvetch	<i>Astragalus bisulcatus</i>	ASB12	T	T	T	T	T
water hemlocks	<i>Cicuta spp.</i>	CICUT	T	T	T	T	T
western yarrow	<i>Achillea lanulosa</i>	ACHIL	UUUU	UUUU	UUUU	UUUU	UUUU
white prairie clover	<i>Dalea candida</i>	DACA7	PPPP	PPPP	PPPP	PPPP	PPPP
wild onion	<i>Allium textile</i>	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD
woodyaster	<i>Xylorhiza spp.</i>	XYLOR	T	T	T	T	T
<b>TREES, SHRUBS &amp; HALF-SHRUBS</b>							
big sagebrush	<i>Artemisia tridentata</i>	ARTR2	UUUU	DDDD	UUUU	DDDD	DDDD
birdfoot sagebrush	<i>Artemisia pedatifida</i>	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU
black greasewood	<i>Sarcobatus vermiculatus</i>	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD
fourwing saltbush	<i>Atriplex canescens</i>	ATCA2	PPPP	PPPP	PPPP	PPPP	PPPP
Gardners saltbush	<i>Atriplex gardneri</i>	ATGA	PPPP	PPPP	DDDD	PPPP	PPPP
green rabbitbrush	<i>Chrysothamnus viscidiflorous</i>	CHVI8	DDDD	DDDD	DDDD	DDDD	DDDD
junipers	<i>Juniperus scopulorum</i>	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU
plains cottonwood (sprouts)	<i>Populus deltoides</i>	PODEM	DDDD	DDDD	DDDD	DDDD	DDDD
ponderosa pine (abortion in cattle)	<i>Pinus ponderosa</i>	PIPO	UUUU	UUUU	UUUU	UUUU	UUUU
rubber rabbitbrush	<i>Ericameria nauseosa</i>	ERNA10	UUUU	DDDD	UUUU	DDDD	DDDD
silver sagebrush	<i>Artemisia cana</i>	ARCA5	DDDD	DDDD	DDDD	PPPP	PPPP
silverberry	<i>Eleagnus commutata</i>	ELCO	UUUU	UUUU	UUUU	DDDD	UUUU
skunkbush sumac	<i>Rhus trilobata</i>	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD
western snowberry	<i>Symphoricarpos occidentalis</i>	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU
wildrose	<i>Rosa woodsii var. woodsii</i>	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD
willows	<i>Salix L.</i>	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU
winterfat	<i>Krascheninnikovia lanata</i>	KRLA2	PPPP	PPPP	PPPP	PPPP	PPPP
yucca	<i>Yucca glauca</i>	YUGL	DDDD	DDDD	DDDD	DDDD	DDDD

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

## **Animal Community – Grazing Interpretations**

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

<b>Plant Community</b>	<b>Production (lb./ac)</b>	<b>Carrying Capacity* (AUM/ac)</b>
Historic Climax Plant Community	1400-2200	.5
Rhizomatous wheatgrasses/greasewood	800-1400	.4
Inland saltgrass/Cheatgrass	600-900	.15

\* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

## **Hydrology Functions**

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderate to rapid. Runoff potential for this site varies from moderate to high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts may be present. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

## **Recreational Uses**

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

## **Wood Products**

No appreciable wood products are present on the site.

## Other Products

None noted.

## Supporting Information

### Associated Sites

Overflow	058BY130WY
Lowland	058BY128WY

### Similar Sites

( ) – Saline Lowland 15-17” Northern Plains P.Z. 058BY238WY  
[Higher production]

### Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel was also used. Those involved in developing this site include: Glen Mitchell, Range Management Specialist, NRCS; Chuck Ring, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

### Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417	12	1971-1994	WY	Campbell & others
Ocular estimates	5	1990-1999	WY	Campbell & others

### State Correlation

This site has been correlated with Montana in MLRA 58B.

### Type Locality

### Field Offices

Buffalo, Douglas, Gillette, Lusk, Newcastle, Sheridan

### Relationship to Other Established Classifications

### Other References

Site Type: Rangeland  
MLRA: 58B – Northern Rolling High Plains

**Saline Lowland 10-14” P.Z.  
R058BY138WY**

## Site Description Approval

\_\_\_\_\_  
State Range Management Specialist

\_\_\_\_\_  
Date

\_\_\_\_\_  
State Range Management Specialist

\_\_\_\_\_  
Date

Your Browser must not be Java enabled, and certain elements may not appear properly, please visit our [text-only version](#).

[About NWCC](#)

[Fact Sheets](#)

[NWCC  
Brochure](#)

[Publications](#)

[Links](#)

[NRCS | USDA](#)

# National Water and Climate Center



[What's New?](#)

[Site Map](#)

[Search](#)

[E-Mail](#)

---

The National Water and Climate Center is part of the [Natural Resources Conservation Service](#), which is an agency of the [US Department of Agriculture](#). ([EEO Statement](#))

[USDA's Privacy Policy Statement](#)

---

[Text Version](#)   [Accessibility Information](#)

---