

Highlands County, Florida
 Nontechnical Soil Descriptions



Nontechnical soil descriptions describe soil properties or management considerations specific to a soil map unit or group of map units. These descriptions are written in terminology that nontechnical users of soil survey information can understand and are used to create reports. By linking the description to the soil survey map units these reports can be generated by conservation planners and other NRCS employees for distribution to land users. These descriptions are available through both TOOLKIT and NASIS.

In this subsection nontechnical descriptions are available through four categories they are Agronomic, ecological community, urban, and Water Quality. Separate map unit to description links are provided for each category.

AGRONOMIC

The following agronomic categories are available and linked through the Land Capability Unit (LCU) that are listed below.

Category

- aSOI - Soil Characteristics
- bSAC - Soil Agronomic Characteristics
- cH2O - Seasonal High Water Table
- dCUL - Cultivation Limitations
- eERO - Erosion Control
- fIRR - Irrigation Needs
- gCIT - Citrus Production
- hPAS - Pasture and Hayland
- iWMG - Water Table Management

<u>Map Symbol</u>	<u>Non hydric LCU</u>	<u>Hydric LCU</u>	<u>Drained LCU</u>	<u>Undrained LCU</u>
1	6s5			
2	7s2			
3		7w3		
4	6s8			
5	6s8			

<u>Map Symbol</u>	<u>Non hydric LCU</u>	<u>Hydric LCU</u>	<u>Drained LCU</u>	<u>Undrained LCU</u>
6	3s7			
7		7w3		
8	4w2			
9	6s5			
10	4w2			
11	6s8			
12		4w21		
13		3w21		
14	6s9			
15	3w12			
16		4w21		
17		4w22		
18			3w27	
19		7w3		
20			3w27	7w1
21			3w27	7w1
22			3w27	7w1
23			3w27	7w1
24		3w21		
25		7w3		
26			3w27	7w1
28	6s8			
29	4w3			
30	4w3			
31		7w3		
32	7e1			
33		3w22(Basinger, St.Johns) 3w21(Placid)		
34	3s6(Tavares)	7w3(Basinger)		7w1(Sanibel)
35			3w27	7w1
36	6s8			
37		7w3		
38	4w3			
39	4w2			
40	7s8			
41		5w1(Anclote) 7w3(Basinger)		
42	6s5(Astatula) 8s1(Urban Land)			

<u>Map Symbol</u>	<u>Non hydric LCU</u>	<u>Hydric LCU</u>	<u>Drained LCU</u>	<u>Undrained LCU</u>
43	8s1			
44	6s9(Satellite) 8s1(Urban Land)	7w3(Basinger)		
45	7s4(Paola)	7w3(Basinger)		
46		7w1		

Map Units without an LCU listed are either not suited to these uses or suitability is so variable that it must be determined on-site.

ECOLOGICAL COMMUNITY

The following categories are available below.

kRNG - Rangeland Suitability
 IWLD - Wildlife Suitability
 mWOD - Woodland Suitability

EC 3 (Sand Scrub) - Map Units: 1, 2, 4, 5, 11, 14, 28, 36, 45*

EC 4 (Longleaf Pine – Turkey Oak Hills) – Map Units: 6, 9, 34*

EC 6 (South Florida Flatwoods) - Map Units: 8, 10, 29, 30, 38, 39

EC 7 (North Florida Flatwoods) – Map Unit 33*

EC 10 (Cutthroat Seeps) – Map Unit: 33*

EC 13 (Cabbage Palm Hammock) - Map Units 15

EC 21 (Swamp Hardwoods) – Map Unit 41*

EC 25 (Freshwater Marshes and Ponds) - Map Units: 3, 7, 18, 19, 20, 21, 22, 23, 25, 26, 31*, 34*, 35, 37, 41*, 45*, 46

EC 26 (Slough) - Map Units: 12, 13, 16, 17, 24, 31*, 34*, 45*

* - These Map Units have more than one type of ecological community.

Map Units without an ecological community listed are not suited to these uses or suitability is so variable that it must be determined on-site.

URBAN USES

The following additional nontechnical descriptions are available for urban interpretations:

oURB - Urban Use Statement
pSEP - Septic Tank Absorption Fields
qLRS - Local Roads and Streets

01 – Map Units – 41, 46
02 - Map Units – 3, 7, 25, 31, 35, 37
03 – Map Units – 8, 10, 12, 13, 15, 16, 17, 24, 29, 30, 33, 38, 39
06 - Map Units – 14
12- Map Units – 1, 2, 6, 9, 11, 28, 36
15 - Map Units – 34, 40, 42, 43, 44, 45
20 – Map Units – 4, 5
21 – Map Units – 18, 19, 20, 21, 22, 23, 26

Map units without a link listed are either not suited to these uses or suitability is so variable that it must be determined on-site.

WATER QUALITY

The last group of nontechnical description in this subsection of this FOTG is that group dealing with water quality, specifically pesticide and nutrient management. The link between the statements and the map units is listed below.

sWQ – Water Quality Statement
tPES – Pesticide Management Statement
uNUT – Nutrient Management Statement

01 - Map Units - None
02 - Map Unit – 1, 2, 4, 5, 6, 9, 11, 14, 28, 36
03 - Map Units – 3, 7, 8, 10, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 29, 30, 31, 33, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46
04 - Map Units – None

Nontechnical Soil Descriptions

3s6 Map Unit 34(Tavares part)

"aSOI", "3s6", "This map unit consists of nearly level and gently sloping, somewhat poorly drained to moderately well drained soils on low ridges. They have moderately to rapidly permeable sandy layers to depths of more than 80 inches."

"bSAC", "3s6", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"cH2O", "3s6", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"dCUL", "3s6", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"eERO", "3s6", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "3s6", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"gCIT", "3s6", "These soils are so variable that no general interpretations for citrus can be given. An on-site investigation will be needed."

"hPAS", "3s6", "These soils are well suited to pastures. Pangola grass, hybrid bermudagrass, and bahiagrasses are well adapted. White clover and lespedezas are also well adapted. These soils produce good yields where nutrient management is practiced. Controlled grazing is needed to maintain vigorous plants for maximum yields."

"iWMG", "3s6", "This map unit is so variable that no interpretations can be given. An on-site investigation will be needed."

3s7 Map Unit 6

"aSOI", "3s7", "This map unit consists of nearly level and gently sloping, moderately well drained soils that occur on narrow to broad ridges and isolated knolls. They have very rapidly permeable sandy layers to depths of more than 80 inches."

"bSAC", "3s7", "The root zone of these soils is limited by a seasonal high water table in wet seasons and by droughtiness during periods of low rainfall. The available water capacity is low in the root zone. Natural fertility is low and crop response to fertilization is low to moderate. Rainfall is rapidly absorbed and there is little runoff. The hazard of erosion is slight."

"cH2O", "3s7", "In normal years these soils have a seasonal high water table at a depth of between 40 and 60 inches for 1 to 4 months. In other months the water table is below these depths. Rarely, only during periods of high rainfall and only for a few days, is the water table above the normal seasonal high water table depth."

"dCUL", "3s7", "These soils have severe limitations for most cultivated crops due to droughtiness and the rapid leaching of plant nutrients. These factors also limit the choice of plants and reduces potential yields of adapted crops. Crop rotations should include close growing crops on the land at least two-thirds of the time. Nutrient management maximizes yields. Soil improving cover crops and all crop residues should be left on the ground."

"fIRR", "3s7", "Irrigation of high value crops is usually feasible where irrigation water is readily available."

"hPAS", "3s7", "These soils are moderately well suited to pastures. Hybrid bermudagrass and bahiagrasses are adapted. White clover and lespedezas are also adapted. These soils produce good yields where nutrient management is practiced. Controlled grazing is needed to maintain vigorous plants for maximum yields."

"iWMG", "3s7", "Tile, or other types of drains, are needed for some crops such as tobacco that are damaged by high water table during the growing season. Tiles can also be used as a source for subirrigation during periods of low rainfall."

3w12 Map Unit 15

"aSOI", "3w12", "This map unit consists of nearly level, poorly drained soils on flatwoods, hammocks, and other flat areas. They have sandy surface and subsurface layers less than 20 inches thick over moderately to moderately rapidly permeable loamy layers."

"bSAC", "3w12", "The root zone is limited by a seasonal high water table that comes to near the surface in wet seasons. The available water capacity averages moderate in the root zone. Natural fertility is low but crop response to fertilization is good. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "3w12", "In normal years these soils have a seasonal high water table at a depth of between 6 and 18 inches for 1 to 4 months. In other months the water table is below these depths. Rarely, only during periods of high rainfall and only for a few days, is the water table above the normal seasonal high water table depth."

"dCUL", "3w12", "These soils have severe limitations for cultivated crops because of wetness. With a total water management system these soils are well suited to a variety of fruit and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. The cover crops and all other crop residue should be returned to the soil. Maximum yields require good soil tilth and nutrient management."

"eERO", "3w12", "Crops produced on these soils do not normally need special erosion control practices."

"fIRR", "3w12", "Highest yields require irrigation during periods of low rainfall. Water can be supplied through subirrigation with a water table management system or by sprinklers."

"gCIT", "3w12", "With proper water table management these soils are suited to citrus crops where they occur in places relatively free from damaging cold in winter. Good management includes adequate water control to maintain the water table at least three feet below the surface. The trees should be planted on beds. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion."

"hPAS", "3w12", "These soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "3w12", "A total water table management system should remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

3w21 Hydric, portions of Map Units 13, 24, 33(Placid part)

"aSOI", "3w21", "This map unit consists of nearly level, poorly drained soils on low flatwoods, low hammocks, and sloughs. They have sandy surface and subsurface layers 20 to 40 inches thick over moderately to moderately rapidly permeable loamy layers."

"bSAC", "3w21", "The root zone is limited by a seasonal high water table that is at or slightly above the surface in wet seasons. The available water capacity averages moderate in the root zone. Natural fertility is low but crop response to fertilization is good. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "3w21", "In normal years these soils have a seasonal high water table at a depth of 6 inches or less for 2 to 6 months. In other months the water table is usually below this depths. During periods of high rainfall the water table may be above the surface for periods of brief duration."

"dCUL", "3w21", "Cultivation of these hydric soils is not recommended. If cultivated, these soils have severe limitations because of wetness. With a total water management system these soils are suited to a variety of fruit and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. The cover crops and all other crop residue should be returned to the soil. Maximum yields require good soil tilth and nutrient management."

"eERO", "3w21", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "3w21", "If cultivated, Highest yields require irrigation during periods of low rainfall either subirrigated through a water table management system or by sprinklers."

"gCIT", "3w21", "With proper water table management these soils are suited to citrus crops where they occur in places relatively free from damaging cold in winter. Good management includes adequate water control to maintain the water table at least three feet below the surface. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion."

"hPAS", "3w21", "These hydric soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "3w21", "If cropped, these hydric soils need a total water table management system to remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

3w27 Drained portions of Map Units 18, 20, 21, 22, 23, 26, 35

"aSOI", "3w27", "This map unit consists of nearly level, very poorly drained organic soils on broad flats and depressions. These hydric soils have been drained and protected from flooding and ponding by means of pumping stations, canals, and dikes for crop production and other uses."

"bSAC", "3w27", "The root zone is limited by water that is above the surface in wet seasons. The available water capacity averages high in the root zone. Natural fertility is high. The internal drainage rate is very slow in the natural condition and seepage water seeps from the soil in wet seasons."

"dCUL", "3w27", "If water control measures are established, these soil would be moderately well to well suited to cultivated crops. Due to the difficulty of installing these measures and the lack of outlets in most areas, they have seldom, if ever, been used for crops."

"eERO", "3w27", "Due to the lack of these soils being cultivated, erosion control is not a management concern."

"fIRR", "3w27", "Due to the lack of cultivation, irrigation is not a normal practice on these soils."

"gCIT", "3w27", "In normal years these soils have a seasonal high water table within 6 inches of the surface for 2 to 6 months of most years. During other months the water table is deeper. These soils are also subject to frequent ponding and/or flooding. Only rarely is the water table below the surface for an extended period."

"hPAS", "3w27", "If water control measures are established, these soil would be moderately well to well suited to improved pastures. Due to the difficulty of installing these measures and the lack of outlets in most areas, they have seldom, if ever, been used for pasture."

"iWMG", "3w27", "Water table management is not a normal practice on these soils because of the lack of cultivation."

4w2 Non-hydric portions of Map Units 8, 10, 39

"aSOI", "4w2", "This map unit consists of nearly level, poorly drained soils on flatwoods, hammocks, and other flat areas. They have sandy layers more than 72 inches thick."

"bSAC", "4w2", "The root zone is limited by a seasonal high water table that is at or slightly above the surface in wet seasons. The available water capacity averages moderate in the root zone. Natural fertility is low but crop response to fertilization is good. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "4w2", "In normal years these soils have a seasonal high water table at a depth of 6 inches or less for 2 to 6 months. In other months the water table is usually below this depths. During periods of high rainfall the water table may be above the surface for periods of brief duration."

"dCUL", "4w2", "Cultivation of these hydric soils is not recommended. If cultivated, these soils have severe limitations because of wetness. With a total water management system these soils are suited to a variety of fruit and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. The cover crops and all other crop residue should be returned to the soil. Maximum yields require good soil tilth and nutrient management."

"eERO", "4w2", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "4w2", "If cultivated, Highest yields require irrigation during periods of low rainfall either subirrigated through a water table management system or by sprinklers."

"gCIT", "4w2", "With proper water table management these soils are suited to citrus crops where they occur in places relatively free from damaging cold in winter. Good management includes adequate water control to maintain the water table at least three feet below the surface. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion."

"hPAS", "4w2", "These hydric soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "4w2", "If cropped, these hydric soils need a total water table management system to remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

4w3 Non-hydric portions of Map Units 29, 30, 38

"aSOI", "4w3", "This map unit consists of nearly level, poorly drained soils on flatwoods, hammocks, and other flat areas. They are dominately sandy with a moderately slowly subsoil layer within 60 inches."

"bSAC", "4w3", "The root zone is limited by a seasonal high water table that comes to near the surface in wet seasons and by droughtiness during periods of low rainfall. The available water capacity averages low to very low in the root zone. Natural fertility is low and crop response to fertilization is moderate. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "4w3", "In normal years these soils have a seasonal high water table at a depth of between 6 and 18 inches for 1 to 4 months. In other months the water table is below these depths. Rarely, only during periods of high rainfall and only for a few days, is the water table above the normal seasonal high water table depth."

"dCUL", "4w3", "These soils have severe limitations for cultivated crops because of wetness in wet seasons and droughtiness during periods of low rainfall. With a total water management system these soils are well suited to a variety of flower and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. All crop residue should be returned to the soil. Maximum yields require nutrient management."

"eERO", "4w3", "Erosion control is not a management concern on these soils."

"fIRR", "4w3", "Highest yields require irrigation during periods of low rainfall either subirrigated through a water table management system or by sprinklers."

"gCIT", "4w3", "With proper water table management these soils are suited to citrus crops. Good management includes adequate water control to maintain the water table at least three feet below the surface. The trees should be planted on beds. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion. Irrigation is required for proper yields."

"hPAS", "4w3", "These soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "4w3" "A total water table management system should remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

4w21 Hydric portions of Map Unit 12, 16

"aSOI", "4w21", "This map unit consists of nearly level, poorly drained soils on low flatwoods, low hammocks, and sloughs. They have sandy layers more than 72 inches thick and a spodic horizon within 30 inches."

"bSAC", "4w21", "The root zone is limited by a seasonal high water table that is at or slightly above the surface in wet seasons. The available water capacity averages moderate in the root zone. Natural fertility is low but crop response to fertilization is good. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "4w21", "In normal years these soils have a seasonal high water table at a depth of 6 inches or less for 2 to 6 months. In other months the water table is usually below this depths. During periods of high rainfall the water table may be above the surface for periods of brief duration."

"dCUL", "4w21", "Cultivation of these hydric soils is not recommended. If cultivated, these soils have severe limitations because of wetness. With a total water management system these soils are suited to a variety of fruit and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. The cover crops and all other crop residue should be returned to the soil. Maximum yields require good soil tilth and nutrient management."

"eERO", "4w21", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "4w21", "Highest yields require irrigation during periods of low rainfall either subirrigated through a water table management system or by sprinklers."

"gCIT", "4w21", "With proper water table management these soils are suited to citrus crops where they occur in places relatively free from damaging cold in winter. Good management includes adequate water control to maintain the water table at least three feet below the surface. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion."

"hPAS", "4w21", "These hydric soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "4w21", "If cropped, these hydric soils need a total water table management system to remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

4w22 Hydric portions of Map Unit 17

"aSOI", "4w22", "This map unit consists of nearly level, poorly drained soils on low flatwoods, low hammocks, and sloughs. They are dominately sandy with a moderately slowly subsoil layer within 60 inches."

"bSAC", "4w22", "The root zone is limited by a seasonal high water table that is at or near the surface in wet seasons and by droughtiness during periods of low rainfall. The available water capacity averages low to very low in the root zone. Natural fertility is low and crop response to fertilization is moderate. Internal drainage is slow but response to artificial drainage is moderate to rapid. The hazard of erosion is slight."

"cH2O", "4w22", "In normal years these hydric soils have a seasonal high water table at a depth of less than 6 inches for 2 to 6 months. In other months the water table is usually below these depths. During periods of high rainfall the water table may be above the surface for periods of brief duration."

"dCUL", "4w22", "Cultivation of these hydric soils is not recommended. If cultivated, severe limitations due to wetness in wet seasons and droughtiness during periods of low rainfall exist. With a total water management system these soils are suited to a variety of flower and vegetable crops. Management should include crop rotations that keep the soil in close growing cover crops at least two-thirds of the time. All crop residue should be returned to the soil. Maximum yields require nutrient management."

"eERO", "4w22", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "4w22", "Highest yields require irrigation during periods of low rainfall either subirrigated through a water table management system or by sprinklers."

"gCIT", "4w22", "With proper water table management these hydric soils are suited to citrus crops. Good management includes adequate water control to maintain the water table at least three feet below the surface. The trees should be planted on beds. Nutrient management is a preferred practice. Close growing vegetation between the trees is needed to protect the soil from erosion. Irrigation is required for proper yields."

"hPAS", "4w22", "These hydric soils are well suited to pastures and hay crops. Improved grasses such as pangola grass and bahiagrasses are well adapted. Several varieties of clovers are also well adapted where properly managed. High yields require nutrient management, water table management, and controlled grazing to prevent overgrazing."

"iWMG", "4w22", "A total water table management system should remove excess water rapidly and provide a means of applying subirrigation. Tile drains, open ditches, and/or tail-race recovery systems may be needed to maintain the preferred water table depths of within 18 inches for vegetables and below four feet for citrus. To obtain adequate drainage, the spacing of tile drains is important. Tile drains may be used for subirrigation during periods of low rainfall."

5w1 Map Unit 41(Anclote part)

"aSOI", "5w1", "This map unit consists of nearly level, poorly drained and very poorly drained soils on flood plains. These soils are subject to common flooding during the growing season."

"bSAC", "5w1", "Wetness and flooding severely limits the use of the root zone of these soils for agronomic crops."

"dCUL", "5w1", "These hydric soils are not suited to cultivated crops without an extensive water table management system."

"eERO", "5w1", "Due to the lack of these soils being cultivated, erosion control is not a management concern."

"fIRR", "5w1", "If cultivated, highest yields require irrigation either subirrigated through the extensive water table management system or by sprinklers"

"hPAS", "5w1", "These hydric soils are not suited to pasture or hay crops without an extensive water table management system."

"iWMG", "5w1", "If these hydric soils are cultivated, an extensive water table management system is needed for crop and pasture production on these soils. It should remove excess water rapidly and provide a means of applying subirrigation. Dikes and a pumping systems are needed for flood control and tile drains and open ditches are needed to maintain the preferred water table depth. Rarely are drainage and flood protection economically feasible and environmentally "

6s5 Map Units 1, 9, 42(Astatula part)

"aSOI", "6s5", "This nearly level to moderately sloping soil is on high sandy ridges in the ridge part of the county. Slopes range from 0 to 8 percent. The available water capacity of this soil is very low. The permeability is very rapid. Depth to the water table is more than 80 inches."

"bSAC", "6s5", "The soils have a loose, well aerated root zone to depths of more than 80 inches. The available water capacity averages very low in the root zone. Natural fertility is very low and nutrients are rapidly leached from the soil. Rainfall is rapidly absorbed on protected areas, and there is little runoff. Erosion is not a serious hazard."

"cH2O", "6s5", "In normal years these soils do not have a seasonal high water table within a depth of 72 inches."

"dCUL", "6s5", "Due to the very low natural fertility, droughtiness, and the rapid leaching of plant nutrients, these soils are not suited to cultivated field crops."

"eERO", "6s5", "If these soils are cultivated, erosion control measures that would adequately protect the soil and water resource base are difficult to install and/or maintain."

"fIRR", "6s5", "Irrigation of high value crops is usually feasible where irrigation water is readily available. The rate of application should be low enough to prevent runoff and erosion. A well designed irrigation system to maintain optimum moisture conditions is needed to assure acceptable citrus yields."

"gCIT", "6s5", "These soils are fairly suited to citrus trees even where they are in places relatively free from freezing temperatures. A good ground cover of close growing plants is needed between the trees to protect the soil from blowing and washing. Poor to fair yields of oranges and grapefruit are usually obtained without irrigation."

"hPAS", "6s5", "These soils have only fair suitability for pastures. Grasses such as hybrid bermudagrass and bahiagrass make only fair growth where an intensive nutrient management system is maintained. Clovers are not adapted."

"iWMG", "6s5", "Water table management is not normally practiced on these soils."

6s8 Map Units 4, 5, 11, 28, 36

"aSOI", "6s8", "This map unit consists of nearly level, somewhat poorly and moderately well drained soils on low ridges of the flatwoods. They have sandy layers to more than 72 inches deep. A layer 20 to 60 inches below the surface is weakly cemented with dark colored organic material."

"bSAC", "6s8", "The root zone is limited by a water table during wet seasons and by droughtiness during periods of low rainfall. The available water capacity is very low in the root zone. Natural fertility is very low and crop response to nutrient management is only fair. The internal drainage rate is slow under natural conditions but response to artificial drainage is rapid."

"cH2O", "6s8", "In normal years these soils have a seasonal high water table at a depth of 18 and 40 inches for 1 to 4 months. In other months the water table is usually below this depth. Only rarely, during periods of high rainfall, is the water table above 18 inches."

"dCUL", "6s8", "Due to the very low natural fertility, wetness in wet seasons, droughtiness during periods of low rainfall, and the rapid leaching of plant nutrients, these soils are not suited to cultivated field crops."

"eERO", "6s8", "Crops produced on these soils do not normally need special erosion control practices."

"fIRR", "6s8", "Irrigation of high value crops is usually feasible where irrigation water is readily available. The rate of water application should be low enough to prevent runoff and erosion. A well designed irrigation system to maintain optimum moisture conditions is needed to assure acceptable citrus yields."

"gCIT", "6s8", "These soils are only fairly to poorly suited for citrus trees even where they are in places relatively free from freezing temperatures. A good ground cover of close growing plants is needed between the trees to protect the soil from blowing and washing. Poor to fair yields of oranges and grapefruit are usually obtained without irrigation."

"hPAS", "6s8", "These soils have only fair suitability for pastures. Grasses such as pangola grass and bahiagrass make only fair growth where an intensive nutrient management system is maintained. Clovers are not adapted."

"iWMG", "6s8", "Water table management is not normally practiced on these soils."

6s9 Map Unit 14, 44(Satellite part)

"aSOI", "6s9", "This map unit consists of nearly level, somewhat poorly and moderately well drained soils on low ridges of the flatwoods. They have sandy layers more than 80 inches deep."

"bSAC", "6s9", "The root zone is limited by a water table during wet seasons and by droughtiness during periods of low rainfall. The available water capacity is very low in the root zone. Natural fertility is very low and crop response to nutrient management is only fair. The internal drainage rate is slow under natural conditions but response to artificial drainage is rapid."

"cH2O", "6s9", "In normal years these soils have a seasonal high water table at a depth of 18 and 40 inches for 1 to 4 months. In other months the water table is usually below this depth. Only rarely, during periods of high rainfall, is the water table above 18 inches."

"dCUL", "6s9", "Due to the very low natural fertility, wetness in wet seasons, droughtiness during periods of low rainfall, and the rapid leaching of plant nutrients, these soils are not suited to cultivated field crops."

"eERO", "6s9", "If these soils are cultivated, erosion control measures are not normally needed."

"fIRR", "6s9", "Irrigation of high value crops is usually feasible where irrigation water is readily available. The rate of water application should be low enough to prevent runoff and erosion. A well designed irrigation system to maintain optimum moisture conditions is needed to assure acceptable citrus yields"

"gCIT", "6s9", "These soils are only fairly to poorly suited to citrus trees even where they are in places relatively free from freezing temperatures. A good ground cover of close growing plants is needed between the trees to protect the soil from blowing and washing. Poor to fair yields of oranges and grapefruit are usually obtained without irrigation."

"hPAS", "6s9", "These soils have only fair suitability for pastures. Grasses such as pangola grass and bahiagrass make only fair growth where an intensive nutrient management system is maintained. Clovers are not adapted."

"iWMG", "6s9", "Water table management is not normally practiced on these soils."

7s4 Map Unit 45(Paola)

"aSOI", "7s4", "This map unit consists of excessively drained, strongly sloping to steep soils on side slopes of upland ridges. They have sandy layers to depths of more than 80 inches."

"bSAC", "7s4", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"cH2O", "7s4", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"dCUL", "7s4", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"eERO", "7s4", "Crops produced on these hydric soils do not normally need special erosion control practices."

"fIRR", "7s4", "These soils are so variable that no interpretations can be given. An on-site investigation will be needed."

"gCIT", "7s4", "These soils are so variable that no general interpretations for citrus can be given. An on-site investigation will be needed."

"hPAS", "7s4", "These soils are well suited to pastures. Pangola grass, hybrid bermudagrass, and bahiagrasses are well adapted. White clover and lespedezas are also well adapted. These soils produce good yields where nutrient management is practiced. Controlled grazing is needed to maintain vigorous plants for maximum yields."

"iWMG", "7s4", "This map unit is so variable that no interpretations can be given. An on-site investigation will be needed."

7s8 Map Unit 40

"aSOI", "7s8", "This map unit consists of nearly level to steep soils on disturbed areas. These soils are variable; however, they are deep and predominately stratified sandy and loamy material."

"bSAC", "7s8", "The available water capacity is low. Natural fertility is low and response to fertilization is low. Rainfall is rapidly absorbed but moves rapidly through the soil and very little is retained. There is severe hazard of gully erosion."

"cH2O", "7s8", "In normal years these soils do not have a seasonal high water table within 72 inches of the surface.."

"dCUL", "7s8", "These soils are too steep and erodible to be suited to cultivated crops."

"eERO", "7s8", "If these soils are cultivated, erosion control measures that would adequately protect the soil and water resource base are difficult to install and/or maintain."

"fIRR", "7s8", "Due to the lack of cultivation, irrigation is not a normal practice on these soils."

"hPAS", "7s8", "These soils are poorly suited to pastures due to steepness of the soil and the hazard of erosion."

"iWMG", "7s8", "Water table management is not a normal practice on these soils because of the lack of cultivation and an available water source."

7w1 Map Unit 46

Undrained portions of Map Units 20, 21, 22, 23, 26, 34(Sanibel), 35

"aSOI", "7w1", "This map unit consists of nearly level, very poorly drained organic soils in depressional areas and floodplains. They have thick layers of partially decomposed remains of aquatic plants."

"bSAC", "7w1", "The root zone is limited by water that is above the surface in wet seasons. The available water capacity averages high in the root zone. Natural fertility is high. The internal drainage rate is very slow in the natural condition and seepage water seeps from the soil in wet seasons."

"dCUL", "7w1", "If water control measures are established, these soil would be moderately well to well suited to cultivated crops. Due to the difficulty of installing these measures and the lack of outlets in most areas, they have seldom, if ever, been used for crops."

"eERO", "7w1", "Due to the lack of these soils being cultivated, erosion control is not a management concern."

"fIRR", "7w1", "Due to the lack of cultivation, irrigation is not a normal practice on these soils."

"gCIT", "7w1", "In normal years these soils have a seasonal high water table within 6 inches of the surface for 2 to 6 months of most years. During other months the water table is deeper. These soils are also subject to frequent ponding and/or flooding. Only rarely is the water table below the surface for an extended period."

"hPAS", "7w1", "If water control measures are established, these soil would be moderately well to well suited to improved pastures. Due to the difficulty of installing these measures and the lack of outlets in most areas, they have seldom, if ever, been used for pasture."

"iWMG", "7w1", "Water table management is not a normal practice on these soils because of the lack of cultivation."

7w3 Map Units 3, 7, 19, 25, 31, 34(Basinger), 37, 41(Basinger), 44(Basinger), 45(Basinger)

"aSOI", "7w3", "This map unit consists of nearly level, very poorly drained soils on depressions. They have sandy or loamy surface layers and sandy, loamy or clayey subsoil layers."

"bSAC", "7w3", "The root zone is limited by water that is above the surface in wet seasons. The available water capacity averages moderate in the root zone. Natural fertility is moderate. The internal drainage rate is very slow in the natural condition and seepage water seeps from the soil in wet seasons."

"cH2O", "7w3", "In normal years these soils have a seasonal high water table within 6 inches of the surface for 2 to 6 months of most years. During other months the water table is deeper. These soils are also subject to frequent ponding. Only rarely is the water table below the surface for an extended period."

"dCUL", "7w3", "Due to extreme wetness, these soils are not suited to cultivated crops."

"eERO", "7w3", "Due to the lack of these soils being cultivated, erosion control is not a management concern."

"fIRR", "7w3", "Due to the lack of cultivation, irrigation is not a normal practice on these soils."

"hPAS", "7w3", "If water control measures are established, these soil would be moderately well suited to improved pastures. Due to the difficulty of installing these measures and the lack of outlets in most areas, they have seldom, if ever, been used for pasture."

"iWMG", "7w3", "Water table management is not a normal practice on these soils because of the lack of cultivation."

8s1 Map Units 42(Urban Land), 43, 44(Urban Land)

"aSOI", "8s1", "This map unit consists of miscellaneous areas where no soil exists and has no value for agricultural uses."

"bSAC", "8s1", "Due to an impervious surface these areas are not vegetated."

"cH2O", "8s1", " These soils have a highly variable water table."

"dCUL", "8s1", "Due to the impervious surface, these soils are not suited to cultivated crops."

"eERO", "8s1", "Due to the lack of these soils being cultivated, erosion control is not a management concern."

"fIRR", "8s1", "Due to the lack of cultivation, irrigation is not a normal practice on these soils."

"hPAS", "8s1", "Due to the impervious surface, actions, these soils are not suited to hay and pasture."

"iWMG", "8s1", "Water table management is not a normal practice on these soils because of the lack of cultivation."

ECOLOGICAL COMMUNITIES

kRNG - Rangeland

IWLD - Wildlife

mWOD - Woodland

Sand Scrub - Map Units 1, 2, 4, 5, 11, 14, 28, 36, 45

"kRNG", "03", "This Sand Scrub range site supports a dense stand of trees and shrubs and has limited potential for producing native forage. Sites in excellent condition produce 1500 to 3500 pounds per acre annually. Fifteen to 40 acres or more are usually needed per animal unit. Little forage will be available if the tree canopy cover exceeds 60%. Forage is usually 75% grasses and grass-like plants, 15% trees and shrubs, and 10% forbs."

"IWLD", "03", "This Sand Scrub site is suited to deer and turkey, especially as escape cover. Many birds inhabit the area including warblers, towhees, flycatchers, scrub jays, and quail. Native legumes furnish food (seeds) for the birds. Fruits of palmetto, gopher apple, and various species of oak are also a good food source. Timber harvest and other disturbances increase wildlife food by increasing the amount and types of Herbaceous plants and by sprout production."

"mWOD", "03", "This Sand Scrub site has a low potential for commercial production of wood and timber. The soils create severe equipment limitations and moderate seedling mortality problems. Sand pine is a commercial species suited to planting. It has a potential annual growth of approximately 0.4 to 0.5 cords per acre."

Longleaf Pine - Turkey Oak Hills – Map Units 6, 9, 34*

"kRNG", "04", "This Longleaf Pine - Turkey Oak Hills range site provides poor quality and low quantity forage and has limited potential for producing native forage. Sites in excellent condition produce 2000 to 4000 pounds per acre annually. Ten to 35 acres or more are usually needed per animal unit. Little forage will be available if the tree canopy cover exceeds 60%. Forage is usually 75% grasses and grass-like plants, 15% trees and shrubs, and 10% forbs."

"IWLD", "04", "This Longleaf Pine - Turkey Oak Hills site is suited to deer and turkey, especially as escape cover. Many birds inhabit the area including warblers, towhees, flycatchers, scrub jays, and quail. Native legumes furnish food (seeds) for the birds. Fruits of palmetto, gopher apple, and various species of oak are also a good food source. Timber harvest and other disturbances increase wildlife food by increasing the amount and types of herbaceous plants and by sprout production."

"mWOD", "04", "This Longleaf Pine - Turkey Oak Hills site has a moderately high potential for commercial production of wood and timber. The soils create moderate equipment limitations and moderate seedling mortality problems. Commercial species suited to planting and their potential annual growth in cords are as follows; Sand pine, 1.2 to 1.0. Slash pine, 1.2 to 1.0. Loblolly pine, 1.0 to 0.8. Longleaf pine, 0.6 to 0.5."

South Florida Flatwoods - Map Units 8, 10, 29, 30, 38, 39

"kRNG", "06", "This South Florida Flatwoods range site has the potential for producing significant amounts of high quality forage from creeping bluestem, chalky bluestem, and indiangrass. Sites in excellent condition produce 3000 to 6000 pounds per acre annually. Three to 16 acres or more are usually needed per animal unit. Little forage will be available if the tree canopy cover exceeds 60%. Forage is usually 75% grasses and grass-like plants, 15% trees and shrubs, and 10% herbaceous plants."

"IWLD", "06", "This South Florida Flatwoods site is well suited to deer, turkey, and quail. It is fairly suited to squirrels and well suited to many songbirds. Palmetto fruit, pine mast, oak acorns, legume seed, and grasses are good sources of wildlife food. Mature hardwoods and snags provide good nesting sites for birds. This site is also well suited to bobcat, raccoons, opossums, and skunks. It is poorly suited to dove."

"mWOD", "06", "This South Florida Flatwoods site has a moderate potential for commercial production of wood and timber. The soils create moderate equipment limitations and moderate seedling mortality rates. Commercial species suited to planting and their potential annual growth in cords are as follows: Slash pine, 0.9 to 0.7. Longleaf pine, 0.5 to 0.4."

North Florida Flatwoods – Map Unit 33*

"kRNG", "07", "This North Florida Flatwoods range site has the potential for producing significant amounts of high quality forage from chalky bluestem, indiangrass, and panicums. Sites in excellent condition produce 3000 to 5500 pounds per acre annually. Five to 15 acres or more are usually needed per animal unit. Little forage will be available if the tree canopy cover exceeds 60%. Forage is usually 75% grasses and grass-like plants, 15% trees and shrubs, and 10% herbaceous plants."

"IWLD", "07", "This North Florida Flatwoods site is well suited to deer, turkey, and quail. It is fairly suited to squirrels and well suited to many songbirds. Palmetto fruit, pine mast, oak acorns, legume seed, and grasses are good sources of wildlife food. This site is also well suited to bobcat, raccoons, opossums, and skunks. It is poorly suited to dove."

"mWOD", "07", "This North Florida Flatwoods site has a moderate potential for commercial production of wood and timber. The soils create moderate equipment limitations and moderate seedling mortality rates. Commercial species suited to planting and their potential annual growth in cords are as follows: Slash pine, 1.0 to 0.8. Longleaf pine, 0.6 to 0.4."

Cutthroat Seep – Map Unit 33*

"kRNG", "10", "This Cutthroat Seep range site has the potential for producing significant amounts of high quality forage from cutthroat grass, bluestems, and panicums. Sites in excellent condition produce 3000 to 5500 pounds per acre annually. Four to 16 acres or more are usually needed per animal unit. Little forage will be available if the tree canopy cover exceeds 60%. Forage is usually 75% grasses and grass-like plants, 10% trees and shrubs, and 15% herbaceous plants."

"IWLD", "10", "This Cutthroat Seep site is well suited to deer, turkey, and many songbirds. It is fairly suited to quail and moderately suited to skunks, opossums, and raccoons. Palmetto fruit, pine mast, legume seed, and grasses are good sources of wildlife food. This site is also valuable as a habitat for reptiles such as ratsnakes and rattlesnakes. It is poorly suited to squirrels and doves."

"mWOD", "10", "This Cutthroat Seep site has a moderately low potential for commercial production of wood and timber. The soils create severe equipment limitations and severe seedling mortality rates. There are no recommended commercial species suited for planting. Potential annual growth of existing slash pine and pond pine is about 0.4 cords per acre."

Cabbage Palm Hammock - Map Units 15

"kRNG", "13", "This Cabbage Palm Hammock range site has a low potential for producing forage due to a dense canopy of and cabbage palms. It provides livestock protection in cold and hot weather. Sites in excellent condition produce 2000 to 4000 pounds per acre each year. Ten to 30 acres per animal unit are needed. If the tree canopy cover exceeds 60%, little forage will be available. The annual forage produced is 55% grasses and grass-like plants, 25% trees and shrubs, and 20% herbaceous plants."

"IWLD", "13", "This Cabbage Palm Hammock site is well suited to deer, turkey, squirrel, black bear, feral and wild hogs, woodpeckers, and owls. Palm and palmetto fruit, pine mast, oak acorns, legume seed, and grasses are good sources of wildlife food. Habitat is poor for quail and dove and fair for most songbirds and squirrels."

"mWOD", "13", "This Cabbage Palm Hammock site has a moderately high to high potential for commercial production of wood and timber. The soils create moderate equipment limitations and moderate seedling mortality rates. Commercial species suited to planting and their potential annual growth in cords are as follows: Slash pine, 1.5 to 1.3. Loblolly pine, 1.2 to 1.0. Sweetgum, 1.5 to 1.3. Sycamore is also suitable for planting."

Swamp Hardwoods – Map Unit 41*

"kRNG", "21", "This Swamp Hardwoods site has little or no range value."

"IWLD", "21", "This Swamp Hardwoods site is well suited to a wide variety of waterfowl, reptiles, amphibians, and mammals. These species must withstand periodic flooding. Inhabitants include squirrels, deer, mink, otter, raccoons, chickadees, titmice, flycatchers, owls, ducks, woodpeckers, wrens, and many other birds. Hardwood vegetation provides good cover and wildlife food for these and many other wildlife species."

"mWOD", "21", "This Swamp Hardwoods site is generally not used for the commercial production of wood and timber; however, it does potential.. The soils create severe equipment limitations and severe seedling mortality rates. Commercial species suited to planting in areas with adequate surface drainage and their potential annual growth in cords are as follows: Slash pine, 1.5 to 1.3. Loblolly pine, 1.2 to 1.0. Cottonwood, 0.8 to 0.6. Sweetgum and sycamore are additional species suitable to plant."

Freshwater Marshes and Ponds - Map Units 3, 7, 18, 19, 20, 21, 22, 23, 25, 26, 31*, 34*, 35, 37, 41*, 45*, 46

"kRNG", "25", "This Freshwater Marsh and Ponds range site has the potential for producing significant amounts of high quality forage from a variety of high quality forage plants. Sites in excellent condition produce 5000 to 10000 pounds per acre annually. Three to 13 acres or more are usually needed per animal unit. Forage is usually 80% grasses and grass-like plants, 5% trees and shrubs, and 15% herbaceous plants."

"IWLD", "25", "This Freshwater Marsh and Ponds site is well suited to a wide variety of wetland wildlife species including waterfowl, reptiles, amphibians, and mammals. These species must withstand ponding of long or very long duration. Inhabitants include mink, otter, raccoons, herons, bitterns, ibis, cranes, snipe, ducks, kites, killdeer, caracara, and hawks. This community also serves as a water source for species from surrounding communities."

"mWOD", "25", "This Freshwater Marsh and Ponds site is seldom used for the commercial production of wood and timber. The soils create very severe limitations that are difficult to overcome."

Slough - Map Units 12, 13, 16, 17, 24, 31*, 34*, 45*

"kRNG", "26", "This Slough range site has the potential for producing significant amounts of high quality forage from a variety of high quality forage plants such as maidencanes, bluestems, and panicums. Sites in excellent condition produce 3000 to 6000 pounds per acre annually. Four to 16 acres or more are usually needed per animal unit. Forage is usually 85% grasses and grass-like plants and 15% herbaceous plants."

"IWLD", "26", "This Slough site is well suited to snakes, frogs, salamanders, raccoons, and wading birds. The grass dominated vegetation is a highly valued food source for quail and deer; however, it provides poor cover for these and most other wildlife species except at its confluence with other communities."

"mWOD", "26", "This Slough site is seldom used for the commercial production of wood and timber. The soils create very severe limitations that are difficult to overcome."

URBAN USES

oURB - Urban Use Statement

pSEP - Septic Tank Absorption Fields

qLRS - Local Roads and Streets

Map Units 41, 46

"oURB", "01", "This soil is generally unsuited to most urban uses because of flooding. Dwellings and small buildings can be constructed on pilings, however, access may be limited during flood events and structural integrity of the building may be threatened by currents and floating debris. Landscaping considerations should include use of species that are adapted to withstanding flood water."

"pSEP", "01", "This soil has very severe limitations for septic tank absorption fields. Flooding interferes with absorption of effluent from septic tanks and poses risks of contamination to adjacent surface waters."

"qLRS", "01", "This soil has severe limitations for local roads and streets. Road surfaces and bases may be eroded by floodwaters and travel is dangerous or impractical during flood events."

Map Units 3, 7, 25, 31, 35, 37

"oURB", "02", "This soil is generally unsuited to most urban uses because of ponding and low bearing strength of the soil. Dwellings and small buildings can be constructed on pilings driven to suitable depths, however, access may be limited during periods when water tables are highest. Drainage may be impractical in many areas because of a lack of suitable outlets. Landscaping considerations should include use of species that are adapted to ponded water and organic soils."

"pSEP", "02", "This soil has severe limitations for septic tank absorption fields. Ponded water tables and organic soil materials interfere with the absorption of effluent from septic tanks and pose risks of contamination to adjacent surface waters."

"qLRS", "02", "This soil has severe limitations for local roads and streets. Road and street surfaces may subside, crack or ripple if sufficient fill is not used as a base. When possible, organic soil material should be removed and filled with suitable soil material to prevent subsidence and damage to road surfaces."

Map Units 8, 10, 12, 13, 15, 16, 17, 24, 29, 30, 33, 38, 39

"oURB", "03", "This soil is poorly suited to most urban uses because of a seasonal high water table at or near the soil surface. Housing pads, driveways, and other home site areas can be elevated using suitable fill. Area drainage can be installed to lower the water table if suitable outlets are available. Fill may also be used to elevate sites for small commercial buildings. Landscaping considerations should include use of species that are adapted to wetness."

"pSEP", "03", "This soil has severe limitations for septic tank absorption fields. High water tables interfere with the absorption of effluent from septic tanks and pose risks of contamination to adjacent surface waters. Septic tank absorption fields can be mounded to maintain the system above the seasonal high water table."

"qLRS", "03", "This soil has severe limitations for local roads and streets. For any construction, care should be taken not to impede natural drainage or impound water on the site and adjacent areas. Well designed culvert placement beneath any fill and use of existing water conveying landscapes can help minimize disturbance to natural drainage."

Map Unit 14

"oURB", "06", "Suitability is poor for most urban land uses because of a seasonal high water table within 40 inches of the soil surface. House and small building pads can be elevated using suitable fill. The fill can be placed with a slight grade to allow water to drain away from the house or building. Irrigation can be helpful in establishing plants and for maintenance during dry periods. Landscaping considerations should include use of species that are adapted to wetness."

"pSEP", "06", "This soil has severe limitations for septic tank absorption fields. High water tables interfere with the absorption of effluent from septic tanks. This poses risks of contamination to adjacent surface waters and system failure. Septic tank absorption fields can be mounded to maintain the system above the seasonal high water table. Absorption field laterals should be installed on a slight downslope gradient. Absorption fields should be placed downslope from dwellings."

"qLRS","06","This soil has severe limitations for local roads and streets. They can be elevated using suitable fill. The fill can be placed with a slight grade to allow water to drain away from the house or building. An engineer or soil scientist should be consulted to determine the shrink-swell potential of near surface soil material. Additional design precautions can be planned if shrink-swell is determined to be a concern."

Map Units 1, 2, 6, 9, 11, 28, 36

"oURB","12","This soil is well suited to most urban uses. There are no significant limitations. Landscaping considerations should include use of species that are adapted to droughty soil conditions. Irrigation can be helpful in establishing and maintaining lawns and landscaping "

"pSEP","12"," Septic tank absorption field laterals should be installed on a slight downslope gradient."

"qLRS","12","This soil has no significant limitations important in the construction of local roads and streets."

Map Units 34, 40, 42, 43, 44, 45

"oURB","15","This soil survey map unit is so variable that no general interpretations can be given. On-site investigation by a soil scientist and/or engineer is recommended."

"pSEP","15","This soil survey map unit is so variable that no general interpretations for the installation of any type on-site sewage disposal system can be given. On-site investigation by a soil scientist and/or engineer is recommended."

"qLRS","15","These soils have no significant limitations for local roads and streets."

Map Units 4, 5

"oURB","20","This soil is moderately suited to most urban land uses. Because of the very rapid permeability of this soil, careful selection of on-site waste disposal areas can help prevent contamination of shallow groundwater and adjacent surface waters. Irrigation, mulching, and fertilizing help establish and maintain lawns and landscaping plants."

"pSEP","20","These soils have no significant limitations for septic tank absorption fields."

"qLRS","20","These soils have no significant limitations for local roads and streets.."

Map Units 18, 19, 20, 21, 22, 23, 26

"oURB", "21", "This soil has a low suitability for urban uses because wetness and the low strength of the organic layers and the likelihood of subsidence if drained."

"pSEP", "21", "This soil has severe limitations for any on-site waste disposal system due to wetness and subsidence of the organic soil material."

"qLRS", "21", "This soil has severe limitations for local roads and streets due to wetness and subsidence of the organic soil material. Excavating and filling is required to assure roads function properly."

WATER QUALITY: PESTICIDE AND NUTRIENT MANAGEMENT

sWQ – Water Quality Statement

tPES – Pesticide Management Statement

uNUT – Nutrient Management Statement

Map Units None

"sWQ", "01", "These soils have a low potential for pesticide leaching to groundwater and a low potential for pesticide runoff to surface water. They have a medium or high potential for nitrogen leaching to groundwater and a low potential for phosphorous runoff to surface runoff."

"tPES", "01", "The Florida Pest Control Guide contains a listing of pesticides suitable for each type of pest and is available from the Cooperative Extension Service. Read and follow pesticide labels."

"uNUT", "01", "A soil test will be used as a guide to determine plant nutrient needs. In addition, a listing of nitrogen and phosphorous requirements by crop type is available from the Cooperative Extension Service. Nutrients should be added at the rate needed by the crop grown or according to the producer's goals, whichever is lower."

Map Units 1, 2, 4, 5, 6, 9, 11, 14, 28, 36

"sWQ", "02", "These soils have a medium or high potential for pesticide leaching to the groundwater and a low potential for pesticide runoff from the field(s) to surface water. They have a medium or high potential for nitrogen leaching to the groundwater and a low potential for phosphorous runoff to surface runoff."

"tPES", "02", "The Florida Pest Control Guide from the Cooperative Extension Service contains a list of pesticides suited to each pest. This list also contains Relative Leaching Potential Index (RLPI) values. While any approved pesticide listed in the guide can be used, the applicator should consider for use pesticides with a larger RLPI value and Health Advisory Level (HAL or HALEQ) value. Read and follow pesticide labels."

"uNUT", "02", "A soil test will be used as a guide to determine plant nutrient needs. In addition, a listing of nitrogen and phosphorous requirements by crop type is available from the Cooperative Extension Service. Nutrients shall be added at the rate needed by the crop grown or according to the producer's goals, whichever is lower."

Map Units 3, 7, 8, 10, 12, 13, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 29, 30, 31, 33, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46

"sWQ", "03", "These soils have a medium or high potential for pesticide leaching to groundwater and a medium to high potential for pesticide runoff to surface water. They have a medium or high potential for nitrogen leaching to the groundwater and a medium or high potential for phosphorous runoff to surface runoff."

"tPES", "03", "The Florida Pest Control Guide from the Cooperative Extension Service contains a list of pesticides suited to each pest. This list also contains Relative Leaching Potential Index (RLPI) and Relative Runoff Potential Index (RRPI) values. While any approved pesticide listed in the guide can be used, the applicator should consider for use pesticides with a larger RLPI value, RRPI value, Health Advisory Level (HAL or HALEQ) value, and Aquatic Toxicity value. Read and follow pesticide labels."

"uNUT", "03", "A soil test will be used as a guide to determine plant nutrient needs. In addition, a listing of nitrogen and phosphorous requirements by crop type is available from the Cooperative Extension Service. Nutrients shall be added at the rate needed by the crop grown or according to the producer's goals, whichever is lower."

Map Units None

"sWQ", "04", "These soils have a low potential for pesticide leaching to groundwater and a medium or high potential for pesticide runoff to surface water. They have a medium or high potential for nitrogen leaching to groundwater and a medium or high potential for phosphorous runoff to surface runoff."

"tPES", "04", "The Florida Pest Control Guide from the Cooperative Extension Service contains a listing of pesticides suited to each pest. This list also contains Relative Runoff Potential Index (RRPI) values. While any approved pesticide listed in the guide can be used, the applicator should consider for use pesticides with a larger RRPI value and a larger Aquatic Toxicity value. Read and follow pesticide labels."

"uNUT", "04", "A soil test will be used as a guide to determine plant nutrient needs. In addition, a listing of nitrogen and phosphorous requirements by crop type is available from the Cooperative Extension Service. Nutrients shall be added at the rate needed by the crop grown, or according to the producer's goals, whichever is lower."