

SECTION II
TABLE OF CONTENTS

II – PASTURELAND AND HAYLAND INTERPRETATIONS

Introduction

Forage Suitability Groups

MLRA _____ 4 ETa Zone _____
MLRA _____ 4 ETa Zone _____
MLRA _____ 4 ETa Zone _____
MLRA _____ 4 ETa Zone _____

II – PASTURELAND AND HAYLAND INTERPRETATIONS

INTRODUCTION

Pastureland refers to grazing lands composed of introduced or domesticated native forage species that are used primarily for the production of domestic livestock. They receive periodic renovation and/or cultural treatments, such as tillage, fertilization, mowing, weed control, and may be irrigated. They are not in rotation with crops. Pastureland is principally harvested by grazing animals, but may be machine harvested to accumulate stored forage.

Hayland refers to land on which perennial plants are managed and harvested for hay for six years or more. Hayland is generally machine harvested, but may be grazed. Land where annual plants are planted for hay or forage crops grown in short-term rotation is cropland.

Landscapes are divided into basic units for study, evaluation, and management. On rangelands and forest lands, these units are called ecological sites; while on forage croplands, hayland, and pasturelands, they are forage suitability groups.

Forage suitability groups (FSGs) are composed of one or more individual soil map units having similar potentials and limitations for forage production. Soils within a forage production suitability group are sufficiently uniform to:

- Support the same adapted forage plants under the same management conditions;
- Require similar conservation treatment and management to produce the forages selected in the quality and quantity desired; and
- Have comparable potential productivity.

Forage suitability groups (FSGs) condense and simplify soils information. They provide the soil and plant science information for planning. The forage suitability groups description contains the soil map units that make up the FSG, adapted forage species and planting mixture, limitations of the FSG, conservation problems associated with the various limitations, annual forage production estimates, and distribution of production during the growing season.

FSGs will be established for each Major Land Resource Area (MLRA) having significant forage production. They will be filed in the *Forage Suitability Groups* subsection, located at the end of Section II of the Technical Guide.

II – PASTURELAND AND HAYLAND INTERPRETATIONS

FORAGE SUITABILITY GROUPS

Forage Suitability Groups (FSGs) have not yet been developed for soils in the Field Office service area. Once developed, they will be located in the *Forage Suitability Groups* subsection at the end of Section II of the FOTG. Data currently available for interpreting soils for pasture and hayland uses are in the published soil survey reports in the section on “Estimated yields and management guides.” FSGs will be developed for the irrigated condition (all 4 ETa Zones) and for the non-irrigated condition (4 ETa zones b, c, d, and e only) in each Major Land Resource Area found in the section on “Estimated yields and management guides.”

Irrigated FSGs will be based on soils according to the following Vegetative Soil Group combinations:

1. A+B+C;
2. D+G;
3. E+F;
4. H+I;

and, according to the following forage mixtures:

1. Perennial grass alone;
2. Perennial grass and a legume;
3. Perennial grass mixture; and
4. Perennial grass mixture and a legume or legume mixture.

Non-irrigated FSGs will be based on soils according to the following Vegetative Soil Group combinations:

1. A+C;
2. B alone;
3. C + E + F;
4. D+G;
5. H + I;

and, according to the following forage mixtures:

1. Perennial grass alone;
2. Perennial grass and a legume;
3. Perennial grass mixture;
4. Perennial grass mixture and a legume or legume mixture;
5. Annual grass alone;
6. Annual grass and legume; and
7. Annual grass mixture and a legume or legume mixture.

II – PASTURELAND AND HAYLAND INTERPRETATIONS

VEGETATIVE SOIL GROUPS

Vegetative Soil Groups are described below and in the table following. Planting recommendations can be found in the Vegetative Guide.

- A. ALL CLIMATICALLY ADAPTED PLANTS SUITED. Soils are deep to very deep, moderately coarse to medium textured, moderately well to well-drained, moderately rapidly to moderately slowly permeable. (Soils in this group can have slight wetness and slight salinity or alkalinity.)
- B. Choice of plants limited by DROUTHINESS AND LOW FERTILITY LEVEL. Soils are coarse to gravelly medium textured, excessively drained, with less than five inches of available water capacity (AWC) in the root zone.
- C. Choice of plants limited by FINE TEXTURES. Soils are deep to very deep, moderately fine to fine-textured, moderately well-drained, moderately slowly to slowly permeable.
- D. Choice of plants limited by VERY SLOWLY PERMEABLE (CLAYPAN) SUBSOILS. Soils are moderately well-drained, with slow or very slow subsoil permeability.
- E. Choice of plants limited by WETNESS. Soils are somewhat poorly to very poorly drained. (Drained soil phases will be placed in appropriate group according to their current drainage status.) Slight salinity and/or alkalinity may be present.
- F. Choice of plants limited by SALINITY OR ALKALINITY. Soils are moderately to strongly saline alkaline, and usually somewhat poorly or poorly drained.
- G. Choice of plants limited by DEPTH. Soils are shallow to moderately deep, well-drained, over hardpan, bedrock or other unfractured dense material.
- H. Choice of plants limited by LOW pH. Soils are strongly to extremely acidic; pH is less than 5.6.
- I. Choice of plants limited by TOXIC PROPERTIES. Soils are usually moderately to strongly serpentinitic.
- J. Choice of plants depends upon ON-SITE INVESTIGATION. Soils include those in the miscellaneous non-arable category, such as river wash, or stony or rocky upland.

Grouping Soils for Vegetative Purposes - California

Veg. Group Symbol	Major Soil Limitation	Effective Depth (in.)	Surface Texture	Subsoil Permeability 1/	Drainage Class 2/	Salinity & Alkalinity 3/	Reaction 4/	Erosion	AWC (in.) 5/
A	None	36 or more	sl through sicl	Moderately rapid through slow	Moderately well through well	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	5 or more
B	Drouthiness	36 or more	s, ls, gls, vg, k	Very rapid through very slow	Excessively through moderately well	None through slight	Strongly acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	5 or less
C	Fine textures	20 or more	c, sic, gc	Moderate through slow	Moderately well through well	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	5 or more
D	Clay Pan 6/	10 through 36	sl through sicl	Slow or very slow	Well through somewhat poorly	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	3 or more
E	Wetness	20 or more	s through c	Rapid through slow	Somewhat poorly through very poorly	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	3 or more
F	Salinity or alkalinity	20 or more	s through c	Rapid through slow	Moderately well through poorly	Moderate through strong	Neutral through very strongly alkaline (pH 6.6-9.0+)	Slight through moderate	3 or more
G	Shallow depth 7/	10 through 36 10 through 20	s, ls, gsl vg, k sl, thr sicl, c, gc	Moderately rapid through very slow	Moderately well through somewhat excessively	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	3 or more
H	Low pH	20 or more	sl through silc	Moderately rapid through mod. Slow	Somewhat poorly through somewhat excessively	None	Strongly acid through extremely acid (pH less than 5.6)	Slight through moderate	3 or more
I	Toxicity (serpentine soils)	10 or more	1 through c	Moderately rapid through mod. Slow	Somewhat poorly through somewhat excessively	None through slight	Medium acid through mod. Alkaline (pH 5.6-8.4)	Slight through moderate	3 or more
J	Severe 8/	any	any	Very rapid through very slow	Excessively through very poor	None through slight	any	Slight through severe	any

TABLE: Grouping Soils for Vegetative Purposes - California

FOOTNOTES:

All terms are standard. For definitions see NRCS TN-Soils-9 (rev.), March 1967, or Agr. Handbook No. 18, USDA-NRCS. Criteria underlined are main soil feature determining vegetative group.

1/ Subsoil permeability refers to permeability of the B horizon(s) or the 10- to 40-inch control section in soils without B horizons.

2/ Drainage class refers to drainage of soils that do not have altered drainage. If the soils have been drained, use class that most nearly reflects growing conditions following drainage improvement.

3/ Use current levels of salinity and alkalinity that are present in the field. Levels may be higher or lower than indicated on maps. Capability unit designations may be based on general assumptions that do not uniformly reflect current, short-term growing conditions on each parcel of land.

4/ Generally applies to the soil to a depth of 20 inches.

5/ Limits are for total available water-holding capacity for that part of the soil profile generally available to roots or to a depth of 60 inches if no severe intervening restrictions of soil or water are present. Refer to California Soil Handbook, Chapter 3, section 3.423.

6/ Soils in this group must have a clay increase of at least 15 percent, absolute, within 1 inch, or an abrupt or very abrupt AB boundary.

7/ Depth to unfractured rock or hardpan. If a claypan over 6 inches thick is present over rock or hardpan, place in Group D. See note 6 for other claypan criteria.

8/ Includes all soils not suitable for routine cultivation, seeding, and planting. Includes all class VII and VIII land, very cobbly soils, soils in class 3, 4, and 5 rockiness, class 2, 3, 4, and 5 stoniness. These soils require on-site recommendations