

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Coarse Upland (CU) 10-14” Foothills and Basins East Precipitation Zone

Site ID: R032XY308WY

Major Land Resource Area: 32 – Northern Intermountain Desertic Basins

Physiographic Features

This site occurs on undulating rolling land.

Landform: Hillsides, alluvial fans, ridges & stream terraces

Aspect: N/A

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	5700	7500
Slope (percent):	0	30
Water Table Depth (inches):	None within 60 inches	
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):	0	0
Frequency:	None	None
Duration:	None	None
Runoff Class:	negligible	high

Climatic Features

Annual precipitation ranges from 10-14 inches per year. The normal precipitation pattern shows the least amount of precipitation in December, January, and February, increasing to a peak during the latter part of May. Amounts decrease through June, July, and August and then increase some in September. Much of the moisture that falls in the latter part of the summer is lost by evaporation and much of the moisture that falls during the winter is lost by sublimation. Average snowfall exceeds 20 inches annually. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Winds are generally not strong as compared to the rest of the state. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 75 mph.

Growth of native cool-season plants begins about April 15 and continues to about July 15. Cool weather and moisture in September may produce some green up of cool season plants that will continue to late October.

The following information is from the “Thermopolis 2” climate station:

	<u>Minimum</u>	<u>Maximum</u>	<u>5 yrs. out of 10 between</u>
Frost-free period (days):	74	149	May 23 – September 16
Freeze-free period (days):	112	180	May 8 – October 1
Annual Precipitation (inches):	7.6	21.9	

Mean annual precipitation: 12.35 inches

Mean annual air temperature: 46.2 °F (30.1°F Avg. Min. to 62.3°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include “Grass Creek 1E”, “Thermopolis”, Thermopolis 25NW”, “Buffalo Bill Dam” and “Black Mountain”.

Influencing Water Features

Wetland Description:	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
None	None	None	None	None

Stream Type: None

Representative Soil Features

The soils of this site are deep to moderately deep (greater than 20" to bedrock), moderately well to somewhat excessively well-drained & moderately slow to moderately rapid permeable. This site consists of bouldery to cobbly coarse fragment soils. The soil surface can be covered extensively with these coarse fragments and as such, plant density can be reduced. The soil characteristics having most influential to the plant community are volume of coarse fragments in the profile that reduces the available moisture and the extensive cover of these coarse fragments, which can reduce the plant density.

Major Soil Series correlated to this site includes: Brownsto

Other Soil Series correlated to this site in MLRA 32 include:

Parent Material Kind: alluvium and residuum

Parent Material Origin: sandstone, shale

Surface Texture: loam, silt loam, very fine sandy loam,

Surface Texture Modifier: cobbly or bouldery

Subsurface Texture Group: loam

Surface Fragments ≤ 3” (% Cover): 5 to 25

Surface Fragments > 3” (%Cover): 0 to10

Subsurface Fragments ≤ 3” (% Volume): 0 to 25

Subsurface Fragments > 3” (% Volume): > 35

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	well	somewhat excessively well
Permeability Class:	moderately slow	moderately rapid
Depth (inches):	20	>60
Electrical Conductivity (mmhos/cm) ≤ 20" :	0	4
Sodium Absorption Ratio ≤ 20" :	0	5
Soil Reaction (1:1 Water) ≤ 20" :	7.4	8.4
Soil Reaction (0.1M CaCl₂) ≤ 20" :	NA	NA
Available Water Capacity (inches) ≤ 30" :	2.8	6.0
Calcium Carbonate Equivalent (percent) ≤ 20" :	0	30

Plant Communities

Ecological Dynamics of the Site:

Potential vegetation on this site is dominated by mid cool-season perennial grasses. Other significant vegetation includes big sagebrush and a variety of forbs. The expected potential composition for this site is about 75% grasses, 10% forbs and 15% woody plants. The composition and production will vary naturally due to historical use, fluctuating precipitation and fire frequency.


As this site deteriorates species such as threadleaf sedge, Sandberg bluegrass, blue grama, and big sagebrush will increase. Plains pricklypear and weedy annuals will invade. Cool-season grasses such as Griffiths and bluebunch wheatgrass, rhizomatous wheatgrasses, and needleandthread will decrease in frequency and production.

Big sagebrush may become dominant on some areas with an absence of fire. Wildfires are actively controlled in recent times so chemical control using herbicides has replaced the historic role of fire on this site. Recently, prescribed burning has regained some popularity.

Due to the amount and pattern of the precipitation, the big sagebrush component typically is not resilient once it has been removed if a healthy and vigorous stand of grass exists and is maintained. The exception to this is where the herbaceous component is severely degraded at the time of treatment, growing conditions are unfavorable after treatment, and/or recovery periods are inadequate.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



- BM** - Brush Management (fire, chemical, mechanical)
- Freq. & Severe Grazing** - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season
- GLMT** - Grazing Land Mechanical Treatment
- LTPG** - Long-term Prescribed Grazing
- MCSLG** - Moderate, Continuous Season-long Grazing
- NU, NF** - No Use and No Fire
- PG** - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)
- VLTPG** - Very Long-term Prescribed Grazing (could possibly take generations)
- WF** - Wildfire (Natural or Human Caused)

Plant Community Composition and Group Annual Production
Reference Plant Community (HCPC)

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Annual Production (Normal Year)		
			Group	lbs./acre	% Comp.
			Total: 650		
GRASSES AND GRASS-LIKES					
GRASSES/GRASSLIKES					
Griffiths wheatgrass or	Elymus albicans	ELAL7	1	163 - 228	25 - 35
Bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6			
Indian ricegrass	Achnatherum hymenoides	ACHY	2	98 - 163	15 - 25
Rhizomatous wheatgrass	Pascopyrum smithii	PASM	3	33 - 98	5 - 15
Spikefescue	Leucopoa kingii	LEKI2	4	0 - 65	0 - 10
Needleandthread grass	Hesperostipa comata	HECO26	5	0 - 65	0 - 10
MISC. GRASSES/GRASSLIKES			6	0 - 65	0 - 10
Blue grama	Bouteloua gracilis	BOGR2	6	0 - 33	0 - 5
Canby bluegrass	Poa canbyi (syn. P. secunda)	POCA(POSE)	6	0 - 33	0 - 5
Prairie junegrass	Koeleria macrantha	KOMA	6	0 - 33	0 - 5
Red threeawn	Aristida purpurea	ARPUL	6	0 - 33	0 - 5
Sandberg bluegrass	Poa secunda	POSE	6	0 - 33	0 - 5
Threadleaf sedge	Carex filifolia	CAFI	6	0 - 33	0 - 5
other perennial grasses (native)		2GP	6	0 - 33	0 - 5
FORBS			7	0 - 65	0 - 10
Aster	Eucephalus spp.	EUCEP2	7	0 - 33	0 - 5
Fringed sagewort	Artemisia frigida	ARFR4	7	0 - 33	0 - 5
Gromwell	Buglossoides arvensis	BUAR3	7	0 - 33	0 - 5
Larkspur	Delphinium spp.	DELPH	7	0 - 33	0 - 5
Miner's candle	Cryptantha virgata	CRVI4	7	0 - 33	0 - 5
Paintbrush	Castilleja spp.	CAST	7	0 - 33	0 - 5
Penstemon	Penstemon spp.	PENST	7	0 - 33	0 - 5
Phlox	Phlox spp.	PHLOX	7	0 - 33	0 - 5
Pussytoes	Antennaria rosea	ANRO2	7	0 - 33	0 - 5
Scarlet globemallow	Sphaeralcea coccinea	SPCO	7	0 - 33	0 - 5
Sulphur flower buckwheat	Eriogonum umbellatum	ERUM	7	0 - 33	0 - 5
Wavyleaf paintbrush	Castilleja applegatei martinii	CAAPM	7	0 - 33	0 - 5
Western yarrow	Achillea lanulosa	ACHIL	7	0 - 33	0 - 5
other perennial forbs (native)		2FP	7	0 - 33	0 - 5
TREES/SHRUBS			8	33 - 98	5 - 15
Antelope bitterbrush	Purshia tridentata	PUTR2	8	0 - 33	0 - 5
Big sagebrush	Artemisia tridentata	ARTR2	8	0 - 33	0 - 5
Black sagebrush	Artemisia nova	ARNO4	8	0 - 33	0 - 5
Green rabbitbrush	Chrysothamnus viscidiflorus	CHVI8	8	0 - 33	0 - 5
Rubber rabbitbrush	Ericameria nauseosa	ERNA10	8	0 - 33	0 - 5
Utah juniper	Juniperus osteosperma	JUOS	8	0 - 33	0 - 5
other shrubs & half shrubs (native)		2SHRUB	8	0 - 33	0 - 5

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

Bluebunch Wheatgrass/Indian Ricegrass Plant Community

This plant community is the interpretive plant community for this site and is considered to be the Historic Climax Plant Community (HCPC). This state evolved with grazing by large herbivores and periodic fires. This plant community can be found on areas that are properly managed with grazing and/or prescribed burning, and on areas receiving occasional short periods of rest. The potential vegetation is about 75% grasses or grass-like plants, 10% forbs, and 15% woody plants. This state is dominated by cool season mid-grasses.

The major grasses include bluebunch wheatgrass, Indian ricegrass, rhizomatous wheatgrasses, spikescue, and needleandthread. Other grasses occurring in this state include blue grama and Sandberg’s bluegrass. A variety of forbs occurs on this site, as shown in the preceding table. Big sagebrush, black sagebrush, and rubber rabbitbrush are conspicuous species in this plant community. Antelope bitterbrush will occur on sites associated with the upper limits of this precipitation zone. These shrubs occur in a mosaic pattern, and make up 15% of the annual production. Plant diversity is high (see Plant Composition Table).

The total annual production (air-dry weight) of this state is about 650 lbs./acre, but it can range from about 400 lbs./acre in unfavorable years to about 900 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	25	40	10	5	10	5	0	0

(Monthly percentages of total annual growth)

This plant community is extremely stable and well adapted to the Northern Intermountain Desertic Basins climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Moderate, continuous season-long grazing will convert the plant community to the *Perennial Grass/Mixed Shrub Plant Community*. Prolonged drought will exacerbate this transition.

Perennial Grass/Mixed Shrub Plant Community

Historically, this plant community evolved under grazing by large ungulates and a low fire frequency. Currently, it is found under moderate, season-long grazing by livestock, and will be exacerbated by prolonged drought conditions. In addition, the fire regime for this site has been modified and extended periods without fire is now common. Big sagebrush and rubber rabbitbrush are significant components of this plant community. Cool-season grasses make up the majority of the understory with the balance made up of short warm-season grasses, annual cool-season grasses, and miscellaneous forbs.

Dominant grasses include bluebunch wheatgrass, needleandthread, and western wheatgrass. Grasses of secondary importance include threadleaf sedge, blue grama, Sandberg bluegrass, and red threeawn. Forbs commonly found in this plant community include scarlet globemallow, Hood's phlox, sulfur flower buckwheat, and penstemon spp. Shrubs can make up to 25% of the total annual production. Plains pricklypear can also occur.

When compared to the Historic Climax Plant Community, sagebrush, rubber rabbitbrush, juniper, blue grama, and threadleaf sedge have increased. Production of cool-season grasses, particularly bluebunch wheatgrass and Indian ricegrass, has been reduced. Some weedy species may have invaded the site but are in small patches such as cheatgrass, kochia, and Russian thistle.

The total annual production (air-dry weight) of this state is about 500 pounds per acre, but it can range from about 300 lbs./acre in unfavorable years to about 700 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	25	40	10	5	10	5	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. The herbaceous component is mostly intact and plant vigor and replacement capabilities are sufficient. Water flow patterns and litter movement may be occurring but only on steeper slopes. Incidence of pedestalling is minimal. Soils are mostly stable and the surface shows minimum soil loss. The watershed is functioning and the biotic community is intact.

Transitions or pathways leading to other plant communities are as follows:

- Prescribed grazing or possibly long-term prescribed grazing will convert this plant community to the HCPC. The probability of this occurring is high especially if rotational grazing along with short deferred grazing is implemented as part of a prescribed method of use. In addition, the removal of fire suppression will allow a somewhat natural fire regime to reoccur to more easily transition between this plant community and the HCPC. A prescribed fire treatment can be useful to hasten this transition if desired.

- Frequent and severe grazing plus no fires will convert the plant community to the *Dense Shrub/Bare Ground Plant community*. The probability of this occurring is high. This is especially evident on areas where drought or heavy browsing does not adversely impact the shrub stand.
- Frequent and severe grazing (yearlong) plus wildfire or brush management, will convert the plant community to a *Threadleaf Sedge and Blue Grama Sod/Bare Ground Plant Community*. The probability for is high especially on areas were the shrubs have been heavily browsed or removed by natural or human causes. Drought can also exasperate this transition.

Dense Shrub/Bare Ground Plant Community

This plant community is the result of frequent and severe grazing and protection from fire. Sagebrush and rubber rabbitbrush dominate this plant community as the annual production of shrubs exceeds 25%. Shrubs comprise the significant component of the plant community and the preferred cool season grasses have been eliminated or greatly reduced.

The dominant grasses are Sandberg bluegrass, threadleaf sedge, and blue grama. Weedy annual species such cheatgrass, kochia, Russian thistle, halogeton and a variety of mustards may occupy the site. Cactus has significantly increased. Noxious weeds such as Russian knapweed, leafy spurge, or Canada thistle may invade the site if a seed source is available. Plant diversity is moderate to poor. Cool-season grasses still exist, but are usually under the shrub canopy or in the cactus patches. The interspaces between plants have expanded leaving the amount of bare ground more prevalent. When compared with the HCPC or the Perennial Grass/ Mixed Shrub Plant Communities, the annual production is less, as the cool season grasses are reduced, but the shrub production compensates for the decline in some of the herbaceous production.

The total annual production (air-dry weight) of this state is about 400 pounds per acre, but it can range from about 250 lbs./acre in unfavorable years to about 650 lbs./acre in above average years.

The following is the growth curve of the plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	25	40	10	5	10	5	0	0

(Monthly percentages of total annual growth)

This plant community is resistant to change as the stand becomes more decadent. These areas may actually be more resistant to fire as less fine fuels are available and the bare ground between the shrubs is increased. The herbaceous component is not as diverse and plant vigor and species regeneration capabilities of cool-season perennials are deficient. The removal of grazing does not seem to affect the plant composition or structure of the plant community.

Soil erosion is accelerated because of increased bare ground. Water flow patterns and pedestalling are obvious. Infiltration is reduced and runoff is increased. Rill channels may be noticeable in the interspaces and gullies may be establishing where rills have concentrated down slope.

Transitions or pathways leading to other plant communities are as follows:

- Brush management, followed by prescribed grazing and seeding, will return this plant community to at or near the *HCPC*. If prescribed fire is used as a means to reduce or remove the shrubs, sufficient fine fuels will need to be present. This may require deferment from grazing prior to treatment. Post management is critical to ensure success. This can range from two or more years of rest to partial growing season deferment, depending on the condition of the understory at the time of treatment and the growing conditions following treatment. In the case of an intense wildfire that occurs when desirable plants are not completely dormant, the length of time required to reach the *HCPC* may be increased and seeding of natives are recommended.
- Brush management, followed by frequent and severe grazing, will convert the plant community to a *Threadleaf Sedge and Blue Grama Sod/Bare Ground Plant community*. The probability of this occurring is high, because of the amount of bare ground exposed to weedy annuals and sod formers as the competition for space, has been removed.

Threadleaf Sedge and Blue Grama Sod/Bare Ground Plant Community

This plant community is created when the Perennial Grass/Mixed Shrub Plant Community is subjected to severe heavy yearlong grazing and the shrub component has been removed by heavy browsing, wildfire or human means. Additionally, this plant community can occur as a result of the Dense Shrub/Bare Ground Plant Community being subjected to fire or brush management and not followed by prescribed grazing. Weedy annuals, threadleaf sedge, and/or blue grama are the most dominant plants. Weedy annuals occupy any open bare ground areas, while threadleaf sedge and/or blue grama form patches of sod in between the exposed boulders or cobbles. Rubber rabbitbrush may or may not be present on this site as this species may quickly re-establish after a fire. However, heavy browsing by large ungulates will significantly reduce or remove this shrub.

Compared to the HCPC, weedy annual species are widespread and may include cheatgrass, kochia, Russian thistle, halogeton and a variety of mustards. Cactus has increase significantly. Noxious weeds such as Russian knapweed or Canada thistle may invade the site if a seed source is available. Virtually all other cool-season mid-grasses are absent or severely decreased. Blue grama and threadleaf sedge have significantly increased from what is found in the HCPC. Shrubs have been removed with the exception of small patches of mostly rubber rabbitbrush.

The total annual production (air-dry weight) of this state is about 150 pounds per acre, but it can range from about 100 lbs./acre in unfavorable years to about 300 lbs./acre in above average years.

The following is the growth curve of the plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	5	25	40	10	5	10	5	0	0

(Monthly percentages of total annual growth)

This plant community is relatively stable and resistant to overgrazing. Annuals and sod forming grasses are effectively competing against the establishment of perennial cool-season grasses. Plant diversity is greatly altered and the herbaceous component is not intact. Recruitment of perennial grasses is not occurring and the replacement potential is absent. The biotic integrity is missing.

On areas with a well established sod plant community, water infiltration will be significantly affected. While this sod protects the area itself, adjacent on-site and off-site areas are impacted by excessive runoff that can cause rill channels and gully erosion. Water flow patterns and pedestalling are obvious. The watershed may or may not be functional.

Transitions or pathways leading to other plant communities are as follows:

- Grazing land mechanical treatment (chiseling, etc.) and pricklypear cactus control (if needed), followed by prescribed grazing, will return this plant community to near *Historic Climax Plant Community* condition. The sod areas are extremely resistant to change and will require grazing land mechanical treatments, such as chiseling to revert to a more preferred state. This may not be possible given the presence of cobbles or boulders on the soil surface, which can also exclude reseeding the area. If applicable, mechanical treatments and reseeding native plant species are recommended. This should be followed by proper grazing management to accelerate recovery where few desirable plants remain.

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

Bluebunch Wheatgrass/Indian Ricegrass Plant Community (HCPC): The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

Perennial Grass/Mixed Shrub Plant Community: The combination of an overstory of sagebrush and an understory of grasses and forbs provide a very diverse plant community for wildlife. The crowns of sagebrush tend to break up hard crusted snow on winter ranges, so mule deer and antelope may use this state for foraging and cover year-round, as would cottontail and jack rabbits. It provides important winter, nesting, brood-rearing, and foraging habitat for sage grouse. Brewer's sparrows' nest in big sagebrush plants, and hosts of other nesting birds utilize stands in the 20-30% cover range.

Dense Shrub/Bare Ground Plant Community: This plant community can provide important winter foraging for elk, mule deer and antelope, as sagebrush can approach 15% protein and 40-60% digestibility during that time. This community provides escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse.

Threadleaf Sedge and Blue Grama Sod/Bare Ground Plant Community: These communities provide limited foraging for antelope and other grazers. They may be used as a foraging site by sage grouse if proximal to woody cover and if the Historic Climax Plant Community or the Perennial Grass/Mixed Shrub Plant Community is limiting. Generally, these are not target plant communities for wildlife habitat management.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 32, 10-14 inch Foothills and Basins East

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope	Elk	Moose	Mtn. Sheep
GRASSES/GRASSLIKES										
Alkali bluegrass	Poa juncea (syn. P. secunda)	POJU (POSE)	DDDD	PPPP	DDDD	PPPP	PPPP	DDDD	DDDD	DDDD
Alkali cordgrass	Spartina gracilis	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	DDDD	UUUU
Alkali sacaton	Sporobolus airoides	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD	DDDD
Baltic rush	Juncus balticus	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	UUUU	UUUU
Basin wildrye	Leymus cinereus	LECI4	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD	PPPP
Big bluegrass	Poa Ampla (syn. P. secunda)	POAM (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Blue grama	Bouteloua gracilis	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	PPPP	DDDD
Bluejoint reedgrass	Calamagrostis canadensis	CACAM	PPPP	DDDD	PPPP	UUUU	UUUU	PPPP	DDDD	DDDD
Bottlebrush squirreltail	Elymus elymoides	ELELE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Bulrush	Scirpus spp.	SCRIP	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD	DDDD	DDDD
Canada wildrye	Elymus canadensis	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	PPPP	PPPP
Canby bluegrass	Poa canbyi (syn. to P. secunda)	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Golden sedge	Carex aurea	CAAU3	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD	UUUU	DDDD
Green needlegrass	Nassella viridula	NAV14	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Indian ricegrass	Achnatherum hymenoides	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Inland saltgrass	Distichlis spicata	DISP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Inland sedge	Carex interior	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD	DDDD	DDDD
Mat muhly	Muhlenbergia richardsonis	MURI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Griffith's wheatgrass	Elymus albicans	ELAL7	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Nebraska sedge	Carex nebrascensis	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD	DDDD
Needleandthread	Hesperostipa comata	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Nuttall's alkaligrass	Puccinellia nuttalliana	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Prairie junegrass	Koeleria macrantha	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Prairie sandreed	Calamovilfa longifolia	CALO	PPPP	DDDD	PPPP	UUUU	UUUU	PPPP	DDDD	DDDD
Sandberg bluegrass	Poa secunda	POSE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Sand dropseed	Sporobolus cryptandrus	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD	UUUU	UUUU
Slender wheatgrass	Elymus trachycaulus	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP	DDDD	DDDD
Slough sedge	Carex obnupta	CAOB3	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Spike fescue	Leucophaea kingii	LEKI2	PPPP	DDDD	PPPP	PPPP	DDDD	DDDD	DDDD	DDDD
Streambank wheatgrass	Elymus lanceolatus	ELLAL3	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Thickspike wheatgrass	Elymus lanceolatus	ELLAL3	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Threadleaf sedge	Carex filifolia	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP	DDDD	DDDD	DDDD
Tufted hairgrass	Deschampsia caespitosa	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP	DDDD	DDDD
Upland sedges	Carex spp.	CAREX	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Water sedge	Carex aquatilis	CAAQ	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Western wheatgrass	Pascopyrum smithii	PASM	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
FORBS										
Alkali seepweed	Suaeda vera	SUVE2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
American bistort	Polygonum bistortoides	POBI16	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Arrowgrass	Triglochin spp.	TRIGL	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Asters	Eucephalus spp.	EUCEP2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Biscuitroots	Lomatium spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD	DDDD
Cinquefoil	Potentilla spp.	POTEN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Deathcamas	Zigadenus Michx.	ZIGAD	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Dock	Rumex spp.	RUMEX	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Evening primrose	Oenothera caespitosa	OECA10	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
False carrot	Turgenia spp.	TURGE	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU	UUUU	DDDD
Fleabanes	Erigeron spp.	ERIGE2	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Fringed sagewort	Artemisia frigida	ARFR4	UUUU	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU
Goldenweed	Stenotus acaulis	STAC	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Gromwell	Buglossoides arvensis	BUAR3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Hawksbeard	Crepis acuminata	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD	UUUU	DDDD	DDDD
Horsetails	Equisetum spp.	EQUIS	UUUU	UUUU	TTTT	UUUU	UUUU	UUUU	UUUU	UUUU
Iris	Iris spp.	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Larkspur	Delphinium spp.	DELPH	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Milkvetch	Astragalus spp.	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Painbrush	Castilleja spp.	CAST	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Penstemons	Penstemon spp.	PENST	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Phlox	Phlox spp.	PHLOX	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Prairie thermopsis	Thermopsis rhombifolia	THRH	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Princessplume	Stanleya spp.	STANL	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Nuttall's povertyweed	Monoecis nuttalliana	MONU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Pussytoes	Antennaria spp.	ANTEN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Salsify	Tragopogon porrifolius	TRPO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Scarlet globemallow	Sphaeralcea coccinea	SPCO	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Stemless hymenoxys	Tetranneuris acaulis	TEACA2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Wild onion	Allium textile	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Winterfat	Krascheninnikovia lanata	KRAL2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Woody aster	Xylorhiza spp.	XYLOR	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Wooly groundsel	Packera cana	PACA15	TTTT	UUUU	TTTT	UUUU	UUUU	TTTT	UUUU	UUUU
TREES, SHRUBS & HALF-SHRUBS										
Antelope bitterbrush	Purshia tridentata	PUTR2	PPPP	PPPP	DDDD	PPPP	PPPP	PPPP	PPPP	PPPP
Boxelder	Acer negundo L. var. interius	ACNE12	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Silver sagebrush	Artemisia cana	ARCA13	DDDD	DDDD	PPPP	PPPP	PPPP	DDDD	DDDD	DDDD
Big sagebrush	Artemisia tridentata	ARTR2	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD	DDDD
Birdfoot sagebrush	Artemisia pedatifida	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Black sagebrush	Artemisia nova	ARNO4	UUUU	PPPP	UUUU	PPPP	PPPP	UUUU	UUUU	DDDD
Cottonwoods (sprouts)	Populus spp.	POPUL	DDDD	DDDD	DDDD	DDDD	UUUU	DDDD	DDDD	UUUU
Curleaf mountainmahogany	Cercocarpus ledifolius	CELE3	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP	DDDD
Gardners saltbush	Atriplex gardneri	ATGA	PPPP	PPPP	DDDD	PPPP	PPPP	PPPP	PPPP	DDDD
Greasewood	Sarcobatus vermiculatus	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	UUUU	UUUU
Green rabbitbrush	Chrysothamnus viscidiflorus	CHVI8	PPPP	DDDD	PPPP	PPPP	PPPP	PPPP	DDDD	DDDD
Limber pine	Pinus flexilis	PINF2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Rubber rabbitbrush	Ericameria nauseosa	ERNA10	UUUU	PPPP	UUUU	DDDD	PPPP	UUUU	UUUU	DDDD
Rocky Mountain juniper	Juniperus scopulorum	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU
Shadscale	Atriplex confertifolia	ATCO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Shrubby cinquefoil	Dasiphora floribunda	DAFL3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	DDDD	UUUU
Silver buffalobery	Shepherdia argentea	SHAR	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
skunkbush sumac	Rhus trilobata	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD	UUUU	UUUU
Snowberry	Symphoricarpos occidentalis	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU
Utah juniper	Juniperus osteosperma	JUOS	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU
Wildrose	Rosa woodsii var. woodsii	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	DDDD	DDDD
Willows	Salix spp.	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU	PPPP	PPPP	DDDD
Winterfat	Krascheninnikovia lanata	KRAL2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Yucca	Yucca spp.	YUCCA	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD	UUUU	DDDD

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (lb./ac)	Carrying Capacity* (AUM/ac)
Bluebunch Wheatgrass/Indian Ricegrass	400-900	.3
Perennial Grass/Mixed Shrub	300-700	.25
Dense Shrub/Bare Ground	250-650	.15
Threadleaf Sedge & Blue Grama Sod/B. Ground	100-300	.07

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration ranges from moderately slow to moderate. Runoff potential for this site varies from low to moderate depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational Uses

This site provides hunting opportunities for upland game species. The wide varieties of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood Products

No appreciable wood products are present on the site.

Other Products

None noted.

Supporting Information

Associated Sites

Shallow Sandy	032XY366WY
Sandy	032XY350WY
Shallow Clayey	032XY358WY
Gravelly	032XY312WY
Lowland	032XY328WY

Similar Sites

- () – Coarse Upland 5-9" Wind River Basin 032XY208WY
Coarse Upland 5-9" Big Horn Basin 032XY108WY
[lower production than Coarse Upland 10-14" E 032XY308WY]

Inventory Data References (narrative)

Information presented here has been derived from NRCS inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Chris Krassin, Range Management Specialist, NRCS and Everet Bainter, Range Management Specialist. Other sources used as references include USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, USDI and USDA Interpreting Indicators of Rangeland Health Version 3, and USDA NRCS Soil Surveys from various counties.

Inventory Data References

Ocular field estimations observed by trained personnel.

State Correlation

This site occurs entirely within Wyoming.

Type Locality

Field Offices

Casper, Cody, Dubois, Fort Washakie, Greybull, Lander, Powell, Riverton, Thermopolis, Worland,

Relationship to Other Established Classifications

Other References

Site Description Approval

State Range Management Specialist

Date