

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Wetland (WL) 15-19” Black Hills Precipitation Zone,

Site ID: 061XY178WY

Major Land Resource Area: 61 – Black Hills Foot Slopes

Physiographic Features

This site normally occurs on level to nearly level bottomlands near springs, seeps and sloughs.

Landform: drainageways, oxbows, and stream terraces.

Aspect: N/A

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	3500	5000
Slope (percent):	0	6
Water Table Depth (inches):	0	18
Flooding:		
Frequency:	occasional	frequent
Duration:	very brief	brief
Ponding:		
Depth (inches):	0	12
Frequency:	frequent	frequent
Duration:	brief	very long
Runoff Class:	negligible	medium

Climatic features

Annual precipitation ranges from 15-19 inches per year. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Strong winds are less frequent than over other areas of Wyoming. Occasional storms, however, can bring brief periods of high winds with gusts exceeding 50 mph.

Growth of native cool season plants begins about April 1 and continues to about July 1. Native warm season plants begin about May 15 and continue to about August 15. Fall green-up may occur in September and last through October.

The following information is from the “Devils Tower 2” climate station:

	<u>Minimum</u>	<u>Maximum</u>	<u>5 yrs. out of 10 between</u>
Frost-free period (days) (32°F):	58	93	June 6 – September 7
Freeze-free period (days) (28°F):	95	125	May 18 – September 20
Annual Precipitation (inches):	14.81	20.17	

Mean annual precipitation: 17.66 inches

Mean annual air temperature: 44.4°F (28.6°F Avg. Min. to 60.1°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include “Hulett” and “Sundance”.

Influencing Water Features

Wetland Description:	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
	Palustrine	None	Emergent Wetland	Persistent

Stream Type: C (Rosgen)

Representative Soil Features

This site consists of deep to very deep poorly drained soils formed in alluvium with a water table above the surface for part but not all of the growing season. They are on nearly level to slightly depressed areas with poor surface drainage. In some places the surface layers have high organic matter content. Layers of the soil most influential to the plant community vary from 3 to 6 inches thick.

Parent Material Kind: alluvium

Parent Material Origin: sandstone, shale

Surface Texture: clay, clay loam, loam, silty clay, silty clay loam, silt loam

Surface Texture Modifier: mucky

Subsurface Texture Group: loam

Surface Fragments ≤ 3” (% Cover): 0

Surface Fragments > 3” (%Cover): 0

Subsurface Fragments ≤ 3” (% Volume): 0

Subsurface Fragments > 3” (% Volume): 0

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	very poorly drained	somewhat poor
Permeability Class:	slow	moderate
Depth (inches):	20	>60
Electrical Conductivity (mmhos/cm) ≤20”:	0	4
Sodium Absorption Ratio ≤20”:	0	5
Soil Reaction (1:1 Water) ≤20”:	6.6	7.8
Soil Reaction (0.1M CaCl2) ≤20”:	NA	NA
Available Water Capacity (inches) ≤30”:	2.2	6.6
Calcium Carbonate Equivalent (percent) ≤20”:	0	5


Plant Communities

Ecological Dynamics of the Site:

As this site deteriorates, species such as spike sedge and Baltic rush increase. Grasses and grass-like plants such as Nebraska sedge, tufted hairgrass, northern, and bluejoint reedgrass will decrease in frequency and production

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



BM - Brush Management (fire, chemical, mechanical)

Freq. & Severe Grazing - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season

GLMT - Grazing Land Mechanical Treatment

LTPG - Long-term Prescribed Grazing

MCSLG - Moderate, Continuous Season-long Grazing

NU, NF - No Use and No Fire

PG - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)

VLTPG - Very Long-term Prescribed Grazing (could possibly take generations)

Na - Moderate Sodium in Soil

**Plant Community Composition and Group Annual Production
Reference Plant Community (HCPC)**

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Annual Production (Normal Year) Total: 6000		
			Group	lbs./acre	% Comp.
GRASSES AND GRASS-LIKES					
GRASSES/GRASSLIKES					
Reedgrasses (Northern, Bluejoint)	Calamagrostis spp.	CALAM	1	600 - 1500	10 - 25
Northern reedgrass	Calamagrostis stricta	CAST13	1		
Bluejoint reedgrass	Calamagrostis canadensis	CACAM	1		
Nebraska sedge	Carex nebrascensis	CANE2	2	1500 - 3000	25 - 50
Tufted hairgrass	Deschampsia caespitosa	DECA18	3	600 - 1500	10 - 25
Baltic rush	Juncus balticus	JUBA	4	300 - 600	5 - 10
Spike sedge	Carex nardina	CANA2	5	300 - 600	5 - 10
Inland sedge	Carex interior	CAIN11	6	0 - 300	0 - 5
other perennial grasses (native)		2GP	7	0 - 300	0 - 5
FORBS			8	300 - 600	5 - 10
Arrowgrass	Triglochin spp.	TRIGL	8	0 - 300	0 - 5
Blue-eyed grass	Sisyrinchium spp.	SISYR	8	0 - 300	0 - 5
cattail, narrow-leaf	Typha angustifolia	TYAN	8	0 - 300	0 - 5
cattail, broad-leaf	Typha latifolia	TYLA	8	0 - 300	0 - 5
Horsetails	Equisetum spp.	EQUIS	8	0 - 300	0 - 5
Iris	Iris spp.	IRIS	8	0 - 300	0 - 5
Water hemlock	Cicuta spp.	CICUT	8	0 - 300	0 - 5
other perennial forbs (native)		2FP	8	0 - 300	0 - 5
TREES/SHRUBS					
Willows	Salix spp.	SALIX	9	0 - 300	0 - 5
other shrubs & half shrubs (native)		2SHRUB	10	0 - 300	0 - 5

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

Nebraska sedge, Northern reedgrass Community

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores and is well suited for grazing by domestic livestock. Potential vegetation is about 90% grasses or grass-like plants and 10% forbs. The major grasses/grass-likes include Nebraska sedge, northern and bluejoint reedgrass, and tufted hairgrass. Grasses/grass-likes of lesser importance are Baltic rush and spike sedge.

The total annual production (air-dry weight) of this state is about 6000 pounds per acre, but it can range from about 4500 lbs/acre in unfavorable years to about 7000 lbs/acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY1603

Growth curve name: 15-19BL, Free Water Sites

Growth curve description: Free Water Sites

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	15	25	25	15	10	10	0	0

(Monthly percentages of total annual growth)

The state is well adapted to the Black Hills Foot Slopes climatic conditions. It is a critical state providing water and habitat for the surrounding area. The diversity in plant species provides a variety of habitats for wildlife. It is resistant to drought due to a dependable water supply. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Moderate, continuous season-long grazing will convert this plant community to the *Baltic rush/Cattail Plant Community*.
- Frequent and Severe grazing will convert this plant community to the *Baltic rush/Smartweed Plant Community*.

Baltic rush/Cattail Plant Community

This plant community evolved under moderate grazing by domestic livestock. Dominant grasses include spike sedge, and Baltic rush. Cattails are present near open water. Willows are present near the dryer edges of this state.

When compared to the Historical Climax Plant Community, northern and bluejoint reedgrass, Nebraska sedge, and tufted hairgrass have decreased. Spike sedge, Baltic rush, and cattails have

increased. The abundant production and proximity to water make this state important for livestock and wildlife such as birds, mule deer, and antelope.

The total annual production (air-dry weight) of this state is about 3000 pounds per acre, but it can range from about 2000 lbs./acre in unfavorable years to about 4000 lbs./acre in above average years.

The following is the growth curve expected during a normal year:

Growth curve number: WY1603
Growth curve name: 15-19BL, Free Water Sites
Growth curve description: Free Water Sites

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	15	25	25	15	10	10	0	0

(Monthly percentages of total annual growth)

The state is stable and protected from excessive erosion. The biotic integrity of this plant community is usually intact. The watershed is usually functioning.

Transitional pathways leading to other plant communities are as follows:

- Prescribed grazing over the long-term will result in a plant community very similar to the *Historic Climax Plant Community*.
- Frequent and Severe grazing will convert this plant community to the *Baltic rush/Smartweed Plant Community*.

Baltic rush/Smartweed Plant Community

This plant community is the result of long-term improper grazing use. Baltic rush, smartweed, and cattails dominate this state. American licorice has invaded.

The total annual production (air-dry weight) of this state is about 1800 pounds per acre, but it can range from about 1200 lbs/acre in unfavorable years to about 2500 lbs/acre in above average years.

The following is the growth curve expected during an average year.

Growth curve number: WY1603
Growth curve name: 15-19BL, Free Water Sites
Growth curve description: Free Water Sites

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	0	15	25	25	15	10	10	0	0

(Monthly percentages of total annual growth)

Bare ground has increased. The soil of this state is not well protected. Degraded stream banks may erode. The watershed is functioning but may produce excessive runoff. The biotic community is at risk due to invasive plants.

Transitional pathways leading to other plant communities are as follows:

- Prescribed Grazing over the long-term will return this state to near *Historic Climax Plant Community*.

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

Nebraska sedge, Northern reedgrass Community: The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. This plant community may provide brood rearing/foraging areas for sage grouse. Other birds that would frequent this plant community include red-wing blackbirds, sandhill cranes, Wilson snipe, western meadowlarks, and golden eagles. Many small mammals would occur here.

Baltic rush/Cattail Plant Community: This plant community may be useful for the same large grazers that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for sage grouse when it occurs proximal to woody cover. Good grasshopper habitat equals good foraging for birds.

Baltic rush/Smartweed Plant Community: This plant community may be useful for the same large grazers that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals. It may provide some foraging opportunities for sage grouse when it occurs proximal to woody cover. Good grasshopper habitat equals good foraging for birds.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 61, 15-19 inch Black Hills

COMMON NAME/	SCIENTIFIC NAME	SCI. SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope
GRASSES/GRASSLIKES							
alkali bluegrass	<i>Poa secunda</i> ssp. <i>juncifolia</i>	POSEJ	DDDD	PPPP	DDDD	PPPP	PPPP
alkali cordgrass	<i>Spartina gracilis</i>	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU
alkali sacaton	<i>Sporobolus airoides</i>	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD
bearded wheatgrass	<i>Elymus caninus</i>	ELCA	PPPP	DDDD	PPPP	DDDD	DDDD
Big bluegrass	<i>Poa ampla</i> (syn. <i>To Poa secunda</i>)	POAM (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
big bluestem	<i>Andropogon gerardii</i>	ANGE	PPPP	PPPP	PPPP	DDDD	DDDD
blue grama	<i>Bouteloua gracilis</i>	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD
Blue wildrye	<i>Elymus glaucus</i>	ELGL	DDDD	DDDD	DDDD	DDDD	DDDD
bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD
bluejoint reedgrass	<i>Calamagrostis canadensis</i>	CACA4	PPPP	DDDD	PPPP	UUUU	UUUU
buffalograss	<i>Buchloe dactyloides</i>	BUDA	DDDD	DDDD	DDDD	DDDD	DDDD
Canada wildrye	<i>Elymus canadensis</i>	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD
Canby bluegrass	<i>Poa canbyi</i> (syn. <i>to Poa secunda</i>)	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
Columbia needlegrass	<i>Achnatherum nelsonii</i>	ACNE9	PPPP	PPPP	DDDD	DDDD	DDDD
Cusick's bluegrass	<i>Poa cusickii</i>	POCU3	PPPP	PPPP	PPPP	PPPP	PPPP
fowl bluegrass	<i>Poa palustris</i>	POPA2	DDDD	DDDD	DDDD	UUUU	UUUU
green needlegrass	<i>Nassella viridula</i>	NAV14	PPPP	PPPP	PPPP	PPPP	PPPP
hairy grama	<i>Bouteloua hirsuta</i>	BOHI2	DDDD	DDDD	DDDD	DDDD	DDDD
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP
inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU
inland sedge	<i>Carex interior</i>	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU
little bluestem	<i>Schizachyrium scoparium</i>	SCSC	PPPP	PPPP	PPPP	DDDD	DDDD
mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	<i>Carex nebraskensis</i>	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD
needleandthread	<i>Hesperostipa comata</i>	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP
needleleaf sedge	<i>Carex duriuscula</i>	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU
northern reedgrass	<i>Calamagrostis stricta</i>	CAS113	PPPP	DDDD	PPPP	UUUU	UUUU
Nuttall's alkaligrass	<i>Puccinellia nuttalliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP
plains reedgrass	<i>Calamagrostis montanensis</i>	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD
prairie cordgrass	<i>Spartina pectinata</i>	SPPE	PPPP	DDDD	PPPP	UUUU	UUUU
prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD
prairie sandreed	<i>Calamovilfa longifolia</i>	CALO	PPPP	DDDD	PPPP	UUUU	UUUU
Pumpelly brome	<i>Bromus inermis</i> spp. <i>pumpellianus</i>	BRINP5	PPPP	PPPP	DDDD	DDDD	UUUU
Richardson's needlegrass	<i>Achnatherum richardsonii</i>	ACRI8	PPPP	DDDD	DDDD	DDDD	DDDD
sand bluestem	<i>Andropogon halli</i>	ANHA	PPPP	DDDD	PPPP	UUUU	UUUU
sand dropseed	<i>Sporobolus cryptandrus</i>	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU
Sandberg bluegrass	<i>Poa secunda</i>	POSE	DDDD	DDDD	DDDD	DDDD	DDDD
sideoats grama	<i>Bouteloua curtipendula</i>	BOCU	PPPP	PPPP	PPPP	DDDD	UUUU
slender wheatgrass	<i>Elymus trachycaulus</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD
spike oatgrass	<i>Helictotrichon hookeri</i>	HEHO8	PPPP	DDDD	PPPP	DDDD	DDDD
spike sedge	<i>Carex nardina</i>	CANA2	DDDD	DDDD	DDDD	UUUU	UUUU
Spikefescue	<i>Leucopoa kingii</i>	LEK12	PPPP	DDDD	PPPP	PPPP	DDDD
stonehills (plains) muhly	<i>Muhlenbergia cuspidata</i>	MUCU3	UUUU	UUUU	UUUU	UUUU	UUUU
switchgrass	<i>Panicum virgatum</i>	PAVI2	UDPD	UDDU	UDPD	UUUU	UUUU
thickspike wheatgrass	<i>Elymus lanceolatus</i>	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD
threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP
threeawn	<i>Aristida</i> spp.	ARIS1	NNNN	NNNN	NNNN	NNNN	NNNN
Timber oatgrass (danthonia)	<i>Danthonia intermedia</i>	DAIN	DDDD	DDDD	DDDD	UUUU	UUUU
tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD
western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DDDD	DDDD	DDDD	DDDD	DDDD
FORBS							
alkali (pursh) seepweed	<i>Suaeda calceoliformis</i>	SUCA2	NNNN	NNNN	NNNN	NNNN	NNNN
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU
American vetch	<i>Vicia americana</i>	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP
arrowgrass	<i>Triglochin</i> spp.	TRIGL	T	T	T	T	T
biscuitroots	<i>Lomatium</i> spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD
bluebells	<i>Mertensia</i>	MERTE	DDDD	PPPP	DDDD	DDDD	DDDD
blue-eyed grass	<i>Sisyrinchium</i> spp.	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD
breadroot scurfpea	<i>Pediemelum esculentum</i>	PEES	DDDD	DDDD	DDDD	DDDD	DDDD
cattail, broad-leaf	<i>Typha latifolia</i>	TYLA	DDDD	UUUU	DDDD	UUUU	UUUU
cattail, narrow-leaf	<i>Typha angustifolia</i>	TYAN	DDDD	UUUU	DDDD	UUUU	UUUU
common comandra (toadflax)	<i>Comandra umbellata</i>	COUMP	UUUU	UUUU	UUUU	UUUU	UUUU
cutweed sagewort	<i>Artemisia ludoviciana</i>	ARLU	UUUU	UUUU	UUUU	UUUU	UUUU
deathcamas	<i>Zigadenus venenosus</i>	ZIVE	TTTT	TTTT	TTTT	TTTT	TTTT
dotted gayfeather	<i>Liatris punctata</i>	LIPU	UPPU	UPPU	UPPU	UPPU	UPPU
erigeron (fleabanes)	<i>Erigeron</i> spp.	ERIGE2	UUUU	UUUU	UUUU	UUUU	UUUU
erigonum (buckwheat)	<i>Eriogonum</i> spp.	ERIGO	UUUU	DDDD	UUUU	UUUU	UUUU
fringed sagewort	<i>Artemisia frigida</i>	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU
goldenrod	<i>Oligoneuron</i>	OLIGO3	UUUU	UUUU	UUUU	UUUU	UUUU
green sagewort	<i>Artemisia dracuncul</i>	ARDR4	UUUU	UUUU	UUUU	UUUU	UUUU
gromwell	<i>Buglossoides arvensis</i>	BUAR3	UUUU	UUUU	UUUU	UUUU	UUUU
groundsel	<i>Tephrosia</i>	TEPHR3	UUUU	UUUU	UUUU	UUUU	UUUU
hawksbeard	<i>Crepis acuminata</i>	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD
horsetails	<i>Equisetum</i> spp.	EQUI5	UUUU	UUUU	UUUU	UUUU	UUUU
iris	<i>Iris</i> spp.	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU
mountain thermopsis	<i>Thermopsis divaricarpa</i>	THDI4	UUUU	UUUU	UUUU	UUUU	UUUU
Nailworts	<i>Paronychia</i> spp.	PARON	UUUU	UUUU	UUUU	UUUU	UUUU
penstemons	<i>Penstemon</i> spp.	PENST	PPPP	PPPP	PPPP	PPPP	PPPP
prairie coneflower	<i>Ratibida columnifera</i>	RACO3	DDDD	PPPP	DDDD	PPPP	PPPP
prairie clovers	<i>Dalea</i> spp.	DALEA	UPPU	UPPU	UPPU	UPPU	UPPU
scurfpeas	<i>Psoraleum</i> spp.	PSORA2	NNNN	UUUU	NNNN	UUUU	UUUU
starwort	<i>Callitriche</i> spp.	CALL16	UUUU	UUUU	UUUU	UUUU	UUUU
stonecrop	<i>Sedum</i> spp.	SEDUM	UUUU	UUUU	UUUU	UUUU	UUUU
twogrooved milkvetch	<i>Astragalus bisulcatus</i>	ASBI2	T	T	T	T	T
violets	<i>Viola</i> spp.	VIOLA	DDDD	DDDD	DDDD	DDDD	DDDD
water hemlocks	<i>Cicuta</i> spp.	CICUT	T	T	T	T	T
western virgin'sbower	<i>Clematis occidentalis</i>	CLOC2	UUUU	DDDD	UUUU	DDDD	DDDD
western wallflower	<i>Erysimum capitatum</i>	ERICAC	DDDD	DDDD	DDDD	DDDD	DDDD
western yarrow	<i>Achillea lanulosa</i>	ACHIL	UUUU	UUUU	UUUU	UUUU	UUUU
wild onion	<i>Allium textile</i>	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD
TREES, SHRUBS & HALF-SHRUBS							
big sagebrush	<i>Artemisia tridentata</i>	ARTR2	UUUU	DDDD	UUUU	DDDD	DDDD
black greasewood	<i>Sarcobatus vermiculatus</i>	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD
green rabbitbrush	<i>Chrysothamnus viscidiflorus</i>	CHVI8	DDDD	DDDD	DDDD	DDDD	DDDD
plains cottonwood (sprouts)	<i>Populus deltoides</i>	PODEM	DDDD	DDDD	DDDD	DDDD	DDDD
rubber rabbitbrush	<i>Encameria nauseosa</i>	ERNA10	UUUU	DDDD	UUUU	DDDD	DDDD
silver sagebrush	<i>Artemisia cana</i>	ARCA5	DDDD	DDDD	DDDD	PPPP	PPPP
skunkbush sumac	<i>Rhus trilobata</i>	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD
western snowberry	<i>Symphoricarpos occidentalis</i>	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU
wildrose	<i>Rosa woodsii</i> var. <i>woodsii</i>	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD
willows	<i>Salix</i> L.	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU
winterfat	<i>Krascheninnikovia lanata</i>	KRLA2	PPPP	PPPP	PPPP	PPPP	PPPP
yucca	<i>Yucca glauca</i>	YUGL	DDDD	DDDD	DDDD	DDDD	DDDD

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (Lbs/acre)	Carrying Capacity* (AUM/ac)
Historic Climax Plant Community	4500-7000	3.0
Baltic rush/Cattail	2000-4000	2.5
Baltic rush/Smartweed	1200-2500	1.5

* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

Hydrology Functions

Climate is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B and C, with localized areas in hydrologic group D. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group and water table. Runoff will be high on this site since the soil may be saturated. (Refer to Part 630, NRCS National Engineering Handbook for detailed hydraulic information.)

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

Recreational Uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Wood Products

No appreciable wood products are present on the site.

Other Products

None noted.

Supporting Information

Associated Sites

Subirrigated	061XY174WY
Overflow	061XY130WY
Lowland	061XY128WY

Similar Sites

(058BY278WY) – Wetland 15-17” Northern Plains P.Z. has lower production.

Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417		1971-1994	WY	Crook & others
Ocular estimates		1990-1999	WY	Crook & others

State Correlation

This site occurs entirely within Wyoming.

Type Locality

Field Offices

Newcastle, Sundance

Relationship to Other Established Classifications

Other References

Site Description Approval

State Range Management Specialist

Date

Ecological Reference Worksheet

Author(s)/participant(s): _____
 Contact for lead author: _____ Reference site used? Yes/No
 Date: 4/05 MLRA: 61 Ecological Site: R061XY178WY Wetland (WL) 15-19"BL

_____ This *must* be verified based on soils and climate (see Ecological Site Description). Current plant community *cannot* be used to identify the ecological site.

<p>Indicators. For each indicator, describe the potential for the site. Where possible, (1) use numbers, (2) include expected range of values for above- and below-average years for each community within the reference state, when appropriate & (3) cite data. Continue descriptions on separate sheet.</p>
<p>1. Number and extent of rills: Rills should not be present</p>
<p>2. Presence of water flow patterns: Barely observable</p>
<p>3. Number and height of erosional pedestals or terracettes: Essentially non-existent</p>
<p>4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are <i>not</i> bare ground): Bare ground is less than 5%</p>
<p>5. Number of gullies and erosion associated with gullies: Active gullies should not be present</p>
<p>6. Extent of wind scoured, blowouts and/or depositional areas: None</p>
<p>7. Amount of litter movement (describe size and distance expected to travel): Little to no plant litter movement. Plant litter remains in place and is not moved by erosional forces.</p>
<p>8. Soil surface (top few mm) resistance to erosion (stability values are averages – most sites will show a range of values for both plant canopy and interspaces, if different): Plant cover and litter is at 95% or greater of soil surface and maintains soil surface integrity. Soil Stability class is anticipated to be 5 or greater.</p>
<p>9. Soil surface structure and SOM content (include type and strength of structure, and A-horizon color and thickness for both plant canopy and interspaces, if different): Use Soil Series description for depth and color of A-horizon</p>
<p>10. Effect of plant community composition (relative proportion of different functional groups) & spatial distribution on infiltration & runoff: Grass canopy and basal cover should reduce raindrop impact and slow overland flow providing increased time for infiltration to occur. Healthy deep rooted native grasses enhance infiltration and reduce runoff. Infiltration varies from moderate to low and runoff is high since the soil is usually saturated.</p>
<p>11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): No compaction layer or soil surface crusting should be present.</p>
<p>12. Functional/Structural Groups (list in order of descending dominance by above-ground weight using symbols: >>>, >, = to indicate much greater than, greater than, and equal to): Tall Grasses and Grasslike > Mid stature Grasses/Grasslike > Forbs > Shrubs/Trees</p>
<p>13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Very Low</p>
<p>14. Average percent litter cover and depth : Average litter cover is 50-55% with depths of 0.75 to 1.5 inches</p>
<p>15. Expected annual production (this is all above-ground production, not just forage production): 6000 lbs/ac</p>
<p>16. Potential invasive (including noxious) species (native and non-native). List species which characterize degraded states and which, after a threshold is crossed, “can, and often do, continue to increase regardless of the management of the site and may eventually dominate the site”: Spike sedge, Baltic rush, Smartweed, American licorice, and Species found on Noxious Weed List</p>
<p>17. Perennial plant reproductive capability: All species are capable of reproducing</p>