

# United States Department of Agriculture Natural Resources Conservation Service

## Ecological Site Description

**Site Type:** Rangeland

**Site Name:** Impervious Clay (IC), 7-9" P.Z., Green River and Great Divide Basins

**Site ID:** R034AY118WY

**Major Land Resource Area:** 34A-Cool Central Desertic Basins and Plateaus

### Physiographic Features

This site will usually occur in a lowland position, on flat to gently sloping land, but can occur in all positions.

**Landform:** alluvial fans & stream terraces

**Aspect:** N/A

	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	6000	7200
<b>Slope (percent):</b>	0	40
<b>Water Table Depth (inches):</b>	none within 60	
<b>Flooding:</b>		
<b>Frequency:</b>	none	none
<b>Duration:</b>	none	none
<b>Ponding:</b>		
<b>Depth (inches):</b>	0	0
<b>Frequency:</b>	none	none
<b>Duration:</b>	none	none
<b>Runoff Class:</b>	negligible	high

### Climatic Features

Annual precipitation ranges from 7-9 inches per year. Wide fluctuations may occur in yearly precipitation and result in more dry years than those with more than normal precipitation.

Temperatures show a wide range between summer and winter and between daily maximums and minimums. This is predominantly due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of native cool season plants begins about April 15 and continues to about July 15. Some green up of cool season plants may occur in late September if moisture is available.

The following information is from the "Green River" climate station:

	<u>Minimum</u>	<u>Maximum</u>	<u>5 yrs. out of 10 between</u>
Frost-free period (days):	68	121	June 2 – September 5
Freeze-free period (days):	97	132	May 23 – September 19

Annual Precipitation (inches): <5.32 >9.34 (2 years in 10)

Average annual precipitation: 7.78 inches

Average annual air temperature: 41.8°F (25.6°F Avg. Min. to 58.1°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/cgibin/state.pl?state=wy> website. Other climate stations representative of this precipitation zone include “Bitter Creek”, “Farson”, “Rock Springs FAA AP”, and “Wamsutter” in Sweetwater County; “Church Buttes Gas PLT”, and “Mountain View” in Uinta County; “Fontenelle”, “La Barge”, and “Sage 4 NNW” in Lincoln County; and “Big Piney” in Sublette County.

### Influencing Water Features

<u>Wetland Description:</u>	<u>System</u>	<u>Subsystem</u>	<u>Class</u>	<u>Sub-class</u>
None	None	None	None	None

Stream Type: None

### Representative Soil Features

The soils of this site are moderately deep (greater than 20” to bedrock) to very deep, well drained soils formed in alluvium. The topsoil, except for thin ineffectual layers, will be heavy clays and/or soils that develop large cracks when dry and are very sticky when wet. These soils are not high in salinity and/or alkalinity but do have high concentrations of exchangeable sodium throughout the profile.

Major Soil Series correlated to this site include: The Kissick and Elkol series.

Parent Material Kind: alluvium

Parent Material Origin: sedimentary rock

Surface Texture: clay, silty clay

Surface Texture Modifier: none

Subsurface Texture Group: clay, silty clay

Surface Fragments ≤ 3” (% Cover): 0

Surface Fragments > 3” (%Cover): 0

Subsurface Fragments ≤ 3” (% Volume): 0

Subsurface Fragments > 3” (% Volume): 0

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	well drained	well drained
Permeability Class:	slow	very slow
Depth (inches):	15	>60
Electrical Conductivity (mmhos/cm) ≤20”:	0	8
Sodium Absorption Ratio ≤20”:	0	5
Soil Reaction (1:1 Water) ≤20”:	6.6	8.4
Soil Reaction (0.1M CaCl2) ≤20”:	NA	NA
Available Water Capacity (inches) ≤30”:	3	4.5
Calcium Carbonate Equivalent (percent) ≤20”:	0	15


## Plant Communities

### Ecological Dynamics of the Site:

As this site deteriorates from improper grazing management, species such as birdfoot sage and unpalatable forbs will increase. Indian ricegrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



BMA – Brush Management (all methods)  
 BMC – Brush Management (chemical)  
 BMF – Brush Management (fire)  
 BMM – Brush Management (mechanical)  
 CSP – Chemical Seedbed Preparation  
 CSLG – Continuous Season-long Grazing  
 DR – Drainage  
 CSG – Continuous Spring Grazing  
 HB – Heavy Browse  
 HCSLG – Heavy Continuous Season-long Grazing  
 HI – Heavy Inundation  
 LPG – Long-term Prescribed Grazing  
 MT – Mechanical Treatment (chiseling, ripping, pitting)

NF – No Fire  
 NS – Natural Succession  
 NWC – Noxious Weed Control  
 NWI – Noxious Weed Invasion  
 NU – Nonuse  
 P&C – Plow & Crop (including hay)  
 PG – Prescribed Grazing  
 RPT – Re-plant Trees  
 RS – Re-seed  
 SGD – Severe Ground Disturbance  
 SHC – Severe Hoof Compaction  
 WD – Wildlife Damage (Beaver)  
 WF - Wildfire

**Plant Community Composition and Group Annual Production**  
**Reference Plant Community (HCPC)**

COMMON NAME/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Annual Production (Normal Year)		
			Group	lbs./acre	% Comp.
			<b>Total: 350</b>		
<b>GRASSES AND GRASS-LIKES</b>					
<b>GRASSES/GRASSLIKES</b>					
western wheatgrass	Pascopyrum smithii	PASM	1	88 - 158	25 - 45
bottlebrush squirreltail	Elymus elymoides	ELEL5	2	35 - 70	10 - 20
Indian ricegrass	Achnatherum hymenoides	ACHY	3	35 - 70	10 - 20
<b>MISC. GRASSES/GRASSLIKES</b>			<b>4</b>	<b>4 - 18</b>	<b>1 - 5</b>
Sandberg bluegrass	Poa secunda	POSE	4	0 - 18	0 - 5
other perennial grasses (native)		2GP	4	0 - 18	0 - 5
<b>FORBS</b>			<b>5</b>	<b>18 - 35</b>	<b>5 - 10</b>
asters	Eucephalus spp.	EUCEP2	5	0 - 18	0 - 5
biscuitroot	Lomatium spp.	LOMAT	5	0 - 18	0 - 5
deathcamas	Zigadenus spp.	ZIGAD	5	0 - 18	0 - 5
fleabane	Erigeron spp.	ERIGE2	5	0 - 18	0 - 5
milkvetches	Astragalus spp.	ASTRA	5	0 - 18	0 - 5
onion	Allium textile	ALTE	5	0 - 18	0 - 5
phlox	Phlox spp.	PHLOX	5	0 - 18	0 - 5
primrose	Oenothera caespitosa	OECA10	5	0 - 18	0 - 5
woody aster	Xylorhiza spp.	XYLOR	5	0 - 18	0 - 5
pussytoes	Antennaria rosea	ANRO2	5	0 - 18	0 - 5
other perennial forbs (native)		2FP	5	0 - 18	0 - 5
<b>TREES/SHRUBS</b>					
birdfoot sagebrush	Artemisia pedatifida	ARPE6	6	88 - 140	25 - 40
<b>MISC. SHRUBS</b>			<b>7</b>	<b>18 - 35</b>	<b>5 - 10</b>
Gardners saltbush	Atriplex gardneri	ATGA	7	0 - 18	0 - 5
winterfat	Krascheninnikovia lanata	KRAL2	7	0 - 18	0 - 5

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

### Plant Community Narratives

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

#### Birdfoot Sage/Rhizomatous Wheatgrass Plant Community (HCPC)

The interpretive plant community for this site is the Historic Climax Plant Community. Potential vegetation is estimated at 50% grasses or grass-like plants, 5% forbs and 45% woody plants. The major grasses include western wheatgrass, bottlebrush squirreltail, Indian ricegrass, and Sandberg bluegrass. Birdfoot sagebrush is the major woody plant. Other woody plants that may occur include Gardner’s saltbush and winterfat.

A typical plant composition for this state consists of western wheatgrass 25-45%, bottlebrush squirreltail 10-20%, Indian ricegrass 10-20%, up to 5% Sandberg bluegrass, perennial forbs 1-5%, birdfoot sagebrush 25-40%, and 5-10% other woody species. Ground cover, by ocular estimate, varies from 30-45%.

The total annual production (air-dry weight) of this state is about 350 pounds per acre, but it can range from about 250 lbs./acre in unfavorable years to about 500 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY0401

Growth curve name: 7-9GR, UPLAND SITES

Growth curve description: ALL UPLAND SITES

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	35	40	10	0	5	0	0	0

(Monthly percentages of total annual growth)

This state is extremely stable and well adapted to the Cool Central Desertic Basins and Plateaus climatic conditions. The diversity in plant species allows for high drought resistance. This is a sustainable plant community (site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Heavy Continuous Season-long Grazing or Severe Hoof Compaction will convert the plant community to the *Heavy Birdfoot Sage State*.

### Rhizomatous Wheatgrass/Birdfoot Sage Plant Community

This plant community is the result of mechanical treatment after frequent and severe grazing. Rhizomatous wheatgrass and bottlebrush squirreltail dominate. Birdfoot sagebrush usually comprises 10-20% of annual production.

The total annual production (air-dry weight) of this state is about 200 pounds per acre, but it can range from about 100 lbs./acre in unfavorable years to about 400 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY0401

Growth curve name: 7-9GR, UPLAND SITES

Growth curve description: ALL UPLAND SITES

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	35	40	10	0	5	0	0	0

(Monthly percentages of total annual growth)

The soil is not protected and erosion will increase if management is not changed. The biotic integrity may be reduced due to low vegetative production. The watershed is functioning at risk.

Transitional pathways leading to other plant communities are as follows:

- Natural Succession will return this state to near *Historic Climax Plant Community (Birdfoot Sage/Rhizomatous Wheatgrass State)*.

### Heavy Birdfoot Sage Plant Community

This plant community is a result of heavy continuous season-long grazing. Severe hoof compaction typically occurs due to fine soil textures. Birdfoot sage increases to 60-80% of the annual production. Cool season bunchgrasses decrease while rhizomatous wheatgrass prevails. Annual forbs increase.

The total annual production (air-dry weight) of this state is about 100 pounds per acre, but it can range from about 50 lbs./acre in unfavorable years to about 250 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number: WY0401

Growth curve name: 7-9GR, UPLAND SITES

Growth curve description: ALL UPLAND SITES

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	35	40	10	0	5	0	0	0

(Monthly percentages of total annual growth)

This state is unstable and vulnerable to excessive erosion. The biotic integrity of this plant community is at risk or non-functioning. The watershed is usually at risk or non-functioning as bare ground increases.

Transitional pathways leading to other plant communities are as follows:

- Mechanical Treatment followed by deferment for 1 to 2 years as part of a Prescribed Grazing plan will convert this plant community to a *Rhizomatous Wheatgrass/Birdfoot Sage State*.

## Ecological Site Interpretations

### Animal Community – Wildlife Interpretations

**Birdfoot Sage/Rhizomatous Wheatgrass Plant Community (HCPC):** Suitable thermal and escape cover for mule deer may be limited due to the low height of woody plants. However, sagebrush provides important winter forage for mule deer and antelope. Year-round habitat is provided for sage grouse and many other sagebrush obligate species such as the sage sparrow, sage thrasher, pygmy rabbit, sagebrush vole, horned lizard, and pronghorn antelope. Open spaces in the sagebrush canopy are potential sage grouse lek locations.

**Rhizomatous Wheatgrass/Birdfoot Sage Plant Community:** This plant community has a low level of diversity. Due to the dominance of grasses, feed for browsing animals is limited. Areas of bare ground may provide leks for sage grouse.

**Heavy Birdfoot Sage Plant Community:** This plant community may be beneficial for the same wildlife that would use the Historic Climax Plant Community. However, the plant community composition is less diverse, and thus, less apt to meet the seasonal needs of these animals.

Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA34A, 7-9 inch Green River & Great Divide Basins

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope	Elk
<b>GRASSES/GRASSLIKES</b>								
Alkali bluegrass	<i>Poa juncea</i> (syn. <i>P. secunda</i> )	POJU (POSE)	DDDD	PPPP	DDDD	PPPP	PPPP	DDDD
Alkali muhly	<i>Muhlenbergia asperifolia</i>	MUAS	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Alkali sacaton	<i>Sporobolus airoides</i>	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Baltic rush	<i>Juncus balticus</i>	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD
Basin wildrye	<i>Leymus cinereus</i>	LEC4	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP
Bluebunch wheatgrass	<i>Pseudoroegneria spicata</i>	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP
Bluejoint reedgrass	<i>Calamagrostis canadensis</i>	CACAM	PPPP	DDDD	PPPP	DDDD	UUUU	PPPP
Bottlebrush squirreltail	<i>Elymus elymoides</i>	ELELE	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Canada wildrye	<i>Elymus canadensis</i>	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP
Canby bluegrass	<i>Poa canbyi</i> (syn. <i>P. secunda</i> )	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Indian ricegrass	<i>Achnatherum hymenoides</i>	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Inland saltgrass	<i>Distichlis spicata</i>	DISP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Inland sedge	<i>Carex interior</i>	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD
James' galleta	<i>Pleuraphis jamesii</i>	PLJA	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD
Letterman needlegrass	<i>Achnatherum lettermanii</i>	ACLE9	PPPP	PPPP	DDDD	DDDD	DDDD	PPPP
Mat muhly	<i>Muhlenbergia richardsonis</i>	MURI	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	<i>Carex nebrascensis</i>	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP
Needleandthread	<i>Hesperostipa comata</i>	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Needleleaf sedge	<i>Carex duriuscula</i>	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Northern reedgrass	<i>Calamagrostis stricta</i> ssp. <i>inexpansa</i>	CASTI3	PPPP	DDDD	PPPP	DDDD	UUUU	PPPP
Nuttall's alkalgrass	<i>Puccinellia nuttalliana</i>	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Plains reedgrass	<i>Calamagrostis montanensis</i>	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Prairie junegrass	<i>Koeleria macrantha</i>	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Reed canarygrass	<i>Phalaris arundinacea</i>	PHAR3	PPPP	UUUU	UUUU	UUUU	UUUU	PPPP
Saline wildrye	<i>Leymus salinus</i>	LESA4	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Sandberg bluegrass	<i>Poa secunda</i>	POSE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Sand dropseed	<i>Sporobolus cryptandrus</i>	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU	DDDD
Slender wheatgrass	<i>Elymus trachycaulis</i>	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD	PPPP
Tall mangrass	<i>Glyceria elata</i> (syn. <i>G. striata</i> )	GLEL (GLST)	DDDD	UUUU	DDDD	UUUU	UUUU	DDDD
Thickspike wheatgrass	<i>Elymus lanceolatus</i> ssp. <i>lanceolatus</i>	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Threadleaf sedge	<i>Carex filifolia</i>	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP	DDDD
Threewaves	<i>Aristida</i> spp.	ARIS	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Tufted hairgrass	<i>Deschampsia caespitosa</i>	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD	PPPP
Western wheatgrass	<i>Pascopyrum smithii</i>	PASM	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
<b>FORBS</b>								
American licorice	<i>Glycyrrhiza lepidota</i>	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Arrowgrass	<i>Triglochin</i> spp.	TRIGL	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Asters	<i>Eucephalus</i> spp.	EUCEP2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Biscuitroot	<i>Lomatium</i> spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD
Blue-eyed grass	<i>Sisyrinchium</i> spp.	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD	DDDD
Buckwheats	<i>Eriogonum</i> spp.	ERIOG	UUUU	DDDD	UUUU	UUUU	UUUU	UUUU
Buttercup	<i>Ranunculus</i> spp.	RANUN	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Clovers	<i>Trifolium</i> spp.	TRIFO	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Deathcamas	<i>Zigadenus</i> spp.	ZIGAD	TTTT	ZIGAD	TTTT	TTTT	TTTT	TTTT
Docks	<i>Rumex</i> spp.	RUMEX	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Elephanthead lousewort	<i>Pedicularis groenlandica</i>	PEGR2	UUUU	DDDD	UUUU	DDDD	UUUU	UUUU
Flax	<i>Linum</i> spp.	LINUM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Fleabanes	<i>Erigeron</i> spp.	ERIGE2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Fringed sagewort	<i>Artemisia frigida</i>	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Goldenpea	<i>Thermopsis</i> spp.	THERM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Goldenweed	<i>Stenotus acaulis</i>	STAC	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Gromwell	<i>Buglossoides arvensis</i>	BUAR3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Groundsel	<i>Tephrosia</i> spp.	TEPHR3	TTTT	UUUU	TTTT	UUUU	UUUU	TTTT
Hawksbeard	<i>Crepis acuminata</i>	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD	UUUU
Horsetails	<i>Equisetum</i> spp.	EQUIS	UUUU	UUUU	TTTT	UUUU	UUUU	UUUU
Iris	<i>Iris</i> spp.	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Milkvetch (locoweed)	<i>Astragalus</i> spp.	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Miners candle	<i>Cryptantha virgata</i>	CRV14	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Paintbrush	<i>Castilleja</i> spp.	CAST	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Penstemons	<i>Penstemon</i> spp.	PENST	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Phlox	<i>Phlox</i> spp.	PHLOX	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Povertyweed	<i>Monolepis</i> spp.	MONOL	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Primrose	<i>Oenothera</i>	OENOT	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Princesplume	<i>Stanleya</i> spp.	STANL	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Pussytoes	<i>Antennaria</i> spp.	ANTEN	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Sagebrush gilia	<i>Leptodactylon pungens</i>	LEPU	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Sandwort	<i>Arenaria</i> spp.	ARENA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Scarlet globemallow	<i>Sphaeralcea coccinea</i>	SPCO	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Scurfpeas	<i>Psoralea</i> spp.	PSORA2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Stoncrop	<i>Sedum</i> spp.	SEDUM	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Tansy	<i>Tanacetum</i> spp.	TANAC	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Toadflax	<i>Comandra umbellata</i>	COUMP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Violets	<i>Viola</i> spp.	VIOLA	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Water hemlock	<i>Cicuta</i> spp.	CICUT	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
Waterleaf	<i>Hydrophyllum</i> spp.	HYDRO4	DDDD	DDDD	DDDD	PPPP	DDDD	DDDD
Western yarrow	<i>Achillea millefolium</i>	ACHMIO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Wild onion	<i>Allium textile</i>	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Woody aster	<i>Xylorhiza</i> spp.	XYLOR	TTTT	TTTT	TTTT	TTTT	TTTT	TTTT
<b>TREES, SHRUBS &amp; HALF-SHRUBS</b>								
Antelope bitterbrush	<i>Purshia tridentata</i>	PUTR2	PPPP	PPPP	DDDD	PPPP	PPPP	PPPP
Big sagebrush	<i>Artemisia tridentata</i>	ARTR2	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD
Birdfoot sagebrush	<i>Artemisia pedatifida</i>	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Bud sagewort	<i>Artemisia spinescens</i>	ARSP5	PPPP	PPPP	DDDD	PPPP	PPPP	PPPP
Buffalobery	<i>Shepherdia</i> spp.	SHEPH	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Cottonwood (sprouts only)	<i>Populus angustifolia</i>	POAN3	PPPP	PPPP	PPPP	PPPP	UUUU	PPPP
Currant	<i>Ribes</i> spp.	RIBES	DDDD	DDDD	DDDD	DDDD	UUUU	DDDD
Early (alkali) sagebrush	<i>Artemisia arbuscula</i> ssp. <i>longiloba</i>	ARARL	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Fourwing saltbush	<i>Atriplex canescens</i>	ATCA2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Gardners saltbush	<i>Atriplex gardneri</i>	ATGA	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP
Greasewood (toxic in large amounts)	<i>Sarcobatus vermiculatus</i>	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD
Greenmolly summercypress	<i>Kochia americana</i>	KOMA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Green rabbitbrush	<i>Chrysothamnus viscidiflorus</i>	CHV18	DDDD	DDDD	UUUU	PPPP	PPPP	DDDD
Hawhorn	<i>Crataegus</i> spp.	CRATA	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Junipers	<i>Juniperus scopulorum</i>	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU	UUUU
Limber pine	<i>Pinus flexilis</i>	PIFL2	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Low sagebrush	<i>Artemisia arbuscula</i>	ARAR8	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD
Rubber rabbitbrush	<i>Ericameria nauseosa</i>	ERNA10	UUUU	DDDD	UUUU	DDDD	PPPP	UUUU
Shadscale	<i>Atriplex confertifolia</i>	ATCO	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Shrubby cinquefoil	<i>Dasiphora floribunda</i>	DAFL3	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Silver sagebrush	<i>Artemisia cana</i>	ARCA13	DDDD	DDDD	DDDD	PPPP	PPPP	DDDD
Skunkbush sumac	<i>Rhus trilobata</i>	RHTR	DDDD	DDDD	UUUU	DDDD	DDDD	DDDD
Spineless horsebrush	<i>Tetradymia canescens</i>	TECA2	UUUU	TTTT	UUUU	UUUU	UUUU	UUUU
Spiny hopsage	<i>Grayia spinesa</i>	GRSP	UUUU	UUUU	UUUU	UUUU	UUUU	UUUU
Spiny horsebrush	<i>Tetradymia spinosa</i>	TESP2	UUUU	DDDD	UUUU	UUUU	DDDD	UUUU
Wildrose	<i>Rosa woodsii</i> var. <i>woodsii</i>	ROWOW	DDDD	DDDD	DDDD	DDDD	DDDD	DDDD
Willows	<i>Salix</i> spp.	SALIX	DDDD	DDDD	DDDD	PPPP	UUUU	DDDD
Winterfat	<i>Krascheninnikovia lanata</i>	KRAL2	PPPP	PPPP	PPPP	PPPP	PPPP	PPPP

N = not used; U = undesirable; D = desirable; P = preferred; T = toxic

## Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

Plant Community	Production (lb./ac)	Carrying Capacity* (AUM/ac)
Birdfoot Sage/Rhizomatous Wheatgrass (HCPC)	250-500	.08
Rhizomatous Wheatgrass/Birdfoot Sage	100-400	.04
Heavy Birdfoot Sage	50-250	.02

\* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

## Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group D. Infiltration ranges from slow to very slow. Runoff potential for this site varies from high to very high depending on ground cover (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns should be barely distinguishable if at all present. Pedestals are only slightly present in association with bunchgrasses. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogammic crusts are present, but only cover 1-2% of the soil surface.

## Recreational Uses

This site provides limited hunting opportunities.

## Wood Products

No appreciable wood products are present on the site.

## Other Products

None noted.

## Supporting Information

### Associated Sites

Clayey

R034AY104WY

Saline Upland R034AY144WY

### Similar Sites

R034AY110WY – Dense Clay (DC) 7-9GR has higher production and does not have alkaline/saline properties.

### Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel were also used. Those involved in developing this site include: Bill Christensen, Range Management Specialist, NRCS; Karen Clause, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist, NRCS. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

### Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417	50	1966-1985	WY	Sweetwater & others

### State Correlation

### Type Locality

### Field Offices

Baggs, Cokeville, Rock Springs/Farson, Lyman, Pinedale, Saratoga

### Relationship to Other Established Classifications

### Other References

### Site Description Approval

\_\_\_\_\_  
State Range Management Specialist

\_\_\_\_\_  
Date