

## **United States Department of Agriculture Natural Resources Conservation Service**

### **Ecological Site Description**

**Site Type:** Rangeland

**Site Name:** Shale (Sh) 15-17” Northern Plains Precipitation Zone,

**Site ID:** 058BY254WY

**Major Land Resource Area:** 58B – Northern Rolling High Plains

### **Physiographic Features**

This site occurs on moderate to steep slopes and ridge tops.

**Landform:** Hill sides, ridges & escarpments      **Aspect:** N/A

	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	3400	4600
<b>Slope (percent):</b>	0	60
<b>Water Table Depth (inches):</b>	None within 60 inches	
<b>Flooding:</b>		
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Ponding:</b>		
<b>Depth (inches):</b>	0	0
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Runoff Class:</b>	negligible	high

### **Climatic features**

Annual precipitation ranges from 15-17 inches per year. Wide fluctuations may occur in yearly precipitation and result in more drought years than those with more than normal precipitation. Temperatures show a wide range between summer and winter and between daily maximums and minimums, due to the high elevation and dry air, which permits rapid incoming and outgoing radiation. Cold air outbreaks from Canada in winter move rapidly from northwest to southeast and account for extreme minimum temperatures. Chinook winds may occur in winter and bring rapid rises in temperature. Extreme storms may occur during the winter, but most severely affect ranch operations during late winter and spring.

Wind speed averages about 8 mph, ranging from 10 mph during the spring to 7 mph during late summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 75 mph.

Growth of native cool-season plants begins about April 1 and continues to about July 1. Native warm-season plants begin growth about May 15 and continue to about August 15. Green up of cool season plants may occur in September and October of most years.

The following information is from the “Echeta 2 NW” climate station:

Site Type: Rangeland  
 MLRA: 58B – Northern Rolling High Plains

**Shale 15-17” P.Z.  
 R058BY254WY**

Frost-free period (32 °F): 70-142 days; (5 yrs. out of 10, these days will occur between June 7 – September 16)

Freeze-free period (28 °F): 106-154 days; (5 yrs. out of 10, these days will occur between May 14 – September 23)

Mean annual precipitation: 15.82 inches

Mean annual air temperature: 45.2 °F (30.0°F Avg. Min. - 60.4°F Avg. Max.)

For detailed information visit the Natural Resources Conservation Service National Water and Climate Center at <http://www.wcc.nrcs.usda.gov/> website. Other climate station(s) representative of this precipitation zone include: “Recluse 14 NNW”.

## Influencing Water Features

<b>Wetland Description:</b>	<b><u>System</u></b>	<b><u>Subsystem</u></b>	<b><u>Class</u></b>	<b><u>Sub-class</u></b>
None	None	None	None	None

**Stream Type:** None

## Representative Soil Features

The soils of this site are very shallow (less than 10 inches to bedrock) well-drained soils formed from residuum. These soils have rapid to slow permeability and can be of any texture. This site usually occurs on steep slopes. The soil is generally less than 10 inches deep with many outcrops of clay shale bedrock. These clay shales are usually saline or alkaline in various degrees, and normally produce some halophytic species. Layers of the soil most influential to the plant community vary from 3 to 6 inches thick.

Major Soil Series correlated to this site include: Samsil, Lismas and Fairburn.

Other Soil Series in MLRA 58B correlated to this site include: Badlands, Rock Outcrop

**Parent Material Kind:** residuum

**Parent Material Origin:** shale

**Surface Texture:** clay loam, loam, silt loam, silty clay loam, clay

**Surface Texture Modifier:** none

**Subsurface Texture Group:** clay loam

**Surface Fragments ≤ 3” (% Cover):** 0 to 10

**Surface Fragments > 3” (%Cover):** 0 to 10

**Subsurface Fragments ≤ 3” (% Volume):** 5 to 20

**Subsurface Fragments > 3” (% Volume):** 0

	<b><u>Minimum</u></b>	<b><u>Maximum</u></b>
<b>Drainage Class:</b>	well	well
<b>Permeability Class:</b>	slow	moderate
<b>Depth (inches):</b>	0	10
<b>Electrical Conductivity (mmhos/cm) ≤20”:</b>	0	4
<b>Sodium Absorption Ratio ≤20”:</b>	0	5
<b>Soil Reaction (1:1 Water) ≤20”:</b>	6.6	8.4
<b>Soil Reaction (0.1M CaCl2) ≤20”:</b>	NA	NA
<b>Available Water Capacity (inches) ≤30”:</b>	1	2
<b>Calcium Carbonate Equivalent (percent) ≤20”:</b>	0	5


## **Plant Communities**

### **Ecological Dynamics of the Site:**

As this site deteriorates, species such as juniper will increase and species such as broom snakeweed, plains pricklypear and annual forbs will invade. Cool season grasses such as bluebunch wheatgrass, little bluestem, and western wheatgrass will decrease in frequency and production.

The Historic Climax Plant Community (description follows the plant community diagram) has been determined by study of rangeland relic areas, or areas protected from excessive disturbance. Trends in plant communities going from heavily grazed areas to lightly grazed areas, seasonal use pastures, and historical accounts have also been used.

The following is a State and Transition Model Diagram that illustrates the common plant communities (states) that can occur on the site and the transitions between these communities. The ecological processes will be discussed in more detail in the plant community narratives following the diagram.



**BM** - Brush Management (fire, chemical, mechanical)

**Freq. & Severe Grazing** - Frequent and Severe Utilization of the Cool-season Mid-grasses during the Growing Season

**GLMT** - Grazing Land Mechanical Treatment

**LTPG** - Long-term Prescribed Grazing

**MCSLG** - Moderate, Continuous Season-long Grazing

**NU, NF** - No Use and No Fire

**PG** - Prescribed Grazing (proper stocking rates with adequate recovery periods during the growing season)

**VLTPG** - Very Long-term Prescribed Grazing (could possibly take generations)

**Na** - found adjacent to a saline site

PLANT COMMUNITY DYNAMICS  
REFERENCE PLANT COMMUNITY

COMMON NAME/ GROUP NAME	SCIENTIFIC NAME	SCIENTIFIC SYMBOL	Grp	Allowable Annual Production			% Comp (MAX.)
				lbs./acre			
				below normal 300	normal 500	above normal 700	
<b>GRASSES/GRASSLIKES</b>							
<b>RHIZOMATOUS WHEATGRASSES:</b>							
thickspike wheatgrass	Elymus lanceolatus	ELLAL	1	75	125	175	25%
western wheatgrass	Pascopyrum smithii	PASM	1	75	125	175	25%
<b>OTHER GRASSES</b>							
alkali sacaton	Sporobolus airoides	SPAI	2	15	25	35	5%
bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6	3	30	50	70	10%
green needlegrass	Nassella viridula	NAV14	4	30	50	70	10%
Indian ricegrass	Achnatherum hymenoides	ACHY	5	30	50	70	10%
plains reedgrass	Calamagrostis montanensis	CAMO	6	30	50	70	10%
little bluestem	Schizachyrium scoparium	SCSC	7	30	50	70	10%
threadleaf sedge	Carex filifolia	CAFI	8	30	50	70	10%
<b>MISCELLANEOUS GRASSES/GRASSLIKES*</b>							
Canby bluegrass	Poa canbyi (syn. to Poa secunda)	POCA (POSE)	9	15	25	35	5%
sideoats grama	Bouteloua curtipendula	BOCU	9	15	25	35	5%
prairie junegrass	Koeleria macrantha	KOMA	9	15	25	35	5%
Sandberg bluegrass	Poa secunda	POSE	9	15	25	35	5%
inland saltgrass	Distichlis spicata	DISP	9	15	25	35	5%
<b>FORBS</b>							
<b>MISCELLANEOUS FORBS*</b>							
American vetch	Vicia americana	VIAM	10	15	25	35	5%
prairie coneflower	Ratibida columnifera	RACO3	10	15	25	35	5%
asters	Asters	ASTER	10	15	25	35	5%
biscuitroots	Lomatium spp.	LOMAT	10	15	25	35	5%
breadroot scurfpea	Pedimelum esculentum	PEES	10	15	25	35	5%
western yarrow	Achillea lanulosa	ACHIL	10	15	25	35	5%
rosy pussytoes	Antennaria rosea	ANRO2	10	15	25	35	5%
milkvetches	Astragalus	ASTRA	10	15	25	35	5%
stemless goldenweed	Haplopappus acaulis	HAAC	10	15	25	35	5%
sulphur flower buckwheat	Eriogonum umbellatum	ERUM	10	15	25	35	5%
scarlet gaura	Gaura coccinea	GACO5	10	15	25	35	5%
prairie thermopsis	Thermopsis rhombifolia	THRHA	10	15	25	35	5%
purple prairie clover	Dalea purpurea	DAPU5	10	15	25	35	5%
white prairie clover	Dalea candida	DACA7	10	15	25	35	5%
bluebells	Mertensia	MERTE	10	15	25	35	5%
wild onion	Allium textile	ALTE	10	15	25	35	5%
hawksbeard	Crepis acuminata	CRAC2	10	15	25	35	5%
<b>TREES, SHRUBS &amp; HALF-SHRUBS*</b>							
winterfat	Krascheninnikovia lanata	KRLA2	11	15	25	35	5%
junipers	Juniperus scoparium	JUSC2	12	30	50	70	10%
ponderosa pine	Pinus ponderosa	PIPO	13	15	25	35	5%
Gardners saltbush	Atriplex gardnerii	ATGA	14	30	50	70	10%
birdfoot sage	Artemisia pedatifida	ARPE6	15	15	25	35	5%

\* Common native perennials are listed. Other native perennials may also be counted but no species in the group may be counted for more than 5%.

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors.

**Plant Community Narratives**

Following are the narratives for each of the described plant communities. These plant communities may not represent every possibility, but they probably are the most prevalent and repeatable plant communities. The plant composition tables shown above have been developed from the best available knowledge at the time of this revision. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. None of these plant communities should necessarily be thought of as “Desired Plant Communities”. According to the USDA NRCS National Range and Pasture Handbook, Desired Plant Communities (DPC’s) will be determined by the decision-makers and will meet minimum quality criteria established by the NRCS. The main purpose for including any description of a plant community here is to capture the current knowledge and experience at the time of this revision.

**Western Wheatgrass, Little bluestem Plant Community**

The interpretive plant community for this site is the Historic Climax Plant Community. This state evolved with grazing by large herbivores and is suited for grazing by domestic livestock. Potential vegetation is about 85% grasses or grass-like plants, 5% forbs, and 10% woody plants. The state is dominated by cool season midgrasses. The major grasses include rhizomatous wheatgrasses, bluebunch wheatgrass, green needlegrass and little bluestem. Other grasses occurring on the state include alkali sacaton, inland saltgrass, Indian ricegrass, Canby and Sandberg bluegrass, and plains reedgrass. Birdfoot sagebrush, winterfat and junipers are conspicuous elements of this state.

The total annual production (air-dry weight) of this state is about 500 pounds per acre, but it can range from about 300 lbs./acre in unfavorable years to about 700 lbs./acre in above average years.

The following is the growth curve of this plant community expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	25	40	10	5	5	5	0	0

(Monthly percentages of total annual growth)

The state is fragile and adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for some drought resistance. This is a sustainable plant community, but is difficult to reestablish when damaged. (Site/soil stability, watershed function, and biologic integrity).

Transitions or pathways leading to other plant communities are as follows:

- Moderate, Continuous Season-Long grazing will convert this plant community to the *Juniper/Inland saltgrass Vegetation State*.
- Frequent and Severe grazing will convert the plant community to the *Juniper/Birdfoot sagebrush Vegetation State*.

**Juniper/Inland saltgrass Plant Community**

Currently this vegetation state is found under moderate, season-long grazing by livestock. Broom snakeweed, junipers, and some conifers, are significant components of this plant community. Saline tolerant grasses make up the majority of the understory with the balance made up of short warm-season grasses, annual cool-season grasses, and miscellaneous forbs.

Dominant grasses include inland saltgrass, Sandberg bluegrass, and threadleaf sedge. Forbs, commonly found in this plant community, include hairy goldaster, prairie thermopsis, and scarlet globemallow. Sagebrush and broom snakeweed canopy ranges from 20% to 30%. Juniper and conifer canopy ranges up to 20%. Plains pricklypear and winterfat can also occur.

When compared to the Historical Climax Plant Community, sagebrush, junipers, and conifers have increased. Rhizomatous wheatgrasses and little bluestem have decreased. Broom snakeweed has invaded the state. The overstory of sagebrush, juniper, conifers, and broom snakeweed has reduced production but does provide a diverse plant community that will support some domestic livestock and wildlife such as mule deer and antelope.

The total annual production (air-dry weight) of this state is about 350 pounds per acre, but it can range from about 250 lbs./acre in unfavorable years to about 450 lbs./acre in above average years.

The following is the growth curve expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	10	25	40	10	5	5	5	0	0

(Monthly percentages of total annual growth)

The state is not well protected from excessive erosion. The biotic integrity of this plant community is usually intact but can be at risk depending on how far a shift has occurred in plant composition toward sagebrush, junipers, and bare ground. The watershed is usually functioning but can become at risk when canopy cover of sagebrush, junipers, conifers and bare ground increases.

Transitional pathways leading to other plant communities are as follows:

- Prescribed grazing use will prevent further deterioration and may return this state to near *Historic Climax Plant Community*.
- Frequent and Severe grazing over the long-term will convert this state to the *Juniper/Birdfoot Vegetation State*.

### **Juniper/Birdfoot sagebrush Plant Community**

Birdfoot sagebrush, junipers and conifers are significant components of this plant community. Cool-season grasses have been reduced. Bare ground and cool-season annual plants dominate the understory.

Perennial grasses are sparse, and include inland saltgrass and Sandberg bluegrass. Cheatgrass is the dominant grass. Forbs, commonly found in this plant community, include hairy goldaster, woody aster, prairie thermopsis, and scarlet globemallow. Birdfoot sagebrush has increased. Juniper canopy ranges up to 50%.

When compared to the Historical Climax Plant Community, birdfoot sagebrush and junipers have increased. Rhizomatous wheatgrasses and little bluestem have decreased. Bare ground has increased. The overstory of sagebrush, juniper, conifers, and broom snakeweed reduce the grass understory and reduce the sites ability to support wildlife.

The total annual production (air-dry weight) of this state is about 300 pounds per acre, but it can range from about 250 lbs./acre in unfavorable years to about 350 lbs./acre in above average years.

The following is the growth curve expected during a normal year:

Growth curve number:

Growth curve name:

Growth curve description:

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	0	15	30	30	10	5	5	5	0	0

(Monthly percentages of total annual growth)

The state is unstable and is not protected from excessive erosion. The biotic integrity of this plant community is not intact. This state is at risk depending on how far a shift has occurred in plant composition toward sagebrush, junipers, broom snakeweed and bare ground. The watershed is not functioning.

Transitional pathways leading to other plant communities are as follows:

- Prescribed grazing use may stabilize this state and return to an upward trend.

## Ecological Site Interpretations

### Animal Community – Wildlife Interpretations

**Historic Climax Plant Community:** The predominance of grasses in this plant community favors grazers and mixed-feeders, such as bison, elk, and antelope. Suitable thermal and escape cover for deer may be limited due to the low quantities of woody plants. However, topographical variations could provide some escape cover. When found adjacent to sagebrush dominated states, this plant community may provide brood rearing/foraging areas for sage grouse, as well as lek sites. Other birds that would frequent this plant community include western meadowlarks, horned larks, and golden eagles. Many grassland obligate small mammals would occur here.

**Juniper/Inland saltgrass:** This plant community can provide important winter foraging for elk, mule deer and antelope. This community provides escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for sage grouse.

**Juniper/Birdfoot sagebrush:** This plant community can provide winter foraging for elk, mule deer and antelope. Forage may be limited by low production. This community provides escape and thermal cover for large ungulates, as well as nesting and brood rearing habitat for tree nesting birds.

**Animal Preferences (Quarterly - 1,2,3,4) for commonly occurring plants in MLRA 58B, 15-17 inch Northern Plains**

COMMON NAME/ GRASSES/GRASSLIKES	SCIENTIFIC NAME	SCI. SYMBOL	Cattle	Sheep	Horses	Mule Deer	Antelope
alkali bluegrass	Poa secunda ssp. juncofolia	POSEJ	DDDD	PPPP	DDDD	PPPP	PPPP
alkali cordgrass	Spartina gracilis	SPGR	DDDD	UUUU	DDDD	UUUU	UUUU
alkali sacaton	Sporobolus airoides	SPA1	PPPP	DDDD	PPPP	DDDD	DDDD
Baltic rush	Juncus balticus	JUBA	DDDD	UUUU	DDDD	UUUU	UUUU
basin wildrye	Leymus cinereus	LEC4	PPPP	PPPP	PPPP	DDDD	DDDD
bearded wheatgrass	Elymus caninus	ELCA	PPPP	DDDD	PPPP	DDDD	DDDD
big bluestem	Andropogon gerardii	ANGE	PPPP	PPPP	PPPP	DDDD	DDDD
blue grama	Bouteloua gracilis	BOGR2	DDDD	DDDD	DDDD	DDDD	DDDD
bluebunch wheatgrass	Pseudoroegneria spicata	PSSP6	PPPP	PPPP	PPPP	DDDD	DDDD
bluejoint reedgrass	Calamagrostis canadensis	CACA4	PPPP	DDDD	PPPP	UUUU	UUUU
bottlebrush squirreltail	Elymus elymoides	ELELE	DDDD	DDDD	DDDD	UUUU	UUUU
buffalograss	Buchloe dactyloides	BUDA	DDDD	DDDD	DDDD	DDDD	DDDD
Canada wildrye	Elymus canadensis	ELCA4	PPPP	PPPP	PPPP	DDDD	DDDD
Canby bluegrass	Poa canbyi (syn. to Poa secunda)	POCA (POSE)	PPPP	PPPP	PPPP	PPPP	PPPP
Cusick's bluegrass	Poa cusickii	POCU3	PPPP	PPPP	PPPP	PPPP	PPPP
Fendler threeawn	Aristida purpurea	ARPUL	UUUU	UUUU	UUUU	UUUU	UUUU
green needlegrass	Nassella viridula	NAV14	PPPP	PPPP	PPPP	PPPP	PPPP
hairly grama	Bouteloua hirsuta	BOH12	DDDD	DDDD	DDDD	DDDD	DDDD
Indian ricegrass	Achnatherum hymenoides	ACHY	PPPP	PPPP	PPPP	PPPP	PPPP
inland saltgrass	Distichlis spicata	DISP	UUUU	UUUU	UUUU	UUUU	UUUU
inland sedge	Carex interior	CAIN11	DDDD	DDDD	DDDD	UUUU	UUUU
little bluestem	Schizachyrium scoparium	SCSC	PPPP	PPPP	PPPP	DDDD	DDDD
mat muhly	Muhlenbergia richardsonis	MURI	UUUU	UUUU	UUUU	UUUU	UUUU
Nebraska sedge	Carex nebraskensis	CANE2	PPPP	PPPP	PPPP	DDDD	DDDD
needleandthread	Hesperostipa comata	HECO26	PPPP	PPPP	PPPP	PPPP	PPPP
needleleaf sedge	Carex duriuscula	CADU6	UUUU	UUUU	UUUU	UUUU	UUUU
northern reedgrass	Calamagrostis stricta	CAST13	PPPP	DDDD	PPPP	UUUU	UUUU
Nuttall's alkaligrass	Puccinellia nuttalliana	PUNU2	PPPP	PPPP	PPPP	PPPP	PPPP
plains muhly	Muhlenbergia cuspidata	MUCU3	DDDD	DDDD	DDDD	UUUU	UUUU
plains reedgrass	Calamagrostis montanensis	CAMO	DDDD	DDDD	DDDD	DDDD	DDDD
prairie cordgrass	Spartina pectinata	SPPE	PPPP	DDDD	PPPP	UUUU	UUUU
prairie junegrass	Koeleria macrantha	KOMA	DDDD	DDDD	DDDD	DDDD	DDDD
prairie sandreed	Calamovilfa longifolia	CALO	PPPP	DDDD	PPPP	UUUU	UUUU
sand bluestem	Andropogon halli	ANHA	PPPP	DDDD	PPPP	UUUU	UUUU
sand dropseed	Sporobolus cryptandrus	SPCR	DDDD	DDDD	DDDD	UUUU	UUUU
Sandberg bluegrass	Poa secunda	POSE	DDDD	DDDD	DDDD	DDDD	DDDD
sideoats grama	Bouteloua curtipendula	BOCU	PPPP	PPPP	PPPP	DDDD	UUUU
slender wheatgrass	Elymus trachycaulus	ELTR7	PPPP	DDDD	PPPP	DDDD	DDDD
spike sedge	Carex nardina	CANA2	DDDD	DDDD	DDDD	UUUU	UUUU
sun sedge	Carex inops ssp. heliophila	CAINH2	PPPP	DDDD	PPPP	UUUU	UUUU
thickspike wheatgrass	Elymus lanceolatus	ELLAL	DDDD	DDDD	DDDD	DDDD	DDDD
threadleaf sedge	Carex filifolia	CAFI	DDDD	DDDD	DDDD	DDDD	PPPP
tufted hairgrass	Deschampsia caespitosa	DECA18	PPPP	PPPP	PPPP	DDDD	DDDD
western wheatgrass	Pascopyrum smithii	PASM	DDDD	DDDD	DDDD	DDDD	DDDD
<b>FORBS</b>							
American licorice	Glycyrrhiza lepidota	GLLE3	UUUU	UUUU	UUUU	UUUU	UUUU
American vetch	Vicia americana	VIAM	PPPP	PPPP	PPPP	PPPP	PPPP
arrowgrass	Triglochin spp.	TRIGL	T	T	T	T	T
asters	Asters	ASTER	UUUU	UUUU	UUUU	UUUU	UUUU
biscuitroots	Lomatium spp.	LOMAT	DDDD	DDDD	UUUU	DDDD	DDDD
bluebells	Mertensia	MERTE	DDDD	PPPP	DDDD	DDDD	DDDD
blue-eyed grass	Sisyrinchium spp.	SISYR	DDDD	PPPP	DDDD	DDDD	DDDD
breadroot scurfpea	Pediemelum esculentum	PEES	DDDD	DDDD	DDDD	DDDD	DDDD
cattail, broad-leaf	Typha latifolia	TYLA	DDDD	UUUU	DDDD	UUUU	UUUU
cattail, narrow-leaf	Typha angustifolia	TYAN	DDDD	UUUU	DDDD	UUUU	UUUU
fringed sagewort	Artemisia frigida	ARFR4	UUUU	UUUU	UUUU	UUUU	UUUU
green sagewort	Artemisia dracuncululus	ARDR4	UUUU	UUUU	UUUU	UUUU	UUUU
hawksbeard	Crepis acuminata	CRAC2	UUUU	PPPP	UUUU	DDDD	DDDD
horsetails	Equisetum spp.	EQUIS	UUUU	UUUU	UUUU	UUUU	UUUU
iris	Iris spp.	IRIS	UUUU	UUUU	UUUU	UUUU	UUUU
milkvetches	Astragalus	ASTRA	DDDD	DDDD	DDDD	DDDD	DDDD
poison hemlock	Conium maculatum	COMA2	T	T	T	T	T
prairie coneflower	Ratibida columnifera	RACO3	DDDD	PPPP	DDDD	PPPP	PPPP
prairie thermopsis	Thermopsis rhombifolia	THRHA	UUUU	UUUU	UUUU	UUUU	UUUU
purple prairie clover	Dalea purpurea	DAPU5	PPPP	PPPP	PPPP	PPPP	PPPP
Pursh seepweed	Suaeda calceoliformis	SUCA2	UUUU	UUUU	UUUU	UUUU	UUUU
rosy pussytoes	Antennaria rosea	ANRO2	UUUU	UUUU	UUUU	UUUU	UUUU
scarlet gaura	Gaura coccinea	GACO5	UUUU	UUUU	UUUU	UUUU	UUUU
stemless goldenweed	Haplopappus acaulis	HAAC	UUUU	UUUU	UUUU	UUUU	UUUU
sulphur flower buckwheat	Eriogonum umbellatum	ERUM	UUUU	UUUU	UUUU	UUUU	UUUU
twogrooved milkvetch	Astragalus bisulcatus	ASBI2	T	T	T	T	T
water hemlocks	Cicuta spp.	CICUT	T	T	T	T	T
western yarrow	Achillea lanulosa	ACHIL	UUUU	UUUU	UUUU	UUUU	UUUU
white prairie clover	Dalea candida	DACA7	PPPP	PPPP	PPPP	PPPP	PPPP
wild onion	Allium textile	ALTE	DDDD	DDDD	DDDD	DDDD	DDDD
woodyaster	Xylorhiza spp.	XYLOR	T	T	T	T	T
<b>TREES, SHRUBS &amp; HALF-SHRUBS</b>							
big sagebrush	Artemisia tridentata	ARTR2	UUUU	DDDD	UUUU	DDDD	DDDD
birdfoot sagebrush	Artemisia pedatifida	ARPE6	UUUU	UUUU	UUUU	UUUU	UUUU
black greasewood	Sarcobatus vermiculatus	SAVE4	DDDD	DDDD	UUUU	DDDD	DDDD
bur oak	Quercus macrocarpa	QUMA2	UUUU	DDDD	UUUU	PPPP	DDDD
fourwing saltbush	Atriplex canescens	ATCA2	PPPP	PPPP	PPPP	PPPP	PPPP
Gardners saltbush	Atriplex gardneri	ATGA	PPPP	PPPP	DDDD	PPPP	PPPP
green rabbitbrush	Chrysothamnus viscidiflorus	CHV18	DDDD	DDDD	DDDD	DDDD	DDDD
junipers	Juniperus scopulorum	JUSC2	UUUU	UUUU	UUUU	DDDD	UUUU
leadplant	Amorpha canescens	AMCA6	PPPP	PPPP	PPPP	PPPP	PPPP
plains cottonwood (sprouts)	Populus deltoides	PODEM	DDDD	DDDD	DDDD	DDDD	DDDD
ponderosa pine (abortion in cattle)	Pinus ponderosa	PIPO	UUUU	UUUU	UUUU	UUUU	UUUU
rubber rabbitbrush	Ericameria nauseosa	ERNA10	UUUU	DDDD	UUUU	DDDD	DDDD
silver sagebrush	Artemisia cana	ARCAC5	DDDD	DDDD	DDDD	PPPP	PPPP
silverberry	Eleagnus commutata	ELCO	UUUU	UUUU	UUUU	DDDD	UUUU
skunkbush sumac	Rhus trilobata	RHTR	DDDD	DDDD	DDDD	DDDD	DDDD
western snowberry	Symphoricarpos occidentalis	SYOC	UUUU	UUUU	UUUU	DDDD	UUUU
wildrose	Rosa woodsii var. woodsii	ROWOW	DDDD	DDDD	UUUU	DDDD	DDDD
willows	Salix L.	SALIX	PPPP	PPPP	DDDD	PPPP	UUUU
winterfat	Krascheninnikovia lanata	KRLA2	PPPP	PPPP	PPPP	PPPP	PPPP
yucca	Yucca glauca	YUGL	DDDD	DDDD	DDDD	DDDD	DDDD

**N = not used; U = undesirable; D = desirable; P = preferred; T = toxic**

## Animal Community – Grazing Interpretations

The following table lists suggested stocking rates for cattle under continuous season-long grazing under normal growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of the conservation planning process. Often, the current plant composition does not entirely match any particular plant community (as described in this ecological site description). Because of this, a field visit is recommended, in all cases, to document plant composition and production. More precise carrying capacity estimates should eventually be calculated using this information along with animal preference data, particularly when grazers other than cattle are involved. Under more intensive grazing management, improved harvest efficiencies can result in an increased carrying capacity. If distribution problems occur, stocking rates must be reduced to maintain plant health and vigor.

<b>Plant Community</b>	<b>Production (lb./ac)</b>	<b>Carrying Capacity* (AUM/ac)</b>
Historic Climax Plant Community	300-700	.2
Juniper/Inland saltgrass	250-450	.15
Juniper/Birdfoot sagebrush	250-350	.05

\* - Continuous, season-long grazing by cattle under average growing conditions.

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland in this area may provide yearlong forage for cattle, sheep, or horses. During the dormant period, the forage for livestock use needs to be supplemented with protein because the quality does not meet minimum livestock requirements.

## Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group D. Infiltration ranges from slow to moderate. Runoff potential for this site varies from moderate to very high depending on soil hydrologic group and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where short-grasses form a strong sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Part 630, NRCS National Engineering Handbook for detailed hydrology information).

Rills and gullies should not typically be present. Water flow patterns may be present but should be barely distinguishable. Pedestals are only slightly present in association with bunchgrasses such as bluebunch wheatgrass. Litter typically falls in place, and signs of movement are not common. Chemical and physical crusts are rare to non-existent. Cryptogamic crusts are present, but only cover 1-2% of the soil surface.

## Recreational Uses

This site provides hunting opportunities for upland game species. The wide variety of plants which bloom from spring until fall have an esthetic value that appeals to visitors.

Site Type: Rangeland  
MLRA: 58B – Northern Rolling High Plains

**Shale 15-17” P.Z.  
R058BY254WY**

## Wood Products

No appreciable wood products are present on the site.

## Other Products

None noted.

## Supporting Information

### Associated Sites

Very Shallow                      058BY276WY

### Similar Sites

() – Shale 10-14” Northern Plains P.Z.      058BY154WY  
[Lower production]

### Inventory Data References (narrative)

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range trained personnel was also used. Those involved in developing this site include: Glen Mitchell, Range Management Specialist, NRCS; Chuck Ring, Range Management Specialist, NRCS; and Everet Bainter, Range Management Specialist. Other sources used as references include: USDA NRCS Water and Climate Center, USDA NRCS National Range and Pasture Handbook, and USDA NRCS Soil Surveys from various counties.

### Inventory Data References

<u>Data Source</u>	<u>Number of Records</u>	<u>Sample Period</u>	<u>State</u>	<u>County</u>
SCS-RANGE-417		1971-1994	WY	Campbell & others
Ocular estimates		1990-1999	WY	Campbell & others

### State Correlation

This site occurs entirely within Wyoming.

### Type Locality

### Field Offices

Gillette, Lusk, Newcastle, Sundance

### Relationship to Other Established Classifications

### Other References

Site Type: Rangeland  
MLRA: 58B – Northern Rolling High Plains

**Shale 15-17” P.Z.  
R058BY254WY**

## Site Description Approval

\_\_\_\_\_  
State Range Management Specialist

\_\_\_\_\_  
Date

## Ecological Reference Worksheet

**Author(s)/participant(s):** \_\_\_\_\_

**Contact for lead author:** \_\_\_\_\_ **Reference site used? Yes/No**

**Date:** 4/05 **MLRA:** 58B **Ecological Site:** R058BY254WY Shale (Sh) 15-17"NP This *must* be verified based on soils and climate (see Ecological Site Description). Current plant community *cannot* be used to identify the ecological site.

**Indicators.** For each indicator, describe the potential for the site. Where possible, (1) use numbers, (2) include expected range of values for above- and below-average years for **each** community within the reference state, when appropriate & (3) cite data. Continue descriptions on separate sheet.

**1. Number and extent of rills:** Rills will be continuous.

**2. Presence of water flow patterns:** Water flow paths will be obvious, regular and continuous with debris dams occurring only on lesser slopes.

**3. Number and height of erosional pedestals or terracettes:** Erosional pedestals present with terracettes present at debris dams.

**4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, plant canopy are *not* bare ground):** Bare ground is 70 to 80%.

**5. Number of gullies and erosion associated with gullies:** Active gullies may be present on steeper slopes.

**6. Extent of wind scoured, blowouts and/or depositional areas:** None.

**7. Amount of litter movement (describe size and distance expected to travel):** Plant litter movement is expected.

**8. Soil surface (top few mm) resistance to erosion (stability values are averages – most sites will show a range of values for both plant canopy and interspaces, if different):** Plant cover and litter is at 20% or greater of soil surface. Stability class anticipated to be 3 or greater.

**9. Soil surface structure and SOM content (include type and strength of structure, and A-horizon color and thickness for both plant canopy and interspaces, if different):** Use soil series description for depth and color of A-horizon.

**10. Effect of plant community composition (relative proportion of different functional groups) & spatial distribution on infiltration & runoff:** Sparse plant canopy, slow infiltration rates, and the high amount of bare ground contribute to a naturally high runoff rate even in HCPC.

**11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** No compaction layer would be expected but soil surface is typically crusted and hard to very hard when dry.

**12. Functional/Structural Groups (list in order of descending dominance by above-ground weight using symbols: >>, >, = to indicate much greater than, greater than, and equal to):** Mid stature bunch grasses > shrubs > Mid stature rhizomatous grasses > short stature grasses/grasslikes > forbs.

**13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** Some plant mortality and decadence (10 to 15%) is expected on this site.

**14. Average percent litter cover (10-20%) and depth ( 0.0 to 0.25\_ inches).** Litter cover is in contact with soil surface with little evidence of biological activity.

**15. Expected annual production (this is TOTAL above-ground production, not just forage production):**  
500 lbs/acre

**16. Potential invasive (including noxious) species (native and non-native). List species which characterize degraded states and which, after a threshold is crossed, “can, and often do, continue to increase regardless of the management of the site and may eventually dominate the site”:** Buckwheats, plains prickly pear, broom snakeweed and species found on noxious weed list.

**17. Perennial plant reproductive capability:** Limited ability to reproduce.