

**NATURAL RESOURCES CONSERVATION SERVICE  
CONSERVATION PRACTICE STANDARD**

**FEED MANAGEMENT**

(No. of Systems and AUs Affected)

CODE 592

**DEFINITION**

Managing the quantity of available nutrients fed to livestock and poultry for their intended purpose.

**PURPOSE**

- Supply the quantity of available nutrients required by livestock and poultry for maintenance, production, growth, performance, animal health, and reproduction; while reducing the quantity of nutrients, especially nitrogen and phosphorus, excreted in manure by minimizing the over-feeding of these and other nutrients.
- Improve net farm income by feeding nutrients more efficiently.

**CONDITIONS WHERE PRACTICE APPLIES**

Confined livestock and poultry operations with a whole farm nutrient imbalance, with more nutrients imported to the farm than are exported and/or utilized by cropping programs.

Confined livestock and poultry operations that have a significant build up of nutrients in the soil due to land application of manure.

Confined livestock and poultry operations that land apply manure and do not have a land base large enough to allow nutrients to be applied at rates recommended by soil test and utilized by crops in the rotation.

Livestock and poultry operations seeking to enhance nutrient efficiencies.

**CRITERIA**

**General Criteria Applicable to All Purposes**

The diets for specific species of animals shall be developed in accordance recommendations from one of the following:

- Standards outlined in the most current recommendations of the National Research Council (NRC).
- Recommendations of the land grant university.
- Standards developed by the professional nutritionists of livestock and poultry production companies, feed companies, and/or feed suppliers.

Laboratory analysis shall be done on the formulated diet, and/or on the feed ingredients used to formulate the diet, to determine its nutrient content.

Feed analyses shall be conducted by laboratories whose tests are accepted by the Land Grant University in the state in which the feeding strategy will be implemented and by the American Feed Industry Association. Data from analyzed feed ingredients and/or appropriate historic feed analysis information for the operation will be used for adjustments of ration formulation.

Diets and feed management strategies shall be developed by professional animal scientists, independent professional nutritionists or other comparably qualified individuals. When required by state policy or regulation, animal nutritionists shall be certified through any certification program recognized within the state. It is recommended that nutritionist be "Professional Animal Scientist"

as certified by the American Registry of Professional Animal Scientists (ARPAS).

Diets shall be formulated to provide the proper quantities and correct relative ratios of available nutrients required by the animal species to meet the species requirements for which the plan is being developed.

Adjustments to nutrient amounts or concentrations shall be provided to meet specific genetic potential, environmental demands, and/or requirements to insure health, well-being and productivity.

One or more of the following feed management practices and/or diet manipulation technologies shall be used to reduce N, P and other excreted nutrients while maintaining the health, well-being and productivity of the animal.

- Formulating diets closer to animal requirements.
- Reducing protein and supplementing with amino acids for non-ruminants and supplementing with protein sources high in rumen undegradable protein or rumen protected amino acids for ruminants.
- Manipulating the crude protein and energy (carbohydrate and fat) content of the diet to enhance the rumen microbial protein synthesis to meet the requirements for amino acids (ruminants).
- Using highly digestible feeds, as appropriate, in the diet.
- Using phytase and reducing the supplemental phosphorus content of the diet (non-ruminants)
- Reducing the phosphorus content of the diet of ruminants when it is being overfed.
- Using selected enzymes or other products to enhance feed digestibility or feed use efficiency.
- Using growth promotants as allowed by law.
- Implementing phase feeding.
- Implementing split-sex feeding.

- Utilizing the Nutrient Balancing Program (NUTBAL) for monitoring of diet levels of protein and phosphorus for animals in pasture situations receiving supplemental feeds.
- Using other feed management or diet manipulation technologies that have demonstrated the ability to reduce the nutrient content of N and P in the manure.

When analysis of manure is done to determine manure nutrient content, the analysis shall be performed by laboratories whose results are accepted by the Land Grant University in the state or the American Feed Industry Association in which the feeding strategy was implemented.

## CONSIDERATIONS

Consider nutrient requirements for production based upon stage of growth, intended purpose of the animal and the type of production (e.g., meat, milk, eggs) involved.

Use management practices described in the NRCS Nutrient Management (Feed Management) Technical Notes for the specific animal species.

Weigh all feed, concentrates, forages, supplements, vitamins, and minerals that are fed and/or mixed to increase accuracy of amounts of nutrients fed so as to more closely follow the feeding plans.

Sample all feeds (purchased or farm raised) and analyzing them for moisture and nutrient values, and adjust the diet to account for changes as seen in the analysis.

Keep records of all feeds fed (forages, grains, concentrates, minerals, vitamins, etc), moisture levels and nutrient levels in order to document the need for feeding changes.

Analyzing the drinking water consumed by the animals to determine its nutrient content, and adjusting the diet to account for this source of nutrients.

Routinely clean water troughs and drinking fountains to minimize build up of bacteria.

Different feed ingredients (e.g. by-products) and their potential impacts on the nutrient content of excreted manure.

The potential impact of feed management on the volume of manure excreted and on manure storage requirements.

The impact of feed management practices, animal management practices, and diet manipulation on manure odors, pathogens, animal health and well-being.

Using concentrates and forages grown on the farm to minimize the quantity of nutrients imported to the farm, and to maximize the recycling of nutrients on the farm.

Analyzing excreted manure or manure from storage facilities to determine manure nutrient content and to estimate the impact of the feeding strategy.

### PLANS AND SPECIFICATIONS

Plans and specifications for feed management shall be in keeping with the requirements of this standard. They shall describe the specific feed management practices and/or technologies that are planned for the operation.

The following components shall be included in the feed management plan:

- The type of technology, or technologies, and/or feeding practices that will be used on the operation.
- Feed analyses and ration formulation information prior to and after implementation of feed management on the operation.
- The estimated, or measured, nutrient content of the manure prior to the implementation of feed management on the operation.
- The estimated impact that feed management will have on manure nutrient content.
- Guidance for how often the feed management plan shall be reviewed and potentially revised.
- The quantities and sources of nitrogen and phosphorus that will be fed.
- Identification of the qualified feed management specialist who developed the plan.

### OPERATION AND MAINTENANCE

The producer/client is responsible for the operation and maintenance of the feed management plan. Operation and maintenance activities address the following:

- Periodic plan review to determine if adjustments or modifications are needed.
- Routine feed analysis to document the rates at which nitrogen and phosphorus were actually fed. When actual rates fed differ from or exceed the planned rates, records will indicate the reasons for the differences.
- Routine fresh manure analysis shall document the rates at which nitrogen and phosphorus were fed to livestock through supplemental feeds and forage consumption while grazing pastures. Tracking manure analysis and stocking rates of animals will document differences in nutrients when supplemental feeding rates are adjusted.
- Maintaining records to document plan implementation. As applicable, records include:
  - ◆ Records of feed analysis and ration formulation, including the record of ration formulation used prior to implementing the feeding strategy.
  - ◆ Records of the initial estimate of the impact the feeding strategy was expected to have on reducing manure nutrient content.
  - ◆ Records of any manure analysis that was done after the feeding strategy was implemented to determine manure nutrient content.
  - ◆ Dates of review and person performing the review, and any recommendations that resulted from the review.

Records of plan implementation shall be maintained for five years, or for a period longer than five years if required by other Federal, state, or local ordinances, program, or contract requirements.