Table N.--Highly Erodible Land (HEL) List

| Map symbol | Soil name | HEL |
|---|---|---|
| BaF | Barkcamp gravelly sandy loam, 40 to 70 percent slopes | Highly erodible land |
| BkD | Berks-Westmoreland silt loams, 15 to 25 percent slopes | Highly erodible land |
| BkE | Berks-Westmoreland silt loams, 25 to 40 percent slopes | Highly erodible land |
| BkF | Berks-Westmoreland silt loams, 40 to 70 percent slopes | Highly erodible land |
| BoD | Bethesda shaly silty clay loam, 8 to 25 percent slopes | Potentially highly erodible land |
| BoE | Bethesda shaly silty clay loam, 25 to 40 percent slopes | Highly erodible land |
| BoF | Bethesda shaly silty clay loam, 40 to 70 percent slopes | Highly erodible land |
| BrC | Brookside silt loam, 8 to 15 percent slopes | Potentially highly erodible land |
| BrD | Brookside silt loam, 15 to 25 percent slopes | Highly erodible land |
| BrE | Brookside silt loam, 25 to 40 percent slopes | Highly erodible land |
| Cd | Chagrin loam, rarely flooded | Not highly erodible land |
| Cg | Chagrin silt loam, frequently flooded | Not highly erodible land |
| CmC | Clymer loam, 8 to 15 percent slopes | Potentially highly erodible land |
| DkE | Dekalb loam, 25 to 40 percent slopes | Highly erodible land |
| DsG | Dekalb and Gilpin stony soils, 25 to 70 percent slopes | Highly erodible land |
| DtD | Dekalb-Westmoreland complex, 15 to 25 percent slopes | Highly erodible land |
| DtE | Dekalb-Westmoreland complex, 25 to 40 percent slopes | Highly erodible land |
| DtF | Dekalb-Westmoreland complex, 40 to 70 percent slopes | Highly erodible land |
| DuF | Dekalb-Westmoreland complex, benched, 40 to 70 percent slopes | Highly erodible land |
| DxA | Doles silt loam, 0 to 3 percent slopes | Not highly erodible land |
| Dy | Dumps, mine | |
| EbF | Elba-Brookside-Berks complex, 40 to 70 percent slopes | Highly erodible land |
| FaD | Fairpoint silt loam, 8 to 25 percent slopes | Highly erodible land |
| FbE | Fairpoint shaly clay loam, 25 to 40 percent slopes | Highly erodible land |
| FbF | Fairpoint shaly clay loam, 40 to 70 percent slopes | Highly erodible land |
| FcA | Fitchville silt loam, 0 to 3 percent slopes | Potentially highly erodible land |
| GaC | Gallia loam, 8 to 15 percent slopes | Potentially highly erodible land |
| GmA | Glenford silt loam, 0 to 3 percent slopes | Not highly erodible land |
| GmB | Glenford silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| GmC | Glenford silt loam, 8 to 15 percent slopes | Potentially highly erodible land |
| GsB | Guernsey silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| GsC | Guernsey silt loam, 8 to 15 percent slopes | Highly erodible land |
| GuC | Guernsey-Upshur complex, 8 to 15 percent slopes | Highly erodible land |
| GuD | Guernsey-Upshur complex, 15 to 25 percent slopes | Highly erodible land |
| HcA | Hackers silt loam, 0 to 3 percent slopes | Potentially highly erodible land |
| LkB | Licking silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| LkC | Licking silt loam, 8 to 15 percent slopes | Highly erodible land |

| | | |
|---|---|---|
| McA | McGary silt loam, 0 to 3 percent slopes | Potentially highly erodible land |
| Mh | Melvin silt loam, frequently flooded | Not highly erodible land |
| Mp | Moshannon silt loam, frequently flooded | Not highly erodible land |
| NeC | Negley loam, 8 to 15 percent slopes | Highly erodible land |
| NgE | Negley gravelly loam, 25 to 40 percent slopes | Highly erodible land |
| Nn | Newark silt loam, frequently flooded | Not highly erodible land |
| No | Nolin silt loam, frequently flooded | Not highly erodible land |
| Or | Orrville silt loam, frequently flooded | Not highly erodible land |
| OtB | Omulga silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| OtC | Omulga silt loam, 8 to 15 percent slopes | Highly erodible land |
| PaB | Parke silt loam, 2 to 6 percent slopes | Potentially highly erodible land |
| Pg | Pits, gravel | |
| RcC | Richland loam, 8 to 15 percent slopes | Potentially highly erodible land |
| RcD | Richland loam, 15 to 25 percent slopes | Highly erodible land |
| RcE | Richland loam, 25 to 40 percent slopes | Highly erodible land |
| StD | Steinsburg sandy loam, 15 to 25 percent slopes | Highly erodible land |
| StE | Steinsburg sandy loam, 25 to 40 percent slopes | Highly erodible land |
| StF | Steinsburg sandy loam, 40 to 70 percent slopes | Highly erodible land |
| Ud | Udorthents, loamy | |
| UpC | Upshur silty clay loam, 8 to 15 percent slopes | Highly erodible land |
| UpD | Upshur silty clay loam, 15 to 25 percent slopes | Highly erodible land |
| UsC | Upshur-Elba silty clay loams, 8 to 15 percent slopes | Highly erodible land |
| VaC | Vandalia silty clay loam, 8 to 15 percent slopes | Potentially highly erodible land |
| VbD | Vandalia-Brookside complex, 15 to 25 percent slopes | Highly erodible land |
| VbE | Vandalia-Brookside complex, 25 to 40 percent slopes | Highly erodible land |
| VcD | Vandalia-Richland complex, 15 to 25 percent slopes | Highly erodible land |
| VcE | Vandalia-Richland complex, 25 to 40 percent slopes | Highly erodible land |
| VqD2 | Vandalia silt loam, 15 to 25 percent slopes, eroded | Highly erodible land |
| VtC | Vincent silt loam, 6 to 12 percent slopes | Highly erodible land |
| W | Water | |
| WdB | Wellston silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| WdC | Wellston silt loam, 8 to 15 percent slopes | Potentially highly erodible land |
| WeB | Westmore silt loam, 3 to 8 percent slopes | Potentially highly erodible land |
| WeC | Westmore silt loam, 8 to 15 percent slopes | Potentially highly erodible land |
| WhC | Westmoreland-Guernsey silt loams, 8 to 15 percent slopes | Highly erodible land |
| WhD | Westmoreland-Guernsey silt loams, 15 to 25 percent slopes | Highly erodible land |
| WhE | Westmoreland-Guernsey silt loams, 25 to 40 percent slopes | Highly erodible land |
| WhF | Westmoreland-Guernsey silt loams, 40 to 70 percent slopes | Highly erodible land |
| WkF | Westmoreland-Guernsey silt loams, benched, 40 to 70 percent slopes | Highly erodible land |
| WmC | Westmoreland-Upshur complex, 8 to 15 percent slopes | Highly erodible land |
| WmD | Westmoreland-Upshur complex, 15 to 25 percent slopes | Highly erodible land |
| WmE | Westmoreland-Upshur complex, 25 to 40 percent slopes | Highly erodible land |
| WmF | Westmoreland-Upshur complex, 40 to 70 percent slopes | Highly erodible land |
| WpB | Wheeling loam, 3 to 10 percent slopes | Potentially highly erodible land |
| WtB | Woodsfield silt loam, 3 to 8 percent slopes | Potentially highly |

| | | |
|---|---|---|
| | | erodible land |
| WtC | Woodsfield silt loam, 8 to 15 percent slopes | Highly erodible land |
| ZnB | Zanesville silt loam, 3 to 8 percent slopes | Potentially highly erodible land |