


# Ecological Site Description

## *Till Exposed Backslope Woodland*

Major Land Resource Area 109  
Iowa and Missouri Heavy Till Plain



**Contact Information for this publication:** *Editors* – Douglas Wallace ([doug.wallace@mo.usda.gov](mailto:doug.wallace@mo.usda.gov)), Ecologist, United States Department of Agriculture (USDA), ACES Program - Natural Resources Conservation Service (NRCS) and Dr. Fred Young ([fred.young@mo.usda.gov](mailto:fred.young@mo.usda.gov)), Soil Scientist, USDA, ACES-NRCS, Columbia, MO; *Project Supervisors* – Jorge Lugo-Camacho ([jorge.lugo-camacho@mo.usda.gov](mailto:jorge.lugo-camacho@mo.usda.gov)), State Soil Scientist, USDA-NRCS and Dwaine Gelnar ([dwaine.gelnar@mo.usda.gov](mailto:dwaine.gelnar@mo.usda.gov)), State Resource Conservationist, USDA-NRCS, Columbia, MO; *Field Data Support* – Alicia Struckhoff ([Alicia.Struckhoff@mdc.mo.gov](mailto:Alicia.Struckhoff@mdc.mo.gov)), ESD project Leader, Missouri Department of Conservation (MDC), and Stribling Stuber ([stribling.stuber@mdc.mo.gov](mailto:stribling.stuber@mdc.mo.gov)), Field Ecologist, MDC, Columbia, MO. Additional field and technical support provided by Tim Nigh, retired MDC Ecologist, and Dennis Meinert, Soil Scientist, Missouri Department of Natural Resources.

**Cover photos:**

*Center photo* (from Missouri Department of Conservation) is an exposed till backslope woodland found on southerly and westerly aspects of steep, dissected slopes in MLRA 109.

*Top right map* is the distribution map for Till Exposed Backslope Woodland (see page 5 for more detail).

*This publication is a multi-agency effort with input from a wide range of natural resource specialists. NRCS and the Missouri Department of Conservation are leading this effort along with the University of Missouri, Missouri Department of Natural Resources, USDA Forest Service, and U.S. Fish and Wildlife Service. Ecological site information is available via the NRCS Web Soil Survey, USDA Ecological Site Information System website, and the Missouri Field Office Technical Guide. The information in this publication was developed using historical data, professional experience, field reviews, and scientific studies. The information is representative of complex communities. Key indicator plants, animals and ecological processes are described to help guide land management decisions.*

## Table of Contents

<b>Section I: Ecological Site Characteristics</b>	5
Ecological Site Identification	5
Introduction	5
Ecological Site Concept	5
Physiographic Features	5
Climatic Features	6
Influencing Water Features	7
Representative Soil Features	7
Ecological Site Dynamics	8
State and Transition Diagram	10
Ecological States and Community Phases	11
STATE 1 – REFERENCE	11
Community Phase 1.1	11
Structure and Cover	12
Community Phase 1.2	18
Structure and Cover	18
STATE 2 – TIMBER MANAGED WOODLAND	20
Community Phase 2.1	21
Community Phase 2.2	21
STATE 3 – FIRE MANAGED WOODLAND	21
Community Phase 3.1	22
Community Phase 3.2	22
STATE 4 – GRASSLAND	22
Community Phase 4.1	23
Community Phase 4.2	23
Community Phase 4.3	24
STATE 5 – HIGH GRADED/GRAZED WOODLAND	24
Community Phase 5.1	24
Community Phase 5.2	25
<b>Section II: Ecological Site Interpretations</b>	26
Forest Site Productivity	26
Animal Community	26
Hydrology Functions	26
Recreational Uses	27
Wood Products	27
Supporting Information	27
Associated Ecological Sites	27
Similar Ecological Sites	27
State Correlation	28
Hierarchical Classification Relationships	28
<b>Section III: Inventory Data Collection</b>	29
Inventory Data References	29
Sampling Methods	29
Type Localities for Reference State Data Plots	29
References	31
Glossary	32


## Section I: Ecological Site Characteristics

### Ecological Site Identification

- *Site common name:* Till Exposed Backslope Woodland
- *Site biotic name:* white oak - black oak / fragrant sumac / hairy woodland brome - hairy sunflower
- *Site ID:* F109XY022MO
- *Major land resource area (MLRA):* 109 – Iowa and Missouri Heavy Till Plain

### Introduction

The Iowa and Missouri Heavy Till Plain (area outlined in red on the map) is an area of rolling hills interspersed with interfluvial divides and alluvial valleys. Elevation ranges from about 660 feet along the lower reaches of rivers, to about 980 feet on stable interfluvial summits in southern Iowa. Relief is about 80 to 160 feet between major streams and adjacent interfluvial summits. Most of the till plain drains south to the Missouri River via the Grand and Chariton River systems, but the northeastern portion drains southeast to the Mississippi River. Loess caps the pre-Illinoian aged till on interfluvial divides, whereas the till is exposed on side slopes. Mississippian aged limestone and Pennsylvanian aged sandstone and shale crop out on lower slopes in some areas.



### Ecological Site Concept


Till Exposed Backslope Woodlands, within the green areas on the map, occupy the southerly and westerly aspects of steep, dissected slopes, and are mapped in complex with the Till Protected Backslope Forest ecological site. These ecological sites occur primarily in the eastern and southeastern portion of the Till Plain. They are typically

downslope from Loess Upland Woodland or Till Upland Woodland ecological sites, and generally occupy the mid to lowest portion of the hillslope. In a few places, a narrow band of Shale Protected Backslope ecological site is downslope. Soils are very deep, with dense till subsoils that are mainly clay loam. The reference plant community is woodland dominated by an overstory of white oak and black oak, with an open understory and a dense, diverse herbaceous ground flora.

### Physiographic Features

This site is on upland backslopes, with slopes of 14 to 45 percent. It is on exposed aspects (south, southwest, and west), which receive significantly more solar radiation than the protected aspects. The site receives runoff from upslope summit and shoulder sites, and generates runoff to adjacent, downslope ecological sites. This site does not flood.

The adjacent figure (adapted from Festervand, 1994) shows the typical landscape position of this ecological site, and landscape relationships among the major ecological sites in the uplands and adjacent floodplains. The site is within the area labeled “3”, and is typically downslope from Till Upland Woodland ecological sites. In areas where the local drainageways have not dissected into the underlying residuum, Upland Drainageway or Floodplain ecological sites are directly downslope.



Landforms: (1) Hill

	<u>Minimum</u>	<u>Maximum</u>
<i>Slope (percent):</i>	14	35
<i>Water table depth (inches):</i>	24	72
<i>Flooding Frequency:</i>	None	None
<i>Ponding Frequency:</i>	None	None
<i>Runoff class:</i>	High	Very high
<i>Aspect:</i>	South, South West, West	

### **Climatic Features**

The Iowa and Missouri Heavy Till Plain MLRA has a continental type of climate marked by strong seasonality. In winter, dry-cold air masses, unchallenged by any topographic barriers, periodically swing south from the northern plains and Canada. If they invade reasonably humid air, snowfall and rainfall result. In summer, moist, warm air masses, equally unchallenged by topographic barriers, swing north from the Gulf of Mexico and can produce abundant amounts of rain, either by fronts or by convectional processes. In some summers, high pressure stagnates over the region, creating extended droughty periods. Spring and fall are transitional seasons when abrupt changes in temperature and precipitation may occur due to successive, fast-moving fronts separating contrasting air masses.

This MLRA experiences small regional differences in climates that grade inconspicuously into each other. The basic gradient for most climatic characteristics is along a line from north to south. Both mean annual temperature and precipitation exhibit fairly minor gradients along this line. Mean January minimum temperature follows the north-to-south gradient. However, mean July maximum temperature shows hardly any geographic variation in the region. Mean July maximum temperatures have a range of only two to three degrees across the region. Mean annual precipitation varies along the same gradient as temperature – lower annual precipitation in the north, higher in the south. Seasonality in precipitation is very pronounced due to strong continental influences. June precipitation, for example, averages four to five times greater than January precipitation. During years when precipitation is normal, moisture is stored in the soil profile during the winter and early spring, when evaporation and transpiration are low. During the summer months the loss of water by evaporation and transpiration is high, and if rainfall fails to occur at frequent intervals, drought will result. Drought directly influences ecological communities by limiting water supplies, especially at times of high temperatures and high evaporation rates. Drought indirectly affects ecological communities by increasing plant and animal susceptibility to the probability and severity of fire. Frequent fires encourage the development of grass/forb dominated communities and understories.

Superimposed upon the basic MLRA climatic patterns are local topographic influences that create topo-climatic, or microclimatic variations. For example, air drainage at nighttime may produce temperatures several degrees lower in valley bottoms than on side slopes. At critical times during the year, this phenomenon may produce later spring or earlier fall freezes in valley bottoms. Slope orientation is an important topographic influence on climate. Summits and south-and-west-facing slopes are regularly warmer and drier, supporting more grass dominated communities than adjacent north- and-east-facing slopes that are cooler and moister that support more woody dominated communities. Finally, the climate within a canopied woodland ecological site is measurably different from the climate of the more open grassland or savanna ecological sites.

Source: University of Missouri Climate Center - <http://climate.missouri.edu/climate.php>; Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin, United States Department of Agriculture Handbook 296 - <http://soils.usda.gov/survey/geography/mlra/>

The following tables are a summary from the following Climate Stations:

- (1) KEOSAUQUA [USC00134389], Van Buren County IA 52565. Period of record 1981-2010
- (2) OSCEOLA [USC00136316], Clarke County IA 50213. Period of record 1981-2010
- (3) CHILLICOTHE 2S [USC00231580], Livingston County MO 64601. Period of record 1981-2010
- (4) UNIONVILLE [USC00238523], Putnam County MO 63565. Period of record 1981-2010

	<u>Averaged</u>
Frost-free period (days):	156
Freeze-free period (days):	183
Mean annual precipitation (inches):	21.37

Monthly Precipitation (Inches):

	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
High	1.80	2.03	3.58	4.78	6.16	6.33	6.65	5.56	5.46	4.40	3.29	2.38
Medium	1.02	1.35	2.35	3.65	4.71	4.54	4.37	3.69	3.25	2.73	2.29	1.62
Low	0.62	0.85	1.52	2.72	3.38	3.27	2.71	2.26	2.18	1.61	1.36	0.93

Monthly Temperature (°F):

	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
High	34.2	39.3	51.8	64.1	73.9	82.8	87.5	86.0	78.0	65.9	51.1	37.3
Low	14.5	18.4	29.0	39.8	50.8	60.7	65.4	63.3	53.4	41.6	30.1	18.4

**Influencing Water Features**

This ecological site is not influenced by wetland or riparian water features.

**Representative Soil Features**

These soils have no major rooting restriction. The soils were formed under woodland vegetation, and have thin, light-colored surface horizons. Parent material is till. They have loam surface layers, with dense subsoils that are mainly clay loam. These soils are not affected by seasonal wetness. Soils in this protected aspect ecological site typically have thicker surface horizons relative to similar soils on exposed aspects (Steele, 2011). Soil series associated with this site include Brevator, Keswick, Lindley, and Winnegan.



The accompanying picture of the Winnegan series shows a thin surface horizon overlying the brown clayey till. Threads and filaments of calcium carbonate are below about 70 cm in this profile, and are typical in soils of this ecological site. (Picture courtesy of Amber Steele); scale is in centimeters.

*Parent materials*

*Kind:* Till

- Surface texture:* (1) Loam  
 (2) Clay loam  
 (3) Silt loam

*Subsurface texture group:* Clayey

	<u>Minimum</u>	<u>Maximum</u>
<i>Surface fragments &lt;=3" (% cover):</i>	0	4
<i>Surface fragments &gt;3" (% cover):</i>	0	1
<i>Subsurface fragments &lt;=3" (% volume):</i>	1	10
<i>Subsurface fragments &gt;3" (% volume):</i>	0	4
<i>Drainage class:</i> Moderately well drained to well drained		
<i>Permeability class:</i> Slow		

	<u>Minimum</u>	<u>Maximum</u>
<i>Depth (inches):</i>	72	>72
<i>Available water capacity (inches):</i>	5.00	6.00
<i>Electrical conductivity (mmhos/cm):</i>	0	2
<i>Sodium adsorption ratio:</i>	0	0
<i>Calcium carbonate equivalent (percent):</i>	0	0
<i>Soil reaction (1:1 water):</i>	4.5	6.5

**Ecological Site Dynamics**

*Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information is representative of very complex vegetational communities. Not all scenarios or plants are included or discussed. Key indicator plants, animals and ecological processes are described to help guide land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. The biological processes on this site are complex. Therefore, representative values are presented in a land management context. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.*

The reference plant community is woodland dominated by an overstory of white oak (*Quercus alba*) and black oak (*Quercus velutina*). This woodland type has a moderate canopy closure (50 to 80 percent), with an open understory and a dense, diverse herbaceous ground flora. Historically, white oak dominated the canopy, along with black oak and occasional hickories, bur oak (*Quercus macrocarpa*) and post oak (*Quercus stellata*). Woodlands are distinguished from forest, by their relatively open understory, and the presence of sun-loving ground flora species (MDC, 2006; Nelson, 2010).



Fire played an important role in the maintenance of these systems. While these ecological sites normally occurred back

from the prairie edge, they typically burned at least once every 5 to 10 years. These periodic fires kept woodlands open, removed the litter, and stimulated the growth and flowering of the grasses and forbs (Frost 1996). During fire free intervals, woody understory species increased and the herbaceous understory diminished. The return of fire would open the woodlands up again and stimulate the abundant ground flora.

Today, this community has either been cleared and converted to pasture, or has grown denser in the absence of fire. Most occurrences today exhibit canopy closure of 80 to 100 percent. In addition, the sub-canopy and understory layers are better developed. Black oak and hickory (*Carya sp.*) now share dominance with white oak and there are considerably more saplings in the understory. Under these denser, more shaded conditions, the original sun-loving ground flora has diminished in diversity and cover. While some woodland species persist in the ground flora, many have been replaced by more shade-tolerant species (Steele et al. 2013).



In the long term absence of fire, woody species, especially hickory, Eastern hop hornbeam (*Ostrya virginiana*) and gooseberry (*Ribes sp.*) encroach into these woodlands. Once established, these woody plants can quickly fill the existing understory increasing shade levels, greatly diminishing ground flora. Opening the canopy, removing the younger understory and applying prescribed fire have proven to be effective restoration means.

Uncontrolled domestic grazing has also impacted these communities, further diminishing the diversity of native plants and introducing species that are tolerant of grazing, such as buckbrush (*Symphoricarpos orbiculatus*), gooseberry, and Virginia creeper (*Parthenocissus quinquefolia*). Grazed sites also have a more open understory. In addition, soil compaction and soil erosion related to over grazing can be a problem and lower site productivity.

This ecological site, if managed properly, can be a source for timber products. Most areas on this ecological site have been repeatedly logged and high graded. Partial selective cutting will perpetuate the overly dense, shaded conditions of current stands. Even-age management, using clearcut, shelterwood or seed tree harvest systems and prescribed fire can restore the more open structure and diversity of ground flora species.


Managed areas show an exceptional resiliency and production. Characteristic plants in the ground flora can be used to gauge the restoration potential of a stand along with remnant open-grown old-age trees, and tree height growth. This type of management can provide timber products, watershed protection, wildlife habitat, and potential native forage.

Finally, on some wooded sites in the northern part of the MLRA, invasive non-native species of earthworms (suborder *Lumbricina*) are beginning to have broad effects on the nutrient cycles in temperate forests. These earthworms increase the cycling and leaching of nutrients by breaking up decaying organic matter and spreading it into the soil. Temperate forests rely on thick layers of decaying organic matter for growth and nutrition. The invasive earthworm presence and activity is diminishing the diversity of native plants in these environments. This change in the plant diversity directly affects the other organisms of the environment and often leads to increased invasions of exotic species as well as overall forest decline. Restoration to a reference state under these conditions will be more difficult if not dramatically reduced or impossible (Hendrix et al. 2006; Nuzzo et al. 2009).

A State and Transition Model Diagram is depicted in Figure 1. Detailed descriptions of each state, transition, plant community, and pathway follow the model. This model is based on available experimental research, field observations, professional consensus, and interpretations. It is likely to change as knowledge increases. The diagram suggests some pathways that the vegetation on this site might take. There may be other states not shown on the diagram. This information is intended to show what might happen in a given set of circumstances. It does not mean that this would happen the same way in every instance. Local professional guidance should always be sought before pursuing a treatment scenario.

State and Transition Diagram (STD)

Till Exposed Backslope Woodland, F109XY022MO



Code	Event/Process/Activity
T1A	Fire suppression; forest timber management
T1B	Prescribed burning; forest stand improvement
T1C, T5A	Clearing; pasture planting; grassland management
T1D, T2B, T3B	High-grading harvesting; uncontrolled grazing
T2A	Prescribed fire; thinning
T3A	Fire suppression; forest timber management
T4A	Long-term succession; no grazing
T5B	Forest stand improvement; no grazing; prescribed burning
T5C	Forest stand improvement; no grazing; thinning

Code	Event/Process/Activity
1.1A	No disturbance (10+ years)
1.2A	Disturbance (fire, wind, ice) < 10 years
2.1A	No activity; idle
2.2A	Forest timber management
3.1A	No activity; idle; fire suppression
3.2A	Prescribed fire; selective thinning
4.1A	Over grazing; no management
4.2A	Brush management; grassland seeding; grassland management
5.1A	No activity; idle
5.2A	High-grading; uncontrolled grazing

Code	Event/Process/Activity
R1A	Extended rotations; selective thinning
R1B	Prescribed fire; extended rotations; selective thinning

## Ecological States and Community Phases

### State 1: Reference

The reference state is dominated by white oak associated with black oak and other mixed hardwoods. Maximum tree age is likely 150 to 300 years. Periodic disturbances from fire, wind or ice maintain the dominance of white oak by opening up the canopy and allowing more light for white oak reproduction. Long disturbance-free periods allowed an increase in more shade tolerant species such as hop hornbeam and hickory. Two community phases are recognized in this state, with shifts between phases based on disturbance frequency.

The reference state is very rare throughout the MLRA. Many sites have been converted to grassland (State 4). Others have been subject to repeated, high-graded timber harvests coupled with domestic livestock grazing (State 5). Fire suppression has resulted in increased canopy density, which has affected the abundance and diversity of ground flora. Some reference sites have been managed for timber harvest, resulting in a timber managed woodland (State 2) or if prescribed fire has been maintained or introduced (State 3) as a fire managed woodland.

### ***Community Phase 1.1: White Oak – Black Oak/Aromatic Sumac/ Woodland Brome – Hairy Sunflower***



*(Photo on left: Reference woodland at Rudolf Bennitt State Wildlife Area near Clark, Missouri - photo from MDC)*

This phase has an overstory that is dominated by white oak and black oak with hickory and post oak also present. This woodland community has a two-tiered structure with an open understory and a dense, diverse herbaceous ground flora.

Periodic disturbances including fire, ice and wind create canopy gaps, allowing white oak and black oak to successfully reproduce and remain in the canopy.

### ***Community Phase Pathway 1.1A***

This pathway results from the absence of disturbance allowing more shade tolerant species such as eastern hop hornbeam, hickories, white ash and others increase in importance and add structural diversity to the system. Some displacement of grasses and forbs may be occurring due to shading and competition from the increased densities of hickories and oaks.

**Structure and Cover – Community Phase 1.1**

Soil Surface Cover

<u>Cover type</u>	<u>Minimum</u>	<u>Maximum</u>
Litter	0.2%	6%

<u>Downed woody material</u>	<u>Minimum</u>	<u>Maximum</u>	<u>Predominant decomposition class*</u>
Downed wood, fine-large (1.00-2.99" diameter; 100-hour fuels)	0.1%	2%	I
Downed wood, coarse-small (3.00-8.99" diameter; 1000-hour fuels)	0.1%	2%	I
Downed wood, coarse-large (>9.00" diameter; 10000-hour fuels)	0%	2%	

Tree snags\*\* per acre

Hard snags***	0	10
Soft snags***	0	0

\* Decomposition classes: N=No or little integration with the soil surface. I=Partial to nearly full integration with the soil surface.

\*\* >4" diameter at 4.5' above ground and >6' height. If diameter or height is smaller, use applicable downed wood type.

\*\*\* Hard=Tree is dead with most or all of bark intact. Soft=Most of bark has sloughed off.

**Ground Cover**

<u>Vegetative cover</u>	<u>Minimum</u>	<u>Maximum</u>
Grasses/grasslikes	0.1%	0.99%
Forbs	0.1%	0.99%
Shrubs/vines	0.1%	0.99%
Trees	1%	2%
Nonvascular plants	0%	0%
Biological crust	0%	0%

<u>Non-vegetative cover</u>	<u>Minimum</u>	<u>Maximum</u>
Litter	50%	95%
Surface fragments >0.25" and <=3"	0%	0.1%
Surface fragments >3"	0%	0.1%
Bedrock	0%	0%
Water	0%	0%
Bare ground	0.01%	2%

**Structure of Canopy Cover**

<u>Height above ground</u>	<u>Grasses/grasslikes</u>		<u>Forbs</u>		<u>Shrubs/vines</u>		<u>Trees</u>	
	<u>Minimum</u>	<u>Maximum</u>	<u>Minimum</u>	<u>Maximum</u>	<u>Minimum</u>	<u>Maximum</u>	<u>Minimum</u>	<u>Maximum</u>
<=0.5 foot	2%	10%	5%	50%	1%	2%	1%	5%
>0.5 to <1 foot	0%	2%	5%	75%	0%	25%	2%	25%
>1 to <=2 feet	0%	2%	2%	50%	0%	5%	2%	25%
>2 to <4.5 feet	0%	0%	0%	2%	0%	10%	0%	5%
>4.5 to <=13 feet	--	--	--	--	--	5%	1%	50%
>13 to <40 feet	--	--	--	--	--	--	0%	50%
<40 to >=80 feet	--	--	--	--	--	--	0%	75%
>80 to <120 feet	--	--	--	--	--	--	--	95%
>=120 feet	--	--	--	--	--	--	--	--

**Woodland Overstory**

White oak and black oak dominate with scattered hickory and post oak.

**Woodland Overstory Characterization Summary (trees >13 feet in height)**

Woodland canopy: Low canopy cover % 50 RV canopy cover % 65 High canopy cover % 80

Overstory plant type: Tree

Name	Symbol	Nativity	Cover low %	Cover high %	Canopy height bottom	Canopy height top	Tree diameter low	Tree diameter high	Basal area low	Basal area high
SHAGBARK HICKORY <i>Carya ovata</i>	<a href="#">CAOV2</a>	N	2.0	10.0	90.0	5.0	11.0	6.7	8.6	
WHITE OAK <i>Quercus alba</i>	<a href="#">QUAL</a>	N	2.0	95.0	90.0	5.0	27.0	90.0	134.0	
NORTHERN RED OAK <i>Quercus rubra</i>	<a href="#">QURU</a>	N	5.0	25.0	90.0	13.0	21.0	9.3	24.0	
BLACK OAK <i>Quercus velutina</i>	<a href="#">QUVE</a>	N	10.0	25.0	85.0	13.0	17.0	25.0	25.0	
MOCKERNUT HICKORY <i>Carya alba(syn)</i>	<a href="#">CAAL27</a>	N	5.0	20.0	70.0					
BUR OAK <i>Quercus macrocarpa</i>	<a href="#">QUMA2</a>	N	5.0	10.0	70.0					
POST OAK <i>Quercus stellata</i>	<a href="#">QUST</a>	N	5.0	10.0	60.0					

**Woodland Understory**

The understory layer is a dense native forb and grass ground layer with scattered shrubs.

**Woodland Understory Canopy Cover Summary (all species <13 feet in height)**

Understory plant type: Grass/grass-like (Graminoids)

Name	Symbol	Nativity	Cover low %	Cover high %	Canopy height bottom	Canopy height top
BIG BLUESTEM <i>Andropogon gerardii</i>	<a href="#">ANGE</a>	N	0.1	1.0	0.0	8.0
LITTLE BLUESTEM <i>Schizachyrium scoparium</i>	<a href="#">SCSC</a>	N	1.0	2.0		4.0
EASTERN BOTTLEBRUSH GRASS <i>Elymus hystrix</i>	<a href="#">ELHY</a>	N	0.1	1.0	0.0	3.0
VIRGINIA WILDRYE <i>Elymus virginicus</i>	<a href="#">ELVI3</a>	N	1.0	2.0		3.0
HAIRY WOODLAND BROME <i>Bromus pubescens</i>	<a href="#">BRPU6</a>	N	0.0	1.0	0.0	2.0
PARASOL SEDGE <i>Carex umbellata</i>	<a href="#">CAUM4</a>	N	1.0	2.0		2.0
HELLER'S ROSETTE GRASS <i>Dichanthelium oligosanthes</i>	<a href="#">DIOL</a>	N	0.1	1.0	0.0	2.0
HAIRY WILDRYE <i>Elymus villosus</i>	<a href="#">ELVI</a>	N	0.1	1.0	1.0	2.0

NODDING FESCUE							
<i>Festuca subverticillata</i>	<a href="#">FESU3</a>	N	0.1	1.0	0.0	2.0	
ROCK MUHLY							
<i>Muhlenbergia sobolifera</i>	<a href="#">MUJO</a>	N	0.1	1.0	0.0	2.0	
WOODLAND SEDGE							
<i>Carex blanda</i>	<a href="#">CABL</a>	N	0.0	0.1	0.0	1.0	
WOODBANK SEDGE							
<i>Carex cephalophora</i>	<a href="#">CACE</a>	N	0.1	1.0	0.0	1.0	
FUZZY-WUZZY SEDGE							
<i>Carex hirsutella</i>	<a href="#">CAHI6</a>	N	0.1	1.0	0.0	1.0	
MUHLENBERG'S SEDGE							
<i>Carex muehlenbergii</i>	<a href="#">CAMU4</a>	N	0.0	0.1	0.0	1.0	
PENNSYLVANIA SEDGE							
<i>Carex pennsylvanica</i>	<a href="#">CAPE6</a>	N	2.0	10.0	0.0	1.0	
BOSC'S PANICGRASS							
<i>Dichanthelium boscii</i>	<a href="#">DIBO2</a>	N	0.0	0.1	0.0	1.0	
RAVENEL'S ROSETTE GRASS							
<i>Dichanthelium ravenelii</i>	<a href="#">DIRA</a>	N	0.1	1.0	0.0	1.0	
POVERTY OATGRASS							
<i>Danthonia spicata</i>	<a href="#">DASP2</a>	N	0.0	5.0	0.0	0.5	
SLIMLEAF PANICGRASS							
<i>Dichanthelium linearifolium</i>	<a href="#">DILI2</a>	N	0.1	1.0	0.0	0.5	

*Understory plant type: Forb/Herb*

<u>Name</u>	<u>Symbol</u>	<u>Nativity</u>	<u>Cover low %</u>	<u>Cover high %</u>	<u>Canopy height bottom</u>	<u>Canopy height top</u>
BUTTON ERYNGO						
<i>Eryngium yuccifolium</i>	<a href="#">ERYU</a>	N	0.0	0.1	2.0	6.0
BALDWIN IRONWEED						
<i>Vernonia baldwinii</i>	<a href="#">VEBA</a>	N	0.1	1.0	0.0	4.0
PURPLE CONEFLOWER						
<i>Echinacea purpurea</i>	<a href="#">ECPU</a>	N	1.0	2.0		3.0
EASTERN BEEBALM						
<i>Monarda bradburiana</i>	<a href="#">MOBR2</a>	N	1.0	2.0		3.0
CANADIAN BLACKSNAKEROOT						
<i>Sanicula canadensis</i>	<a href="#">SACA15</a>	N	1.0	2.0		3.0
WIDOWSFRILL						
<i>Silene stellata</i>	<a href="#">SIST</a>	N	0.1	1.0	0.0	3.0
DOWNY AGRIMONY						
<i>Agrimonia pubescens</i>	<a href="#">AGPU</a>	N	0.1	1.0	0.0	2.0
HEMP DOGBANE						
<i>Apocynum cannabinum</i>	<a href="#">APCA</a>	N	0.0	0.1	0.0	2.0
SMOOTH BLUE ASTER						
<i>Aster laevis(syn)</i>	<a href="#">ASLA5</a>	N	1.0	2.0		2.0
SMOOTH YELLOW FALSE FOXGLOVE						
<i>Aureolaria flava</i>	<a href="#">AUFL</a>	N	0.1	1.0	1.0	2.0
TALL COREOPSIS						
<i>Coreopsis tripteris</i>	<a href="#">COTR4</a>	N	0.0	0.1	1.0	2.0

PANICLELEAF TICKTREFOIL <i>Desmodium paniculatum</i>	<a href="#">DEPA6</a>	N	0.1	2.0	0.0	2.0
PERPLEXED TICKTREFOIL <i>Desmodium perplexum</i>	<a href="#">DEPE80</a>	N	0.0	5.0	0.0	2.0
WHITE AVENS <i>Geum canadense</i>	<a href="#">GECA7</a>	N	0.0	0.1	0.0	2.0
HAIRY SUNFLOWER <i>Helianthus hirsutus</i>	<a href="#">HEHI2</a>	N	0.1	25.0	0.0	2.0
SLENDER LESPEDEZA <i>Lespedeza virginica</i>	<a href="#">LEVI7</a>	N	1.0	2.0		2.0
WILD QUININE <i>Parthenium integrifolium</i>	<a href="#">PAIN3</a>	N	0.0	0.1	0.0	2.0
NARROW-LEAF MOUNTAIN-MINT <i>Pycnanthemum tenuifolium</i>	<a href="#">PYTE</a>	N	0.0	0.1	1.0	2.0
CLUSTERED BLACKSNAKEROOT <i>Sanicula odorata</i>	<a href="#">SAOD</a>	N	0.1	1.0	0.0	2.0
WHOLELEAF ROSINWEED <i>Silphium integrifolium</i>	<a href="#">SIIN2</a>	N	0.0	0.1	0.0	2.0
BROADLEAF GOLDENROD <i>Solidago flexicaulis</i>	<a href="#">SOFL2</a>	N	0.1	1.0	0.0	2.0
ELMLEAF GOLDENROD <i>Solidago ulmifolia</i>	<a href="#">SOUL2</a>	N	5.0	10.0	0.0	2.0
DRUMMOND'S ASTER <i>Symphotrichum drummondii</i>	<a href="#">SYDR</a>	N	0.0	1.0	0.0	2.0
SMOOTH BLUE ASTER <i>Symphotrichum laeve</i>	<a href="#">SYLA3</a>	N	0.1	1.0	0.0	2.0
CALICO ASTER <i>Symphotrichum lateriflorum</i>	<a href="#">SYLA4</a>	N	0.1	1.0	0.0	2.0
WOODLAND AGRIMONY <i>Agrimonia rostellata</i>	<a href="#">AGRO3</a>	N	0.1	1.0	0.0	1.0
WESTERN ROCKJASMINE <i>Androsace occidentalis</i>	<a href="#">ANOC2</a>	N	0.0	1.0	0.0	1.0
VIRGINIA-SNAKEROOT <i>Aristolochia serpentaria</i>	<a href="#">ARSE3</a>	N	0.0	0.1	0.0	1.0
CANADA MILKVETCH <i>Astragalus canadensis</i>	<a href="#">ASCA11</a>	N	0.1	1.0	0.0	1.0
FOUR-LEAF MILKWEED <i>Asclepias quadrifolia</i>	<a href="#">ASQU</a>	N	0.0	0.1	0.0	1.0
FINGER COREOPSIS <i>Coreopsis palmata</i>	<a href="#">COPA10</a>	N	0.0	1.0	0.0	1.0
LARGE FLOWER TICKCLOVER <i>Desmodium glutinosum</i>	<a href="#">DEGL5</a>	N	0.1	2.0	0.0	1.0
NAKED-FLOWER TICK-TREFOIL <i>Desmodium nudiflorum</i>	<a href="#">DENU4</a>	N	10.0	25.0	0.0	1.0
FLOWERING SPURGE <i>Euphorbia corollata</i>	<a href="#">EUCO10</a>	N	0.1	1.0	0.0	1.0
WILD STRAWBERRY <i>Fragaria virginiana</i>	<a href="#">FRVI</a>	N	0.1	1.0	0.0	1.0
SHRUBBY LESPEDEZA <i>Lespedeza frutescens</i>	<a href="#">LEFR5</a>	N	1.0	50.0	0.0	1.0

FEATHERY FALSE SOLOMON'S-SEAL <i>Maianthemum racemosum</i>	<a href="#">MARA7</a>	N	0.1	1.0	0.0	1.0
WILD PETUNIA <i>Ruellia humilis</i>	<a href="#">RUHU</a>	N	0.0	0.1	0.0	1.0
BABY WHITE ASTER <i>Symphyotrichum anomalum</i>	<a href="#">SYAN2</a>	N	0.0	0.1	0.0	1.0
SMOOTH VIOLET PRAIRIE ASTER <i>Symphyotrichum turbinellum</i>	<a href="#">SYTU2</a>	N	0.0	0.1	0.0	1.0
RUE-ANEMONE <i>Thalictrum thalictroides</i>	<a href="#">THTH2</a>	N	0.1	1.0	0.0	1.0
BLUEJACKET <i>Tradescantia ohiensis</i>	<a href="#">TROH</a>	N	0.0	0.1	0.0	1.0
HOGPEANUT <i>Amphicarpaea bracteata</i>	<a href="#">AMBR2</a>	N	1.0	5.0	0.0	0.5
PARLIN'S PUSSYTOES <i>Antennaria parlinii</i>	<a href="#">ANPA9</a>	N	0.0	1.0	0.0	0.5
BASTARD TOADFLAX <i>Comandra umbellata</i>	<a href="#">COUM</a>	N	0.1	1.0	0.0	0.5
SHINING BEDSTRAW <i>Galium concinnum</i>	<a href="#">GACO3</a>	N	0.1	1.0	0.0	0.5
VIRGINIA STICKTIGHT <i>Hackelia virginiana</i>	<a href="#">HAVI2</a>	N	0.0	0.1	0.0	0.5
AMERICAN GINSENG <i>Panax quinquefolius</i>	<a href="#">PAQU</a>	N	0.1	1.0	0.0	0.5
WILD BLUE PHLOX <i>Phlox divaricata</i>	<a href="#">PHDI5</a>	N	0.1	1.0	0.0	0.5
LOPSEED <i>Phryma leptostachya</i>	<a href="#">PHLE5</a>	N	0.0	1.0	0.0	0.5
VIRGINIA GROUNDCHERRY <i>Physalis virginiana</i>	<a href="#">PHVI5</a>	N	0.0	1.0	0.0	0.5
GRAYHEAD PRAIRIE CONEFLOWER <i>Ratibida pinnata</i>	<a href="#">RAPI</a>	N	0.0	0.1	0.0	0.5

Understory plant type: Shrub/Subshrub

<u>Name</u>	<u>Symbol</u>	<u>Nativity</u>	<u>Cover low %</u>	<u>Cover high %</u>	<u>Canopy height bottom</u>	<u>Canopy height top</u>
HAZELNUT <i>Corylus americana</i>	<a href="#">COAM3</a>	N	0.1	2.0	0.0	8.0
FRAGRANT SUMAC <i>Rhus aromatica</i>	<a href="#">RHAR4</a>	N	2.0	50.0	0.0	8.0
NEW JERSEY TEA <i>Ceanothus americanus</i>	<a href="#">CEAM</a>	N	5.0	20.0		3.0
BUCKBRUSH <i>Symphoricarpos orbiculatus</i>	<a href="#">SYOR</a>	N	0.1	2.0	0.0	3.0
STIFF DOGWOOD <i>Cornus foemina</i>	<a href="#">COFO</a>	N	0.0	0.1	0.0	2.0
DEWBERRY <i>Rubus flagellaris</i>	<a href="#">RUFL</a>	N	0.1	2.0	0.0	2.0

BLACKHAW							
<i>Viburnum prunifolium</i>	<a href="#">VIPR</a>	N	0.0	0.1	0.0	2.0	
SHRUBBY ST. JOHNSWORT							
<i>Hypericum prolificum</i>	<a href="#">HYPR</a>	N	2.0	5.0	0.0	1.0	
CAROLINA ROSE							
<i>Rosa carolina</i>	<a href="#">ROCA4</a>	N	0.0	0.1	0.0	1.0	

*Understory plant type: Tree*

<u>Name</u>	<u>Symbol</u>	<u>Nativity</u>	<u>Cover low %</u>	<u>Cover high %</u>	<u>Canopy height bottom</u>	<u>Canopy height top</u>
EASTERN REDCEDAR						
<i>Juniperus virginiana</i>	<a href="#">JUVI</a>	N	0.0	0.1	0.0	16.0
EASTERN HOPHORNBEAM						
<i>Ostrya virginiana</i>	<a href="#">OSVI</a>	N	0.1	25.0	0.0	16.0
BLACK HICKORY						
<i>Carya texana</i>	<a href="#">CATE9</a>	N	0.1	1.0	0.0	8.0
JUNEBERRY						
<i>Amelanchier arborea</i>	<a href="#">AMAR3</a>	N	0.1	1.0	0.0	2.0
MOCKERNUT HICKORY						
<i>Carya alba(syn)</i>	<a href="#">CAAL27</a>	N	0.1	1.0	0.0	2.0
PIGNET HICKORY						
<i>Carya cordiformis</i>	<a href="#">CACO15</a>	N	0.0	0.1	0.0	2.0
PERSIMMON						
<i>Diospyros virginiana</i>	<a href="#">DIVI5</a>	N	0.1	1.0	0.0	2.0
BLACK CHERRY						
<i>Prunus serotina</i>	<a href="#">PRSE2</a>	N	0.1	1.0	0.0	2.0
SHINGLE OAK						
<i>Quercus imbricaria</i>	<a href="#">QUIM</a>	N	0.0	2.0	0.0	2.0

*Understory plant type: Vine*

<u>Name</u>	<u>Symbol</u>	<u>Nativity</u>	<u>Cover low %</u>	<u>Cover high %</u>	<u>Canopy height bottom</u>	<u>Canopy height top</u>
POISON IVY						
<i>Toxicodendron radicans</i>	<a href="#">TORA2</a>	N	0.1	2.0	0.0	16.0
BITTERSWEET						
<i>Celastrus scandens</i>	<a href="#">CESC</a>	N	0.0	1.0	0.0	1.0
VIRGINIA CREEPER						
<i>Parthenocissus quinquefolia</i>	<a href="#">PAQU2</a>	N	1.0	5.0	0.0	1.0
WINTER GRAPE						
<i>Vitis vulpina</i>	<a href="#">VIVU</a>	N	0.0	1.0	0.0	1.0

**Community Phase 1.2: White Oak-Black Oak-Hickory/Eastern Hop Hornbeam – Aromatic Sumac/Woodland Brome – Hairy Sunflower**

(Photo below: White oak dominated reference site at Rebels Cove Conservation Area, Schuyler County, MO - photo from MDC)



This phase is similar to community phase 1.1 but oak and hickory densities are increasing due to longer periods of fire suppression. Displacement of some grasses and forbs (see species lists below) may be occurring due to shading and competition from the increased densities of oak and hickory saplings in the understory.

**Community Phase Pathway 1.2A**

This pathway results from disturbance intervals such as fire that are increasing and occur more than once every 10 years. Over time, these disturbance events result in a community phase transition back to the

phase 1.1.

**Woodland Overstory**

The overstory of this community phase is a mixture of white oak and black oak, with increasing densities of hickory and other more shade tolerant species.

**Woodland Overstory Characterization Summary (tree and vine species >13 feet in height)**

Woodland canopy: Low canopy cover % 60 RV canopy cover % 70 High canopy cover % 90

Overstory plant type: Tree

Name	Symbol	Nativity	Cover low %	Cover high %	Canopy height bottom	Canopy height top
BLACK OAK <i>Quercus velutina</i>	<a href="#">QUVE</a>	N	20.0	40.0		80.0
MOCKERNUT HICKORY <i>Carya tomentosa</i>	<a href="#">CATO6</a>	N	10.0	30.0		70.0
WHITE OAK <i>Quercus alba</i>	<a href="#">QUAL</a>	N	40.0	70.0		70.0
BUR OAK <i>Quercus macrocarpa</i>	<a href="#">QUMA2</a>	N	5.0	10.0		70.0
SHAGBARK HICKORY <i>Carya ovata</i>	<a href="#">CAOV2</a>	N	10.0	30.0		60.0
POST OAK <i>Quercus stellata</i>	<a href="#">QUST</a>	N	5.0	10.0		60.0

**Woodland Understory**

This woodland community has a multi-tiered structure due to lack of disturbance activities. Ground cover densities are decreasing due increasing overstory cover.

**Woodland Understory Canopy Cover Summary** (all species <13 feet in height)

*Understory plant type: Grass/grass-like*

Name	Symbol	Nativity	Cover low %	Cover high %	Canopy height bottom	Canopy height top
BIG BLUESTEM <i>Andropogon gerardii</i>	<a href="#">ANGE</a>	N	0.1	2.0		5.0
LITTLE BLUESTEM <i>Schizachyrium scoparium</i>	<a href="#">SCSC</a>	N	0.1	2.0		4.0
HAIRY WOODLAND BROME <i>Bromus pubescens</i>	<a href="#">BRPU6</a>	N	0.1	2.0		3.0
EASTERN BOTTLEBRUSH GRASS <i>Elymus hystrix</i>	<a href="#">ELHY</a>	N	0.1	2.0		3.0
VIRGINIA WILD RYE <i>Elymus virginicus</i>	<a href="#">ELVI3</a>	N	0.1	5.0		3.0
FOUR-LEAF MILKWEED <i>Asclepias quadrifolia</i>	<a href="#">ASQU</a>	N	0.1	2.0		2.0
PENNSYLVANIA SEDGE <i>Carex pensylvanica</i>	<a href="#">CAPE6</a>	N	0.1	2.0		2.0
PARASOL SEDGE <i>Carex umbellata</i>	<a href="#">CAUM4</a>	N	0.1	2.0		2.0
ROCK MUHLY <i>Muhlenbergia sobolifera</i>	<a href="#">MUSO</a>	N	0.1	5.0		2.0

*Understory plant type: Forb/Herb*

Name	Symbol	Nativity	Cover low %	Cover high %	Canopy height bottom	Canopy height top
CULVER'S-ROOT <i>Veronicastrum virginicum</i>	<a href="#">VEVI4</a>	N	0.1	1.0		4.0
PURPLE CONEFLOWER <i>Echinacea purpurea</i>	<a href="#">ECPU</a>	N	0.1	2.0		3.0
HAIRY SUNFLOWER <i>Helianthus hirsutus</i>	<a href="#">HEHI2</a>	N	0.1	5.0		3.0
EASTERN BEEBALM <i>Monarda bradburiana</i>	<a href="#">MOBR2</a>	N	0.1	2.0		3.0
CANADIAN BLACKSNAKEROOT <i>Sanicula canadensis</i>	<a href="#">SACA15</a>	N	0.1	2.0		3.0
ELMLEAF GOLDENROD <i>Solidago ulmifolia</i>	<a href="#">SOUL2</a>	N	1.0	5.0		3.0
SMOOTH BLUE ASTER <i>Aster laevis(syn)</i>	<a href="#">ASLA5</a>	N	0.1	2.0		2.0
NAKED-FLOWER TICK-TREFOIL <i>Desmodium nudiflorum</i>	<a href="#">DENU4</a>	N	0.1	2.0		2.0
SLENDER LESPEDEZA <i>Lespedeza virginica</i>	<a href="#">LEVI7</a>	N	0.1	2.0		2.0
BLUEJACKET <i>Tradescantia ohiensis</i>	<a href="#">TROH</a>	N	0.1	5.0		2.0

*Understory plant type: Shrub/Subshrub*

Name	Symbol	Nativity	Cover	Cover	Canopy	Canopy
			low %	high %	height bottom	height top
HAZELNUT <i>Corylus americana</i>	<a href="#">COAM3</a>	N	1.0	2.0		4.0
NEW JERSEY TEA <i>Ceanothus americanus</i>	<a href="#">CEAM</a>	N	1.0	2.0		3.0
FRAGRANT SUMAC <i>Rhus aromatica</i>	<a href="#">RHAR4</a>	N	1.0	5.0		3.0

*Understory plant type: Tree*

Name	Symbol	Nativity	Cover	Cover	Canopy	Canopy
			low %	high %	height bottom	height top
EASTERN HOPHORNBEAM <i>Ostrya virginiana</i>	<a href="#">OSVI</a>	N	1.0	5.0		40.0
WHITE ASH <i>Fraxinus americana</i>	<a href="#">FRAM2</a>	N	0.1	1.0		20.0
SLIPPERY ELM <i>Ulmus rubra</i>	<a href="#">ULRU</a>	N	1.0	2.0		20.0

Transition T1A

Fire suppression and Woodland management will result in a transition to community phase 2.1.

Transition T1B

Fire suppression, Woodland management and harvesting will result in a transition to community phase 3.1.

Transition 1C

Clearing, pasture planting and grassland management will result in a transition to community phase 4.1.

Transition 1D

High grade logging and uncontrolled grazing will result in a transition to community phase 5.1.

**State 2: Timber Managed Woodland**

This state develops from mixed oak woodlands, which over time, increase in canopy cover, eventually transitioning into forest-like communities. These woodlands tend to be rather dense, with a higher developed understory but a more under developed ground flora. Thinning can increase overall tree vigor and improve and control understory density. Continual timber management, depending on the practices used, should maintain this state. Over time white oak will become less dominant with black oak increasing. With cessation of harvesting and no other management inputs this community phase will slowly increase in more shade tolerant species such as hickories, white ash and maple and transition to community phase 2.2. Over time white oak may become less dominant. Re-initiation of harvesting and other forest management activities will transition this community phase back to community phase 2.1. Repeated, high-grading timber harvests coupled with introduction of uncontrolled domestic livestock grazing will transition this state to state 5, High-Graded/Grazed Woodland.

This state can be restored to a reference state by modifying or eliminating timber harvests, extending rotations, incorporating selective thinning, and re-introducing prescribed fire. (See Ecological Dynamics section for caution on sites with invasive non-native species of earthworms)

**Community Phase 2.1: Black Oak – White Oak /Serviceberry – Hazelnut / Aster – Pennsylvania Sedge**  
 (Photo below: Timber managed site with recent harvest (notice stump in foreground) - photo from MDC)

These woodlands tend to be rather dense, with a higher developed understory but a more under developed ground flora. Thinning can increase overall tree vigor and improve and control understory density. Continual timber management, depending on the practices used, should maintain this state. Over time white oak will become less dominant. Canopy cover ranges from 70 to 85 percent. Most stands are 50 to 80 years of age.



**Community Phase Pathway 2.1A**

This pathway results in a cessation or reduction of harvesting frequencies.

**Woodland Overstory**

Black oak and white oak are common overstory species along with hickory. Canopy levels can approach 90 percent.

**Woodland Understory**

A multi-layered understory is present. Due to increased shading, the ground flora is somewhat under developed.

**Community Phase 2.2: Black Oak – White Oak - Hickory/Serviceberry - Maple Saplings/ Aster – Pennsylvania Sedge**

With cessation of harvesting and no other management inputs this community phase will slowly increase in more shade tolerant species such as hickories, white ash and maple. Over time white oak will become less dominant.

**Community Phase Pathway 2.2A**

Re-initiation of harvesting and other Woodland management activities will transition this community phase back to community phase 2.1.

**Transition 2A**

This state will transition to a fire managed woodland state by incorporating selective thinning and re-introducing prescribed fire.

**Transition 2B**

This state will transition to a high-graded/grazed woodland state with high-grade logging and allowing uncontrolled grazing.

**Restoration Pathway 1B**

This state can be restored to a reference state by modifying or eliminating timber harvests, extending rotations, incorporating selective thinning, and re-introducing prescribed fire.

**State 3: Fire Managed Woodland**

The fire managed woodland state most closely resembles the reference state. The principal difference is tree age with most canopy trees being only 50 to 90 years old. Composition and reduction in tree size are also likely altered from the reference state depending on tree selection during any previous harvest cycles.

Without a regular fire regime or harvesting, this community phase will slowly increase in more shade tolerant species such as hickories and other more shade tolerant species and transition to community phase 3.2. Over time white oak may become less dominant. Repeated, high-grading timber harvests coupled with introduction of uncontrolled domestic livestock grazing and cessation of prescribed fire will transition this state to state 5, High-Graded/Grazed Woodland.

This state can be restored to a reference state by modifying timber harvests, extending rotations, incorporating selective thinning, and possibly increasing fire frequencies. (See Ecological Dynamics section for caution on sites with invasive non-native species of earthworms)

**Community Phase 3.1: Black Oak – White Oak – Hickory/ Buckbrush/ Woodland Brome**



*(Photo on left: Fire managed woodland at Union Ridge Conservation Area near Green City, Missouri - photo from MDC)*

This is a fire managed phase. Fire frequencies occur at a 5 to 10 year interval usually preceding a timber harvest. Logging activities are removing higher volumes of white oak causing a decrease in white oak in the canopy and an increase in black oak.

**Community Phase Pathway 3.1A**  
This pathway results in reduced fire frequencies and logging activity. No disturbance periods can exceed 10 years.

**Woodland Overstory**

Black oak and white oak dominate with occasional hickories. Canopy cover ranges from 60 percent to 80 percent.

**Woodland Understory**

Dense rich ground cover dominated by forbs and shrubs.

**Community Phase 3.2: Black Oak – White Oak – Hickory/Oak Saplings – Black Cherry/ Woodland Brome**

In the long term absence of fire, woody species, especially oak and hickory saplings, black cherry and hornbeam encroach into the understory of these woodlands. Hickory species will increase in the overstory.

**Community Phase Pathway 3.2A**

This pathway results in increased prescribed fire frequencies and the reinstating of selective thinning.

**Transition 3A**

This state will transition to a timber managed state by modifying timber harvests and suppressing prescribed fire.

**Transition 3B**

High-grade harvesting and introduction of uncontrolled grazing will cause a transition to community phase 5.1

**Restoration Pathway 1A**

This state can be restored to a reference state by maintaining prescribed fires, modifying or eliminating timber harvests, extending rotations, and incorporating selective thinning.

**State 4: Grassland**

Conversion of Woodlands to planted, non-native cool season grasses and legumes has been common. Without proper grassland management these ecological sites are challenging to maintain in a healthy, productive state.

With over grazing and cessation of active pasture management, tall fescue, white clover and multi-flora rose will increase in density. Over a long period of time with no grazing, phase 4.2 will eventually transition to a phase that resembles community phase 5.2, Black Oak-Hickory/Black Cherry-Buckbrush-Gooseberry- Multi-flora Rose/ Black Snakeroot - Geranium.

In some instances, this state has been converted to native warm season grasses, primarily big bluestem, switchgrass, and Indian grass or pure stands of single species. These sites are typically converted through a federal cost share program such as the Conservation Reserve Program (CRP) or the Environmental Quality Incentives Program (EQIP). Some sites are associated with an active rotational grazing system. Without active management with prescribed fire and grassland management these sites will transition to phase 5.2.

**Community Phase 4.1: Cool Season Grasses – Legumes**

This phase is well-managed grassland, composed of non-native cool season grasses and legumes. Grazing and haying is occurring. *(Photo on left: Well-*



*managed cool season pasture near Lake Wapello State Park, Iowa - photo from MDC).* The effects of long-term liming on soil pH, and calcium and magnesium content, is most evident in this phase. Studies show that these soils have higher pH and higher base status in soil horizons as much as two feet below the surface, relative to poorly managed grassland (phase 4.2) and to woodland communities (where liming is not practiced).

**Community Phase Pathway 4.1A**

This pathway results from over grazing and cessation of active pasture management.

**Community Phase 4.2: Tall fescue – White Clover/Multi-flora Rose**

*(Photo below: Cool season pasture showing weedy invasion due to poor management near Bloomfield, Iowa - photo from MDC)*



This phase is the result of poor grassland management. Over grazing and inadequate or no fertility application has allowed tall fescue, multi-flora rose, thistle and other weedy species to increase in cover and density reducing overall forage quality and site productivity. White clovers such as ladino and alsike will decrease or go away with no fertilization and overgrazing although Dutch white clover will leave last. Soil pH and bases such as calcium and magnesium are lower, relative to well-managed pastures (Phase 4.1).

**Community Phase Pathway 4.2A**

To return to Community Phase 4.1, requires brush management, grassland seeding, rotational grazing, and integrated pest management.

*Community Phase 4.3: Native Warm Season Grasses*

In some instances, this state has been converted to native warm season grasses, primarily big bluestem, switchgrass, and Indian grass or pure stands of single species. These sites are typically converted through a federal cost share program such as the Conservation Reserve Program (CRP) or the Environmental Quality Incentives Program (EQIP). Some sites are associated with an active rotational grazing system.



*(Photo on left: Native grasses and legumes on CRP land in southern Iowa - photo from NRCS)*

**Transition 4A**

This state will transition to a high-graded/grazed woodland idle phase with long term succession allowing woody species to become established and little to no grazing.

**State 5: High-Graded/Grazed Woodland**

Reference or managed wooded states subjected to repeated, high-grading timber harvests and uncontrolled cattle grazing transition to this degraded state. This state exhibits an over-abundance of hickory and other less economically desirable tree species and weedy understory species such as buckbrush, gooseberry, poison ivy and multi-flora rose. The vegetation offers little nutritional value for cattle, and excessive livestock stocking damages tree boles, degrades understory species composition and results in soil compaction and accelerated erosion and runoff. Browsing by goats using good rotational management can open up the shrub layer, eliminate many of the weedy species and increase both native herbaceous vegetation and may induce regeneration of oak and hickory species. Cessation of active logging and exclusion of livestock from sites in this state will create an idle phase that experiences an increase in black cherry and Ohio buckeye in the understory layer.

Transition back to either an even-age managed or uneven-age managed woodland will required dynamic and sustained forest stand improvements, cessation of grazing, and selective thinning of overstory and understory canopies. (See Ecological Dynamics section for caution on sites with invasive non-native species of earthworms)

*Community Phase 5.1: Black Oak-Hickory/Hickory saplings-Gooseberry-Multiflora Rose/Black Snakeroot-Geranium*



Due to high-grade logging and uncontrolled grazing, this community phase exhibits an over-abundance of hickory and other less economically desirable tree species and weedy understory species such as buckbrush, gooseberry, poison ivy and multi-flora rose. The understory vegetation offers little nutritional value for cattle, and excessive livestock stocking damages tree boles, degrades understory species composition and results in soil compaction and accelerated erosion and runoff.

*(Photo on left: Actively grazed woodland creating an open understory - photo from NRCS)*

**Community Phase Pathway 5.1A**

This pathway results from cessation of active logging and periodic exclusion of livestock or reduction in grazing intensity.

**Woodland Overstory**

Canopy exhibits an over-abundance of hickory and other less economically desirable tree species such as black oak.

**Woodland Understory**

Over time, understory species such as buckbrush, gooseberry, poison ivy and multi-flora rose increase in density.

*Community Phase 5.2: Black Oak-Hickory/Black Cherry-Buckbrush-Gooseberry-Multiflora Rose/Black Snakeroot-Geranium*



Cessation of active logging and reduction in grazing intensity will create an idle phase that experiences an increase in black cherry and Ohio buckeye and weedy species such as buckbrush and gooseberry in the understory layer.

*(Photo on left: Lamson Woods State Preserve showing high-graded formerly grazed woodland near Fairfield, Iowa - photo from MDC)*

*Community Phase Pathway 5.2A*

This pathway results in a return to unrestricted logging activity and increased grazing intensities and frequencies.

**Woodland Overstory**

Canopy exhibits an over-abundance of hickory and other less economically desirable tree species such as black oak.

**Woodland Understory**

Understory densities levels are increasing. Species such as black cherry, Ohio buckeye, buckbrush, gooseberry, poison ivy and multi-flora rose are common.

Transition 5A

This state will transition to a grassland state with clearing, pasture planting, and grassland management.

Transition 5B

This state will transition to a fire managed woodland state with timber management, forest stand improvement, suppression of grazing and prescribed burning.

Transition 5C

This state will transition to a timber managed woodland state with timber management, forest stand improvement, and suppression of grazing.

## Section II: Ecological Site Interpretations

### Forest Site Productivity

<u>Common Name</u>	<u>Symbol</u>	<u>Site</u>	<u>Site</u>	<u>CMAI</u>	<u>CMAI</u>	<u>Age of</u>	<u>Site</u>	<u>Site</u>	<u>Citation</u>
		<u>Index</u>	<u>Index</u>				<u>Index</u>	<u>Index</u>	
		<u>Low</u>	<u>High</u>	<u>Low</u>	<u>High</u>	<u>CMAI</u>	<u>Curve</u>	<u>Curve</u>	
WHITE OAK	<a href="#">QUAL</a>	48	60	32	46	50	820	50TA	Schnur, G. Luther. 1937. Yield, stand, and volume tables for even-aged upland oak forests. United States Department of Agriculture Technical Bulletin 560
BLACK OAK	<a href="#">QUVE</a>	42	73	32	48	50	820	50TA	Schnur, G. Luther. 1937. Yield, stand, and volume tables for even-aged upland oak forests. United States Department of Agriculture Technical Bulletin 560

### Animal Community

#### *Wildlife:*

Wild turkey, white-tailed deer, and eastern gray squirrel depend on hard and soft mast food sources and are typical upland game species of this type. (Pitts and McGuire 2000; Schwartz, et al. 2001)

Oaks provide hard mast; scattered shrubs provide soft mast; native legumes provide high-quality wildlife food.

Sedges and native cool-season grasses provide green browse; patchy native warm-season grasses provide cover and nesting habitat; and a diversity of forbs provides a diversity and abundance of insects.

Post-burn areas can provide temporary bare-ground – herbaceous cover habitat important for turkey poults and quail chicks.

Bird species associated with mature communities include Indigo Bunting, Red-headed Woodpecker, Eastern Bluebird, Northern Bobwhite, Eastern Wood-Pewee, Broad-winged Hawk, Great-Crested Flycatcher, Summer Tanager, and Red-eyed Vireo (Fitzgerald and Pashley 2000b; Jacobs 2001).

Reptiles and amphibians associated with these Woodlands include: ringed salamander, spotted salamander, marbled salamander, central newt, long-tailed salamander, dark-sided salamander, southern red-backed salamander, small-mouthed salamander, three-toed box turtle, ground skink, western worm snake, western earth snake, American toad, and eastern timber rattlesnake (Johnson 2000).

#### *Domestic livestock:*

Currently, domestic cattle grazing is occurs on sites but concerns with lack of water, access issues, and high amounts of acorns on the ground in the fall that are detrimental to cattle, limits its usefulness. On sites that have been cleared of trees and converted to cool-season grasses cattle grazing is common.

### Hydrology Functions

Most precipitation on sites in reference or well-managed timber states infiltrates the soil, and either recharges the local groundwater or moves slowly as lateral flow, surfacing in headwaters of ephemeral streams. The trees, the shrub and herbaceous understories, and the litter provide nearly 100 percent soil cover. Little or no surface runoff occurs on these sites, except for rare, high-intensity storms. These sites provide high yields of good-quality groundwater, which is released slowly into ephemeral streams over time.

In high graded woodland or poorly managed pasture states, soil compaction and reduced surface cover generally results in reduced infiltration and increased runoff. Groundwater recharge is reduced. Surface runoff results in soil

erosion, which degrades water quality. The rapid release of runoff into ephemeral streams increases the risk of downstream flooding, and shortens the time when ephemeral streams are active.

**Recreational Uses**

Hunting, bird watching, horseback riding, camping, and hiking are recreational uses of this ecological site. Reference and well managed sites provide good hunting for turkey, white-tailed deer, and squirrel. Recreational uses are reduced in the heavily grazed grassland state and high-graded woodland state. In many areas of this predominantly agricultural MLRA, these sites provide the only Woodlands available for recreational use.

**Wood Products**

This ecological site is productive. Timber harvesting can occur but care should be taken to maintain the integrity and character of the site. Potential products include lumber, oak staves, pallet materials, and in some cases oak and walnut veneer (only on well managed or old growth sites).

**Other Information**

Woodland management: Site index values range from 50 to 60 for oak. Timber management opportunities are good. These groups respond well to even-aged management. Create group openings of at least 2 acres. Large clearcuts should be minimized if possible to reduce impacts on wildlife and aesthetics. Uneven-aged management using single tree selection and group selection cuttings of ½ to 1 acre are other options that can be used if clear cutting is not desired or warranted. Prescribed fire is an effective management tool. Using prescribed fire as a management tool however, could have a negative impact on timber quality and should be used with caution on a particular site if timber management is the primary objective. Favor white oak, post oak, chinkapin oak, and black oak.

**Supporting Information**

**Associated Sites**

<u>Site name</u>	<u>Site ID</u>	<u>Site narrative</u>
Loess Upland Woodland	F109XY003MO	Loess Upland Woodlands are often upslope from Till Exposed Backslope Woodlands.
Till Upland Woodland	F109XY007MO	Till Upland Woodlands are often upslope from Till Exposed Backslope Woodlands.
Shale Exposed Backslope Woodland	F109XY013MO	Shale Exposed Backslope Woodlands are downslope from Till Exposed Backslope Woodlands in some places.
Till Protected Backslope Forest	F109XY022MO	Till Protected Backslope Forests are mapped in complex with the Till Exposed Backslope Woodlands, on northerly and easterly aspects

**Similar Sites**

<u>Site name</u>	<u>Site ID</u>	<u>Site narrative</u>
Till Upland Woodland	F109XY007MO	Till Upland Woodlands are on upper slopes and shoulders, and are not as steep. Canopy composition is similar to Till Exposed Backslope Woodlands but more open with a ground flora having a higher grass density grass.
Till Backslope Savanna	R109XY008MO	Till Backslope Savannas have thicker, darker surface layers from historic grassland vegetation. Bur oak and post oak dominated the much more open canopy. Ground flora was similar to adjacent prairie site and fire frequencies were higher.
Shale Exposed Backslope Woodland	F109XY025MO	Shale Exposed Backslope Woodlands are downslope from Till Exposed Backslope Woodlands in some places. Canopy composition and structure are similar to Till Exposed Backslope Woodlands. Fire frequency was generally less.

**State Correlation**

This site has been correlated with the following states: IA MO

**Hierarchical Classification Relationships**

Atlas of Missouri Ecoregions (Nigh & Schroeder, 2002): This ecological site occurs in many Land Type Associations within the following Subsections:

- Chariton River Hills
- Claypan Till Plains
- Mississippi River Hills
- Wyaconda River Dissected Till Plains

Terrestrial Natural Community Type in Missouri (Nelson, 2010):

The reference state for this ecological site is most similar to a Dry-Mesic Loess/Glacial Till Forest.

Missouri Department of Conservation Forest and Woodland Communities (Missouri Department of Conservation, 2006): The reference state for this ecological site is most similar to White Oak Loess/Glacial Till Forest.

National Vegetation Classification System Vegetation Association (NatureServe, 2010): The reference state for this ecological site is within the North-Central Interior Dry-Mesic Oak Forest and Woodland (CES202.046), and is most similar to *Quercus alba* - (*Carya ovata*)/ *Carex pensylvanica* Glaciated Woodland (CEGL002134).

**Data Source of ECS-5 plots (for Lindley, Keswick and Winnegan soils)**

<u>Data source</u>	<u>Number</u>	<u>Year</u>	<u>State code</u>	<u>County code</u>	<u>State</u>	<u>County</u>
ECS-5	2	1990	29	01	Missouri	Adair
	7	1990	29	01	Missouri	Adair
	8	1990	29	01	Missouri	Adair
	2	1964	29	019	Missouri	Boone
	10	1984	29	027	Missouri	Callaway
	8	1984	29	027	Missouri	Callaway
	8	1991	29	041	Missouri	Chariton
	11	1984	29	113	Missouri	Lincoln
	11	1984	29	115	Missouri	Linn
	12	1984	29	115	Missouri	Linn
	8	1988	29	121	Missouri	Macon
	1	1975	29	127	Missouri	Marion
	1	1983	29	175	Missouri	Randolph
	2	1983	29	175	Missouri	Randolph
	7	1985	29	195	Missouri	Saline
	9	1990	29	211	Missouri	Sullivan

## Section III: Inventory Data Collection

The data contained in this document is derived from analysis of inventories, ecological interpretation from field evaluations, and various reference papers and books. Destructive plant sampling was not allowed on the public reference sites. Site index information on woody species was collected to provide estimates of site productivity.

### Inventory Data References

Site concept was developed from Paul Nelson's Terrestrial Natural Communities and other works and refined with Amber Steele's thesis.

- Steele, Amber M.; Kabrick, John M.; Miles, Randall J. 2013. Regional and geomorphic influence on the productivity, composition, and structure of oak ecosystems in the western central hardwoods region. In: Miller, Gary W.; Schuler, Thomas M.; Gottschalk, Kurt W.; Brooks, John R.; Grushecky, Shawn T.; Spong, Ben D.; Rentch, James S., eds. Proceedings, 18th Central Hardwood Forest Conference; 2012 March 26-28; Morgantown, WV; Gen. Tech. Rep. NRS-P-117. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 80-92.
- Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.
- Yatskievych, George A. 1999/2006/2013. Flora of Missouri. Missouri Dept. of Conservation in cooperation with Missouri Botanical Garden Press, Volumes 1-3.

### Sampling methods (nested plots/transects/releve)

#### *Level 2 and reconnaissance inventory:*

- 2007: Kolaks/Meinert reconnaissance at Little Lost Creek CA and Rudolph Bennitt CA
- 2009/10: Steele/Steele/Kabrick reconnaissance for Till Backslope thesis study site selection (Atlanta CA, Hidden Hollow CA, Hungry Mother CA, Rebel's Cove CA, Union Ridge CA, Sugar Creek CA, others)
- 2014: Reconnaissance in Iowa: (Lamson Woods, Lacey Keosaqua SP. others)
- 2014: Reconnaissance and Tier III plot establishment in Missouri (Dark Hollow CA, Mineral Hills CA, Union Ridge CA, Thousand Hills SP, Rudolph Bennitt CA)

#### *Reference Inventory Plots:*

- THHISP02 Thousand Hills State Park
- DAHOCA02 Dark Hollow Natural Area
- RUBECA02 Rudolf Bennitt Wildlife Conservation Area

#### *Timber Managed Woodland State Inventory Plots:*

- RECOCA01 Rebel's Cove Conservation Area
- HNMOCA02 Hungry Mother Conservation Area

#### *Fire Managed Woodland State Inventory Plots:*

- UNRICA03 Union Ridge Conservation Area

### Type Localities for Reference State Data Plots:

<b>State:</b>	<b>MO</b>
County:	Adair
Township:	62N
Range:	15W
Section:	18
Datum:	NAD83
Zone:	15
Northing:	4447630.34

*Easting:* 531737.44  
*General legal description:* Plot THHISP02 Thousand Hills State Park  
*Latitude degrees:* 40  
*Latitude minutes:* 10  
*Latitude seconds:* 42  
*Latitude decimal:* 42  
*Longitude degrees:* 92  
*Longitude minutes:* 37  
*Longitude seconds:* 37  
*Longitude decimal:* 99  
*Universal Transverse Mercator (UTM) system:* NAD83154447630.34531737.44  
**State:** **MO**  
*County:* Randolph  
*Township:* 52N  
*Range:* 14W  
*Section:* 36  
*Datum:* NAD83  
*Zone:* 15  
*Northing:* 4345191.15  
*Easting:* 547626.93  
*General legal description:* Plot RUBECA02 Rudolf Bennitt Wildlife Conservation Area  
*Latitude degrees:* 39  
*Latitude minutes:* 15  
*Latitude seconds:* 17  
*Latitude decimal:* 6  
*Longitude degrees:* 92  
*Longitude minutes:* 26  
*Longitude seconds:* 52  
*Longitude decimal:* 80  
*Universal Transverse Mercator (UTM) system:* NAD83154345191.15547626.93  
**State:** **MO**  
*County:* Sullivan  
*Township:* 64N  
*Range:* 18W  
*Section:* 28  
*Datum:* NAD83  
*Zone:* 15  
*Northing:* 4463745.93  
*Easting:* 506076.80  
*General legal description:* Plot DAHOCA01 Dark Hollow Natural Area Winnegan pedon  
*Latitude degrees:* 40  
*Latitude minutes:* 19  
*Latitude seconds:* 27  
*Latitude decimal:* 19  
*Longitude degrees:* 92  
*Longitude minutes:* 55  
*Longitude seconds:* 42  
*Longitude decimal:* 49  
*Universal Transverse Mercator (UTM) system:* NAD83154463745.93506076.80

## References

- Ecological Site Information System: Soils Ecological Site Inventory-Forestland. Data (Keswick, Lindley, and Winnegan soils) accessed June 2013. Retrieved from [https://esi.sc.egov.usda.gov/ESI\\_Forestland/pgFSWelcome.aspx](https://esi.sc.egov.usda.gov/ESI_Forestland/pgFSWelcome.aspx).
- Festervand, D.F. 1994. Soil Survey of Putnam County, Missouri. U.S. Dept. of Agric. Soil Conservation Service.
- Fitzgerald, J.A. and D.N. Pashley. 2000b. Partners in Flight bird conservation plan for the Dissected Till Plains. American Bird Conservancy.
- Frost, C., 1996. Pre-settlement Fire Frequency Regimes of the United States: A First Approximation. Pages 70-81, Proceedings of the 20nd Tall Timbers Fire Ecology Conference: Fire in Ecosystem Management: Shifting the Paradigm from Suppression to Prescription. Tall Timbers Research Station, Tallahassee, FL.
- Hendrix, P. F., G. H. Baker, M. A. Callahan Jr, G. A. Damoff, C. Fragoso, G. Gonzalez, S. W. James, S. L. Lachnicht, T. Winsome and X. Zou. 2006. Invasion of exotic earthworms into ecosystems inhabited by native earthworms. *Biol Invasions* (2006) 8:1287–1300
- Jacobs, B. 2001. Birds in Missouri. Missouri Department of Conservation, Jefferson City, Missouri.
- Johnson, T.R. 2000. The amphibians and reptiles of Missouri. 2nd ed. Missouri Department of Conservation, Jefferson City, Missouri.
- Maerz, J., V. A. Nuzzo, and B. Blossey. 2009. Declines in Woodland Salamander Abundance Associated with Non-Native Earthworm and Plant Invasions. *Conservation Biology*, Volume 23, No. 4, 975–981
- Missouri Department of Conservation, 2006. Missouri Forest and Woodland Community Profiles. Jefferson City, Missouri.
- NatureServe. 2010. Vegetation Associations of Missouri (revised). NatureServe, St. Paul, Minnesota.
- Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri. 550p.
- Nigh, Timothy A., and Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.
- Nuzzo, V., J. C. Maerz, and B. Blossey. 2009. Earthworm invasion as the Driving Force Behind Plant Invasion and Community Change in Northeastern North American Forests. *Conservation Biology* 23: No. 4, 966–974.
- Pitts, D.E. and W.D. McGuire. 2000. Wildlife management for Missouri landowners. 3rd ed. Missouri Department of Conservation, Jefferson City, Missouri.
- Schwartz, C.W., E.R. Schwartz and J.J. Conley. 2001. The wild mammals of Missouri. University of Missouri Press, Columbia and Missouri Department of Conservation, Jefferson City, Missouri.
- Steele, Amber M. 2011. Regional and geomorphic influence on soil genesis and oak ecosystems in the Chariton River Hills of Missouri. MS Thesis, University of Missouri, Columbia, Missouri.
- Steele, Amber M.; Kabrick, John M.; Miles, Randall J. 2013. Regional and geomorphic influence on the productivity, composition, and structure of oak ecosystems in the western central hardwoods region. In: Miller, Gary W.; Schuler, Thomas M.; Gottschalk, Kurt W.; Brooks, John R.; Grushecky, Shawn T.; Spong, Ben D.; Rentch, James S., eds. Proceedings, 18th Central Hardwood Forest Conference; 2012 March 26-28; Morgantown, WV; Gen. Tech.

Rep. NRS-P-117. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 80-92.

### **Glossary**

*Backslope* – a hillslope profile position that forms the steepest and generally linear, middle portion of the slope.

*Backswamp* – marshy or swampy, depressed areas of flood plains between natural levees and valley sides or terraces

*Calcareous* – the presence of calcium carbonate in the soil parent material within the rooting zone; relatively alkaline

*Claypan* – a dense, compact, slowly permeable layer in the subsoil having much higher clay content than the overlying material

*Chert* – hard, extremely dense or compact crystalline sedimentary rock, consisting dominantly of interlocking crystals of quartz

*Cliff* – a significant vertical, or near vertical, rock exposure

*Deep Loess* – a geographic area characterized by thick, dissected loess deposits, formed immediately adjacent to the edges of the Missouri and Mississippi River floodplains

*Dolomite* – a type of sedimentary rock that is a carbonate mineral composed of calcium magnesium carbonate

*Drainageway* – the upper most reach of a stream channel system characterized by little meandering

*Dry* – a site where soil moisture is limiting during the growing season; low available water capacity

*Dune* – a low mound, ridge, bank or hill of loose, wind-blown sand

*Exposed* – steep, south and west-facing slopes, which are warmer and drier than other slope aspects

*Flatwoods* – a type of woodland that occurs on soils with a root restricting subsoil layer within 20 to 30 inches, resulting in very slow runoff and ponding that remains saturated for most of the winter and early spring months but dries out and becomes very dry in the summer months; plants that grow there must be adapted to both conditions

*Floodplain* – the nearly level plain that borders a stream and is subject to inundation under flood-stage conditions

*Footslope* – a hillslope position at the base of a slope where hillslope sediment (colluvium) accumulates

*Forest* – a vegetative community dominated by trees forming a closed canopy and interspersed with shade-tolerant understory species

*Fragipan* – a dense, brittle subsoil horizon that is extremely hard and compact when dry

*Glade* – open, rocky, barren vegetative community dominated by drought-adapted forbs and grasses, typically with scattered, stunted woody plants

*Igneous* – bedrock created by cooling and crystallization of magma forming igneous rock. Granite and rhyolite are typical igneous bedrocks in Missouri

*Knob* – precambrian ancient exposed igneous rocks of prominent rounded mountain tops

*Limestone* – a type of sedimentary rock composed largely of calcium carbonate

*Loess* – material transported and deposited by wind and consisting predominantly of silt-size particles

*Loamy* – soil material containing a relatively equal mixture of sand and silt and a somewhat smaller proportion of clay

*Marsh* – a type of wetland that is dominated by herbaceous rather than woody plant species

*Moist* – a site that is moderately well to well drained and has high available water capacity, resulting in a well-balanced supply of moisture (neither too dry nor too wet).

*Mudstone* – blocky or massive, fine-grained sedimentary rock in which the proportions of clay and silt are approximately equal

*Natric* – a soil horizon that displays a blocky, columnar, or prismatic structure and has a subhorizon with an exchangeable-sodium saturation of over 15%

*Outwash* – stratified sediments of sand and gravel removed or “washed out” from a glacier by melt-water streams

*Prairie* – a vegetative community dominated by perennial grasses and forbs with scattered shrubs and very few trees

*Protected* – steep, north- and east-facing slopes, which are generally cooler and moister than other slope aspects

*Residuum* - unconsolidated, weathered, or partly weathered mineral material that accumulates by disintegration of bedrock in place

*Riser* – a component of terraces and flood-plain steps consisting of the steep side slope; the escarpment

*Riverfront* – a vegetative community in the floodplain immediately adjacent and generally parallel to a river or stream channel

*Sandy* – a coarse-sized soil containing a large mixture of sand and gravels and a somewhat smaller proportion of silts and clays with excessive drainage

*Sandstone* – a sedimentary rock containing dominantly sand-size particles

*Savanna* – grasslands interspersed with open-grown scattered trees, groupings of trees, and shrubs

*Shale* – a sedimentary rock formed from clay, silty clay, or silty clay loam deposits and having the tendency to split into thin layers

*Shallow* – a site with bedrock within 20 inches of the surface

*Shoulder* – the slope profile position that forms the convex surface near the top of a hill slope; it comprises the transition zone from summit to backslope

*Sinkhole* – a closed, circular or elliptical depression, commonly funnel-shaped, characterized by subsurface drainage and formed either by dissolution of the surface of underlying bedrock or by collapse of underlying caves within bedrock

*Summit* – the top or highest area of a hillslope

*Swale* – shallow, closed depressions irregularly spaced across a floodplain or terrace with an irregularly undulating surface.

*Swamp* – an area of low, saturated ground, intermittently or permanently covered with water, and predominantly vegetated by shrubs and trees.

*Talus* – rock fragments of any size or shape (usually coarse and angular) derived from and lying at the base of a cliff or very steep rock slope.

*Terrace* – a step-like surface, bordering a valley floor that represents the former position of a flood plain

*Till* – dominantly unsorted and unstratified soil material deposited directly by a glacier

*Upland* – a general term for the higher ground of a region, in contrast with a low-lying, adjacent land such as a valley or floodplain

*Wet* – a somewhat poorly, poorly or very poorly drained site that has an oversupply of moisture during the growing season

*Woodland* – a highly variable vegetative community with a canopy of trees ranging from 30 to 100 percent closure with a sparse midstory and a dense ground flora of grasses, sedges and forbs

*USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex (including gender identity and expression), marital status, familial status, parental status, religion, sexual orientation, political beliefs, genetic information, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) To file a complaint of discrimination, complete, sign and mail a program discrimination complaint form, available at any USDA office location or online at [www.ascr.usda.gov](http://www.ascr.usda.gov) or call toll free at (866) 632-9992 (voice) to obtain additional information, the appropriate office or to request documents. Individuals who are deaf, hard of hearing or have speech disabilities may contact USDA through the Federal Relay service at (800) 877-8339 or (800) 845-6136 (in Spanish). USDA is an equal opportunity provider, employer and lender.*

**Helping People Help the Land**