

Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description (Brief, Generated)

Okeechobee County, Florida

Map Unit: 2—Basinger fine sand, 0 to 2 percent slopes

Component: Basinger (90%)

The Basinger component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during July, August. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

Component: EauGallie (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Margate (3%)

Generated brief soil descriptions are created for major components. The Margate soil is a minor component.

Component: Placid, depressional (3%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 3—Basinger and Placid soils, depressional**Component: Basinger, depressional (50%)**

The Basinger, depressional component makes up 50 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Placid, depressional (40%)

The Placid, depressional component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 6 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: St. Johns (5%)

Generated brief soil descriptions are created for major components. The St. Johns soil is a minor component.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Map Unit: 4—Bradenton fine sand

Component: Bradenton (85%)

The Bradenton component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY012FL Wetland Hardwood Hammock ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Riviera (4%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Component: Ft. Drum (4%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Component: Parkwood (4%)

Generated brief soil descriptions are created for major components. The Parkwood soil is a minor component.

Component: Wabasso (3%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 5—Valkaria fine sand

Component: Valkaria (90%)

The Valkaria component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ft. Drum (4%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Riviera (3%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Map Unit: 6—Manatee loamy fine sand, depressional

Component: Manatee, depressional (80%)

The Manatee, depressional component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 10 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana, depressional (7%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Parkwood (7%)

Generated brief soil descriptions are created for major components. The Parkwood soil is a minor component.

Component: Placid, depressional (6%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 7—Floridana, Riveria, and Placid soils, depressional**Component:** Floridana, depressional (40%)

The Floridana, depressional component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 11 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Riviera, depressional (30%)

The Riviera, depressional component makes up 30 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Placid, depressional (20%)

The Placid, depressional component makes up 20 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 6 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelanta, depressional (5%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Manatee, depressional (5%)

Generated brief soil descriptions are created for major components. The Manatee soil is a minor component.

Map Unit: 8—Pineda fine sand, 0 to 2 percent slopes

Component: Pineda (93%)

The Pineda component makes up 93 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 9—Riviera fine sand**Component: Riviera (90%)**

The Riviera component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bradenton (3%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Valkaria (2%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Wabasso (2%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 10—Ft. Drum fine sand**Component: Ft. Drum (80%)**

The Ft. Drum component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY005FL Cabbage Palm Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 15 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 20 within 30 inches of the soil surface.

Component: Bradenton (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Parkwood (5%)

Generated brief soil descriptions are created for major components. The Parkwood soil is a minor component.

Component: Pineda (5%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Map Unit: 11—Immokalee fine sand, 0 to 2 percent slopes

Component: Immokalee (90%)

The Immokalee component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats, coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Margate (3%)

Generated brief soil descriptions are created for major components. The Margate soil is a minor component.

Component: Placid, depressional (2%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 12—Udorthents, 2 to 35 percent slopes**Component: Udorthents (100%)**

The Udorthents component makes up 100 percent of the map unit. Slopes are 2 to 35 percent. This component is on fills on marine terraces on coastal plains. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 13—Manatee, Floridana, and Tequesta soils, frequently flooded**Component: Manatee, frequently flooded (45%)**

The Manatee, frequently flooded component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on marshes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 7 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana, frequently flooded (25%)

The Floridana, frequently flooded component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on marshes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 11 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Tequesta (15%)

The Tequesta component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on marshes on marine terraces on coastal plains. The parent material consists of organic material over sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 48 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 1 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelanta, frequently flooded (4%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Placid, frequently flooded (4%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Component: Basinger (4%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Riviera (3%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Map Unit: 14—Myakka fine sand, 0 to 2 percent slopes**Component: Myakka (90%)**

The Myakka component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods, coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: EauGallie, non-hydric (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Placid, depressional (1%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 15—Okeelanta muck**Component: Okeelanta, depressional (90%)**

The Okeelanta, depressional component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Terra Ceia (5%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Component: Placid, depressional (5%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 17—Orsino fine sand**Component:** Orsino (90%)

The Orsino component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of eolian deposits and/or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 54 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY001FL Sand Pine Scrub ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pomello (5%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

Component: Immokalee (5%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Map Unit: 18—Parkwood fine sand**Component:** Parkwood (90%)

The Parkwood component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September. Organic matter content in the surface horizon is about 4 percent. This component is in the R155XY012FL Wetland Hardwood Hammock ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 15 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ft. Drum (5%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Component: Bradenton (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Map Unit: 19—Floridana, Placid, and Okeelanta soils, frequently flooded

Component: Floridana (40%)

The Floridana component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 11 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Placid, frequently flooded (25%)

The Placid, frequently flooded component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 6 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelanta, frequently flooded (20%)

The Okeelanta, frequently flooded component makes up 20 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is frequently flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: St. Johns (3%)

Generated brief soil descriptions are created for major components. The St. Johns soil is a minor component.

Component: Basinger (3%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (3%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Riviera (3%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Component: Myakka (3%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Map Unit: 20—Pomello fine sand, 0 to 5 percent slopes

Component: Pomello (90%)

The Pomello component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY001FL Sand Pine Scrub ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Immokalee (3%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Myakka (3%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: St. Johns (2%)

Generated brief soil descriptions are created for major components. The St. Johns soil is a minor component.

Component: Orsino (2%)

Generated brief soil descriptions are created for major components. The Orsino soil is a minor component.

Map Unit: 21—Adamsville fine sand, organic substratum

Component: Adamsville (90%)

The Adamsville component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY005FL Cabbage Palm Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Map Unit: 23—St. Johns fine sand

Component: St. Johns (85%)

The St. Johns component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Immokalee (5%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Map Unit: 24—Terra Ceia muck**Component: Terra Ceia (80%)**

The Terra Ceia component makes up 80 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Placid, depressional (10%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Component: Okeelanta, depressional (10%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Map Unit: 25—Wabasso fine sand**Component: Wabasso (90%)**

The Wabasso component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Valkaria (3%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Riviera (3%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Map Unit: 99—Water

Component: Water (100%)

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

Data Source Information

Soil Survey Area: Okeechobee County, Florida

Survey Area Data: Version 9, Dec 18, 2013