

## Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

## Report—Map Unit Description (Brief, Generated)

### Seminole County, Florida

**Map Unit:** 2—Adamsville-Sparr fine sands

**Component:** Adamsville (54%)

The Adamsville component makes up 54 percent of the map unit. Slopes are 0 to 2 percent. This component is on rises on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Sparr (36%)**

The Sparr component makes up 36 percent of the map unit. Slopes are 0 to 2 percent. This component is on rises on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during July, August, September. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Immokalee (10%)**

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

**Map Unit: 3—Arents, 0 to 5 percent slopes****Component: Arents (100%)**

The Arents component makes up 100 percent of the map unit. Slopes are 0 to 5 percent. This component is on fills, rises on marine terraces on coastal plains. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during June, July, August, September. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Map Unit: 4—Astatula fine sand, 0 to 5 percent slopes****Component: Astatula (90%)**

The Astatula component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on hillslopes on marine terraces on coastal plains, ridges on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Apopka (5%)**

Generated brief soil descriptions are created for major components. The Apopka soil is a minor component.

**Component: Pomello (5%)**

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

**Map Unit: 5—Astatula fine sand, 5 to 8 percent slopes**

**Component: Astatula (97%)**

The Astatula component makes up 97 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Paola (3%)**

Generated brief soil descriptions are created for major components. The Paola soil is a minor component.

**Map Unit: 6—Astatula-Apopka fine sands, 0 to 5 percent slopes**

**Component: Astatula (65%)**

The Astatula component makes up 65 percent of the map unit. Slopes are 0 to 5 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Apopka (22%)**

The Apopka component makes up 22 percent of the map unit. Slopes are 0 to 5 percent. This component is on hillslopes on marine terraces on coastal plains, ridges on marine terraces on coastal plains. The parent material consists of eolian deposits and/or sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Tavares (7%)**

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

**Component: Pomello (6%)**

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

**Map Unit: 7—Astatula-Apopka fine sands, 5 to 8 percent**

**Component: Astatula (63%)**

The Astatula component makes up 63 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Apopka (27%)**

The Apopka component makes up 27 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian deposits and/or sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Millhopper (5%)**

Generated brief soil descriptions are created for major components. The Millhopper soil is a minor component.

**Component: Tavares (5%)**

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

**Map Unit: 8—Astatula-Apopka fine sands, 8 to 12 percent slopes**

**Component: Astatula (72%)**

The Astatula component makes up 72 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Apopka (25%)**

The Apopka component makes up 25 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian deposits and/or sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Millhopper (3%)**

Generated brief soil descriptions are created for major components. The Millhopper soil is a minor component.

**Map Unit: 9—Basinger and Delray fine sands**

**Component: Basinger (60%)**

The Basinger component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Delray (32%)**

The Delray component makes up 32 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 4 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Malabar (4%)**

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

**Component: Wabasso (4%)**

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

**Map Unit: 10—Basinger, Samsula, and Hontoon soils, depressional**

**Component: Basinger (58%)**

The Basinger component makes up 58 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 14 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Samsula (15%)**

The Samsula component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 60 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Hontoon (15%)**

The Hontoon component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 80 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Felda (3%)**

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

**Component: EauGallie (3%)**

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component: Smyrna (2%)**

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

**Component: Myakka (2%)**

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

**Component: St. Johns (2%)**

Generated brief soil descriptions are created for major components. The St. Johns soil is a minor component.

**Map Unit: 11**—Basinger and Smyrna fine sands, depressional

**Component: Basinger (63%)**

The Basinger component makes up 63 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 14 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Smyrna (28%)**

The Smyrna component makes up 28 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: EauGallie (5%)**

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component: Malabar (4%)**

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

**Map Unit: 12**—Canova and Terra Ceia mucks

**Component: Canova, drained (75%)**

The Canova, drained component makes up 75 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September. Organic matter content in the surface horizon is about 55 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Terra Ceia, drained (25%)

The Terra Ceia, drained component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September. Organic matter content in the surface horizon is about 75 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Map Unit:** 13—EauGallie and Immokalee fine sands

**Component:** EauGallie (56%)

The EauGallie component makes up 56 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Immokalee (35%)

The Immokalee component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Malabar (9%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

**Map Unit:** 14—Felda fine sand, saline, frequently flooded

**Component:** Felda, flooded (90%)

The Felda, flooded component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, July, August, September, October, November, December. Organic matter content in the surface horizon is about 15 percent. This component is in the R155XY009FL Salt Marsh ecological site. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Pineda (10%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

**Map Unit:** 15—Felda and Manatee mucky fine sands, depressional

**Component:** Felda (56%)

The Felda component makes up 56 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 15 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Manatee (38%)**

The Manatee component makes up 38 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 23 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Delray (3%)**

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

**Component: Wabasso (3%)**

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

**Map Unit: 16—Immokalee sand**

**Component: Immokalee (90%)**

The Immokalee component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** EauGallie (10%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Map Unit:** 17—Brighton, Samsula, and Sanibel mucks

**Component:** Brighton, drained (47%)

The Brighton, drained component makes up 47 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of woody organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September. Organic matter content in the surface horizon is about 75 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Samsula, drained (35%)

The Samsula, drained component makes up 35 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September. Organic matter content in the surface horizon is about 60 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Sanibel, drained (15%)**

The Sanibel, drained component makes up 15 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of thin organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September. Organic matter content in the surface horizon is about 45 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Delray (2%)**

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

**Component: Basinger (1%)**

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Map Unit: 18—Malabar fine sand****Component: Malabar (86%)**

The Malabar component makes up 86 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: EauGallie (5%)**

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component: Basinger (5%)**

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Component:** Felda (4%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

**Map Unit:** 19—Manatee, Floridana, and Holopaw soils, frequently flooded

**Component:** Manatee, flooded (61%)

The Manatee, flooded component makes up 61 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 7 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Floridana, flooded (21%)

The Floridana, flooded component makes up 21 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 11 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Holopaw, flooded (15%)

The Holopaw, flooded component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Basinger, flooded (3%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Map Unit:** 20—Myakka and EauGallie fine sands

**Component:** Myakka (58%)

The Myakka component makes up 58 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 4 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** EauGallie (32%)

The EauGallie component makes up 32 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Basinger (5%)**

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Component: Pompano, flooded (5%)**

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

**Map Unit: 21—Nittaw mucky fine sand, depressional****Component: Nittaw (91%)**

The Nittaw component makes up 91 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is high. Shrink-swell potential is high. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 15 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Basinger (9%)**

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Map Unit: 22—Nittaw muck, occasionally flooded****Component: Nittaw, flooded (100%)**

The Nittaw, flooded component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is high. Shrink-swell potential is high. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 55 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Map Unit: 23—Nittaw, Okeelanta, and Basinger soils, frequently flooded****Component: Nittaw, flooded (45%)**

The Nittaw, flooded component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is high. Shrink-swell potential is high. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 55 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Okeelanta, flooded (34%)**

The Okeelanta, flooded component makes up 34 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 73 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Basinger, flooded (19%)**

The Basinger, flooded component makes up 19 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Pompano, flooded (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

**Map Unit:** 24—Paola-St. Lucie sands, 0 to 5 percent slopes**Component:** Paola (52%)

The Paola component makes up 52 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 0 percent. This component is in the R155XY001FL Sand Pine Scrub ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** St. Lucie (43%)

The St. Lucie component makes up 43 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Tavares (5%)

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

**Map Unit:** 25—Pineda fine sand**Component:** Pineda (89%)

The Pineda component makes up 89 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** EauGallie (6%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component:** Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

**Map Unit:** 26—Udorthents, excavated

**Component:** Udorthents (90%)

The Udorthents component makes up 90 percent of the map unit. Slopes are 0 to 8 percent. This component is on — Error in Exists On —. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Aquents (10%)

Generated brief soil descriptions are created for major components. The Aquents soil is a minor component.

**Map Unit:** 27—Pomello fine sand, 0 to 5 percent slopes

**Component:** Pomello (88%)

The Pomello component makes up 88 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** EauGallie (6%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component:** Millhopper (6%)

Generated brief soil descriptions are created for major components. The Millhopper soil is a minor component.

**Map Unit:** 28—Pompano fine sand, occasionally flooded

**Component:** Pompano, flooded (90%)

The Pompano, flooded component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Nittaw, flooded (10%)

Generated brief soil descriptions are created for major components. The Nittaw soil is a minor component.

**Map Unit:** 29—St. Johns and EauGallie fine sands

**Component:** St. Johns (57%)

The St. Johns component makes up 57 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: EauGallie (36%)**

The EauGallie component makes up 36 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Felda (7%)**

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

**Map Unit: 30—Seffner fine sand**

**Component: Seffner (91%)**

The Seffner component makes up 91 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Immokalee (9%)**

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

**Map Unit:** 31—Tavares-Millhopper fine sands, 0 to 5 percent slopes

**Component:** Tavares (45%)

The Tavares component makes up 45 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Millhopper (41%)

The Millhopper component makes up 41 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component:** Astatula (5%)

Generated brief soil descriptions are created for major components. The Astatula soil is a minor component.

**Component:** Felda (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

**Component:** Pomello (4%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

**Map Unit: 32**—Tavares-Millhopper fine sands, 5 to 8 percent slopes

**Component: Tavares (58%)**

The Tavares component makes up 58 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Millhopper (40%)**

The Millhopper component makes up 40 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Astatula (2%)**

Generated brief soil descriptions are created for major components. The Astatula soil is a minor component.

**Map Unit: 33**—Terra Ceia muck, frequently flooded

**Component: Terra Ceia, flooded (100%)**

The Terra Ceia, flooded component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of herbaceous organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is very high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 80 percent. This component is in the R155XY010FL Freshwater Marshes And Ponds ecological site. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Map Unit: 34—Urban land**

**Component: Urban land (85%)**

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

**Component: Astatula (2%)**

Generated brief soil descriptions are created for major components. The Astatula soil is a minor component.

**Component: Paola (2%)**

Generated brief soil descriptions are created for major components. The Paola soil is a minor component.

**Component: Millhopper (2%)**

Generated brief soil descriptions are created for major components. The Millhopper soil is a minor component.

**Component: Apopka (2%)**

Generated brief soil descriptions are created for major components. The Apopka soil is a minor component.

**Component: EauGallie (2%)**

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

**Component: Myakka (2%)**

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

**Component: Pomello (1%)**

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

**Component: Tavares (1%)**

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

**Component: St. Lucie (1%)**

Generated brief soil descriptions are created for major components. The St. Lucie soil is a minor component.

**Map Unit: 35—Wabasso fine sand****Component: Wabasso (90%)**

The Wabasso component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a slightly sodic horizon within 30 inches of the soil surface.

**Component: Pineda (10%)**

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

**Map Unit: 99—Water****Component: Water (100%)**

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

**Data Source Information**

Soil Survey Area: Seminole County, Florida

Survey Area Data: Version 11, Dec 7, 2013