

Map Unit Description

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. All the soils of a series have major horizons that are similar in composition, thickness, and arrangement. Soils of a given series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Additional information about the map units described in this report is available in other soil reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the soil reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description

Walton County, Florida

1—Albany-Pactolus loamy sands, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Albany and similar soils: 55 percent

Pactolus and similar soils: 30 percent
Minor components: 15 percent

Description of Albany

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)
Depth to water table: About 12 to 30 inches
Frequency of flooding: Rare
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.3 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3e
Hydrologic Soil Group: A/D
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Typical profile

0 to 11 inches: Loamy sand
11 to 45 inches: Loamy sand
45 to 80 inches: Sandy loam

Description of Pactolus

Setting

Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy marine and fluvial deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 18 to 36 inches
Frequency of flooding: Rare

Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A/D
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP129FL)

Typical profile

0 to 12 inches: Loamy sand
12 to 28 inches: Loamy sand
28 to 80 inches: Sand

Minor Components

Blanton

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvial
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Escambia

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluvial, talus
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Leefield

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluvial
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Bonifay

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluvial
Down-slope shape: Convex

Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Chipley

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Stilson

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

2—Bonifay loamy sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 350 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Bonifay and similar soils: 80 percent
Minor components: 20 percent

Description of Bonifay

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 48 to 60 inches

Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.6 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Typical profile

0 to 7 inches: Loamy sand
7 to 44 inches: Loamy sand
44 to 80 inches: Sandy loam

Minor Components

Fuquay

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Blanton

Percent of map unit: 4 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Albany

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Foxworth

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

3—Bonifay loamy sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 20 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Bonifay and similar soils: 75 percent
Minor components: 25 percent

Description of Bonifay

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 5 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 48 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.4 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 4s
Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Typical profile

0 to 5 inches: Loamy sand
5 to 48 inches: Loamy sand
48 to 67 inches: Sandy clay loam
67 to 80 inches: Sandy loam

Minor Components

Fuquay

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Lakeland

Percent of map unit: 3 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP195FL)

Albany

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluvium, talus
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Blanton

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Troup

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Lucy

Percent of map unit: 3 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Leefield

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

4—Chipley sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 10 to 350 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Chipley and similar soils: 80 percent
Minor components: 20 percent

Description of Chipley

Setting

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 24 to 36 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.7 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Typical profile

0 to 6 inches: Sand
6 to 80 inches: Sand

Minor Components

Albany

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Foxworth

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Mandarin

Percent of map unit: 3 percent
Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Hurricane

Percent of map unit: 3 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Blanton

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Leon

Percent of map unit: 2 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Eglin

Percent of map unit: 2 percent

Landform: Marine terraces, rises

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

Lakeland

Percent of map unit: 2 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

5—Chipley sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Chipley and similar soils: 80 percent

Minor components: 20 percent

Description of Chipley

Setting

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 5 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: About 24 to 36 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 3.7 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 4s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP080FL)

Typical profile

0 to 5 inches: Sand

5 to 80 inches: Sand

Minor Components

Albany

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Blanton

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Foxworth

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Floralia

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Lakeland

Percent of map unit: 2 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP195FL)

Troup

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvium

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Leefield

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluvium

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

6—Escambia sandy loam, 2 to 5 percent slopes**Map Unit Setting**

Elevation: 50 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Escambia and similar soils: 75 percent

Minor components: 25 percent

Description of Escambia**Setting**

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluvium, talus

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 2 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately low to moderately high (0.06 to 0.57 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: High (about 9.9 inches)

Interpretive groups

Farmland classification: Prime farmland if drained

Land capability (nonirrigated): 2e

Hydrologic Soil Group: C

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Typical profile

0 to 9 inches: Sandy loam

9 to 17 inches: Sandy loam

17 to 67 inches: Sandy loam

67 to 80 inches: Loam

Minor Components**Floralia**

Percent of map unit: 5 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Leefield

Percent of map unit: 4 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Fuquay

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Stilson

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Malbis

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Kinston

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Bibb

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

8—Dorovan-Pamlico association, frequently flooded

Map Unit Setting

Elevation: 0 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Dorovan and similar soils: 60 percent

Pamlico and similar soils: 20 percent

Minor components: 20 percent

Description of Dorovan**Setting**

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Organic material

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: Frequent

Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Very high (about 22.7 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 7w

Hydrologic Soil Group: B/D

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Typical profile

0 to 60 inches: Muck

Description of Pamlico**Setting**

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Concave

Parent material: Herbaceous organic material over sandy marine deposits

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 5.95 in/hr)

Depth to water table: About 0 inches
Frequency of flooding: Frequent
Frequency of ponding: Frequent
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very high (about 14.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7w
Hydrologic Soil Group: A/D
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Typical profile

0 to 30 inches: Muck
30 to 80 inches: Sand

Minor Components

Bibb

Percent of map unit: 6 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Kinston

Percent of map unit: 6 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Leon

Percent of map unit: 5 percent
Landform: Flats on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

9—Dothan loamy sand, 0 to 2 percent slopes

Map Unit Setting

Elevation: 100 to 500 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Dothan and similar soils: 75 percent
Minor components: 25 percent

Description of Dothan

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.7 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 1
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Typical profile

0 to 9 inches: Loamy sand
9 to 57 inches: Sandy clay loam
57 to 80 inches: Sandy clay loam

Minor Components

Floralia

Percent of map unit: 5 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Fuquay

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Orangeburg

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Tifton

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Malbis

Percent of map unit: 5 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

10—Dothan loamy sand, 2 to 5 percent slopes

Map Unit Setting

Elevation: 100 to 400 feet

Mean annual precipitation: 40 to 69 inches

Mean annual air temperature: 55 to 70 degrees F

Frost-free period: 190 to 310 days

Map Unit Composition

Dothan and similar soils: 80 percent

Minor components: 20 percent

Description of Dothan

Setting

Landform: Interfluves

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Marine deposits

Properties and qualities

Slope: 2 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 32 to 55 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Farmland classification: All areas are prime farmland

Land capability (nonirrigated): 2e

Hydrologic Soil Group: C

Typical profile

0 to 12 inches: Loamy sand

12 to 24 inches: Sandy clay loam

24 to 34 inches: Sandy clay loam

34 to 48 inches: Sandy clay loam

48 to 65 inches: Sandy clay loam

Minor Components

Fuquay

Percent of map unit: 5 percent

Landform: Interfluves

Down-slope shape: Convex

Across-slope shape: Linear

Nankin

Percent of map unit: 5 percent

Landform: Broad interstream divides

Landform position (two-dimensional): Backslope, shoulder, summit

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Cowarts

Percent of map unit: 5 percent

Landform: Broad interstream divides

Down-slope shape: Convex

Across-slope shape: Linear

Clarendon

Percent of map unit: 5 percent

Landform: Flats on broad interstream divides

Down-slope shape: Linear

Across-slope shape: Linear

11—Dothan loamy sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 30 to 700 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Dothan and similar soils: 75 percent

Minor components: 25 percent

Description of Dothan

Setting

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 5 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 36 to 60 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 3e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Typical profile

0 to 7 inches: Loamy sand
7 to 40 inches: Sandy clay loam
40 to 80 inches: Sandy clay loam

Minor Components

Norfolk

Percent of map unit: 5 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Orangeburg

Percent of map unit: 5 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on ridges and side slopes of mesic uplands (G133AA312FL), Unnamed (G133AP140FL)

Malbis

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP328FL)

Tifton

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Cowarts

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Angie

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluvial, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Floral

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

12—Foxworth sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 350 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Foxworth and similar soils: 85 percent
Minor components: 15 percent

Description of Foxworth**Setting**

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Eolian or sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Very high
(19.84 to 50.02 in/hr)
Depth to water table: About 42 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.5 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Typical profile

0 to 7 inches: Sand
7 to 54 inches: Sand
54 to 80 inches: Sand

Minor Components**Blanton**

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Lakeland

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

Troup

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Chipley

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Albany

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

13—Fuquay loamy sand, 0 to 5 percent slopes**Map Unit Setting**

Elevation: 20 to 400 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Fuquay and similar soils: 80 percent
Minor components: 20 percent

Description of Fuquay**Setting**

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex

Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 48 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 6.6 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 2s
Hydrologic Soil Group: B
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Typical profile

0 to 5 inches: Loamy sand
5 to 26 inches: Loamy sand
26 to 74 inches: Sandy loam
74 to 80 inches: Coarse sandy loam

Minor Components

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Bonifay

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Bonneau

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Albany

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Escambia

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Floral

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Lucy

Percent of map unit: 2 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

14—Fuquay loamy sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 40 to 400 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Fuquay and similar soils: 75 percent

Minor components: 25 percent

Description of Fuquay

Setting

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvium

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy marine deposits

Properties and qualities

Slope: 5 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 48 to 72 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 6.0 inches)

Interpretive groups

Farmland classification: Farmland of local importance

Land capability (nonirrigated): 3s

Hydrologic Soil Group: B

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Typical profile

0 to 5 inches: Loamy sand

5 to 34 inches: Loamy sand

34 to 43 inches: Sandy loam

43 to 80 inches: Sandy clay loam

Minor Components

Bonifay

Percent of map unit: 4 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Lucy

Percent of map unit: 4 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Convex

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP144FL)

Dothan

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Lakeland

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP195FL)

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Floralia

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

15—Kinston-Johnston-Bibb complex, frequently flooded

Map Unit Setting

Elevation: 0 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Kinston and similar soils: 30 percent
Johnston and similar soils: 20 percent
Bibb and similar soils: 15 percent
Minor components: 35 percent

Description of Kinston

Setting

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Loamy alluvium

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 9.0 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 6w
Hydrologic Soil Group: B/D
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Typical profile

0 to 6 inches: Loam
6 to 42 inches: Sandy clay loam
42 to 48 inches: Sand
48 to 80 inches: Clay loam

Description of Johnston

Setting

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Loamy and sandy alluvium

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)
Depth to water table: About 0 inches
Frequency of flooding: Frequent
Frequency of ponding: Frequent
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: High (about 9.7 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7w
Hydrologic Soil Group: A/D
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Typical profile

0 to 37 inches: Mucky loam

37 to 65 inches: Sand

Description of Bibb**Setting**

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Loamy and sandy alluvium

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 6 to 18 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 8.0 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 5w

Hydrologic Soil Group: B/D

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Typical profile

0 to 12 inches: Loam

12 to 37 inches: Sandy loam

37 to 65 inches: Sand

Minor Components**Dorovan**

Percent of map unit: 5 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Pamlico

Percent of map unit: 5 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Rutlege

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Albany

Percent of map unit: 4 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Chiple

Percent of map unit: 4 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Escambia

Percent of map unit: 4 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Floral

Percent of map unit: 4 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Leefield

Percent of map unit: 4 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

16—Kureb sand, 0 to 8 percent slopes**Map Unit Setting**

Elevation: 0 to 40 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Kureb and similar soils: 80 percent

Minor components: 20 percent

Description of Kureb**Setting**

Landform: Dunes on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Eolian deposits or sandy fluvial or marine deposits

Properties and qualities

Slope: 0 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Very low (about 1.8 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 7s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G152AP192FL)

Typical profile

0 to 4 inches: Sand
 4 to 17 inches: Sand
 17 to 68 inches: Sand
 68 to 80 inches: Sand

Minor Components**Resota**

Percent of map unit: 10 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G152AP142FL)

Mandarin

Percent of map unit: 5 percent
Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Corolla

Percent of map unit: 3 percent
Landform: Rises on dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP180FL)

Newhan

Percent of map unit: 2 percent
Landform: Dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G152AP198FL)

17—Lakeland sand, 0 to 5 percent slopes**Map Unit Setting**

Elevation: 30 to 300 feet
Mean annual precipitation: 59 to 69 inches
Mean annual air temperature: 63 to 72 degrees F
Frost-free period: 252 to 295 days

Map Unit Composition

Lakeland and similar soils: 77 percent
Minor components: 23 percent

Description of Lakeland**Setting**

Landform: Hills on marine terraces
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy eolian deposits and/or marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability classification (irrigated): 3s
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Ecological site: Longleaf Pine-Turkey Oak Hills (R133AY002FL)
Other vegetative classification: Unnamed (G152AT141FL)

Typical profile

0 to 7 inches: Sand
7 to 80 inches: Sand

Minor Components**Troup**

Percent of map unit: 14 percent
Landform: — error in exists on —
Landform position (two-dimensional): Summit
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Ecological site: Longleaf Pine-Turkey Oak Hills (R133AY002FL)
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G152AT141FL)

Bonifay

Percent of map unit: 9 percent

Landform: Hills on marine terraces

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex, linear

Across-slope shape: Linear, convex

Ecological site: Longleaf Pine-Turkey Oak Hills (R133AY002FL)

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AT141FL)

18—Lakeland sand, 5 to 12 percent slopes**Map Unit Setting**

Elevation: 20 to 300 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Lakeland and similar soils: 80 percent

Minor components: 20 percent

Description of Lakeland**Setting**

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Eolian or sandy marine deposits

Properties and qualities

Slope: 5 to 12 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 3.1 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 6s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP195FL)

Typical profile

0 to 3 inches: Sand

3 to 80 inches: Sand

Minor Components**Troup**

Percent of map unit: 10 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interflue

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP144FL)

Bonifay

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interflue, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP141FL)

Foxworth

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interflue

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Chipley

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interflue

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP080FL)

19—Lakeland sand, 12 to 30 percent slopes

Map Unit Setting

Elevation: 20 to 300 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Lakeland and similar soils: 75 percent

Minor components: 25 percent

Description of Lakeland

Setting

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Eolian or sandy marine deposits

Properties and qualities

Slope: 12 to 30 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 3.1 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 7s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP195FL)

Typical profile

0 to 5 inches: Sand

5 to 80 inches: Sand

Minor Components

Troup

Percent of map unit: 10 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Side slope, interfluvium

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP147FL)

Foxworth

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP142FL)

Bonifay

Percent of map unit: 4 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP141FL)

Chipleay

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP080FL)

Dorovan

Percent of map unit: 2 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Pamlico

Percent of map unit: 2 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

20—Leefield-Stilson loamy sands, 0 to 5 percent slopes

Map Unit Setting

Elevation: 30 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Leefield and similar soils: 70 percent

Stilson and similar soils: 15 percent

Minor components: 15 percent

Description of Leefield

Setting

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 5.1 inches)

Interpretive groups

Farmland classification: Farmland of local importance

Land capability (nonirrigated): 2w

Hydrologic Soil Group: C/D

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Typical profile

0 to 7 inches: Loamy sand

7 to 26 inches: Loamy sand

26 to 34 inches: Fine sandy loam

34 to 80 inches: Sandy clay loam

Description of Stilson

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 30 to 36 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.3 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 2s
Hydrologic Soil Group: A
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Typical profile

0 to 7 inches: Loamy sand
7 to 25 inches: Loamy sand
25 to 32 inches: Fine sandy loam
32 to 80 inches: Sandy clay loam

Minor Components

Escambia

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Fuquay

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Blanton

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Dothan

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Floralia

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Pantego

Percent of map unit: 2 percent

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

21—Leon sand, 0 to 2 percent slopes

Map Unit Setting

Elevation: 0 to 300 feet

Mean annual precipitation: 60 to 69 inches

Mean annual air temperature: 63 to 72 degrees F

Frost-free period: 252 to 306 days

Map Unit Composition

Leon and similar soils: 80 percent

Minor components: 20 percent

Description of Leon**Setting**

Landform: Marine terraces, flatwoods

Landform position (three-dimensional): Tread, talf

Down-slope shape: Linear, convex

Across-slope shape: Linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.20 to 1.98 in/hr)

Depth to water table: About 2 to 18 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 5.6 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability classification (irrigated): 4w

Land capability (nonirrigated): 4w

Hydrologic Soil Group: A/D

Ecological site: North Florida Flatwoods (R152AY004FL)

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AT013FL)

Typical profile

0 to 5 inches: Sand

5 to 18 inches: Sand

18 to 26 inches: Sand

26 to 65 inches: Sand

65 to 80 inches: Sand

Minor Components**Leon, hydric**

Percent of map unit: 5 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Tread, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AT013FL)

Pottsburg

Percent of map unit: 4 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AT003FL)

Hurricane

Percent of map unit: 4 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AT080FL)

Mandarin

Percent of map unit: 3 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G152AA131FL), Unnamed (G152AT077FL)

Pickney

Percent of map unit: 2 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Concave, linear

Across-slope shape: Concave, linear

Ecological site: North Florida Flatwoods (R153AY004FL)

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AT800FL)

Rutlege

Percent of map unit: 2 percent

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Ecological site: North Florida Flatwoods (R155XY004FL)

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G152AA145FL), Unnamed (G152AT800FL)

22—Lucy loamy sand, 0 to 5 percent slopes**Map Unit Setting**

Elevation: 50 to 500 feet

Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Lucy and similar soils: 80 percent
Minor components: 20 percent

Description of Lucy

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine and fluvial deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 6.0 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 2s
Hydrologic Soil Group: B
Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Typical profile

0 to 8 inches: Loamy sand
8 to 33 inches: Loamy sand
33 to 39 inches: Sandy loam
39 to 80 inches: Sandy clay loam

Minor Components

Orangeburg

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Fuquay

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Bonifay

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Dothan

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Stilson

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

23—Lucy loamy sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 30 to 500 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Lucy and similar soils: 75 percent

Minor components: 25 percent

Description of Lucy

Setting

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluvial, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine and fluvial deposits

Properties and qualities

Slope: 5 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 6.3 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 3s

Hydrologic Soil Group: B

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Typical profile

0 to 9 inches: Loamy sand

9 to 28 inches: Loamy sand

28 to 33 inches: Sandy loam

33 to 80 inches: Sandy clay loam

Minor Components

Orangeburg

Percent of map unit: 10 percent

Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on ridges and side slopes of mesic uplands (G133AA312FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 2 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

25—Orangeburg sandy loam, 1 to 5 percent slopes

Map Unit Setting

Elevation: 30 to 500 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Orangeburg and similar soils: 80 percent

Minor components: 20 percent

Description of Orangeburg

Setting

Landform: Ridges on marine terraces

Landform position (two-dimensional): Summit

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 1 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Farmland classification: All areas are prime farmland

Land capability classification (irrigated): 2e

Land capability (nonirrigated): 2e

Hydrologic Soil Group: B

Ecological site: Upland Hardwood Hammock (R133AY008FL)

Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AA311FL)

Typical profile

0 to 10 inches: Sandy loam

10 to 17 inches: Sandy loam

17 to 57 inches: Sandy clay loam

57 to 80 inches: Sandy clay loam

Minor Components

Dothan

Percent of map unit: 8 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 5 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Convex, linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Lucy

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

26—Orangeburg sandy loam, 5 to 8 percent slopes

Map Unit Setting

Elevation: 30 to 500 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Orangeburg and similar soils: 75 percent

Minor components: 25 percent

Description of Orangeburg**Setting**

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 5 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 7.5 inches)

Interpretive groups

Farmland classification: All areas are prime farmland

Land capability (nonirrigated): 3e

Hydrologic Soil Group: B

Other vegetative classification: Loamy and clayey soils on ridges and side slopes of mesic uplands (G133AA312FL), Unnamed (G133AP140FL)

Typical profile

0 to 6 inches: Sandy loam

6 to 20 inches: Sandy clay loam

20 to 80 inches: Sandy clay loam

Minor Components**Lucy**

Percent of map unit: 5 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 5 percent

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 5 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 4 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP144FL)

Tifton

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

27—Rutlege fine sand

Map Unit Setting

Elevation: 0 to 300 feet
Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Rutlege and similar soils: 70 percent
Minor components: 30 percent

Description of Rutlege

Setting

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Parent material: Sandy marine deposits and/or fluvio-marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 0 to 6 inches
Frequency of flooding: Frequent
Frequency of ponding: Frequent
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7w
Hydrologic Soil Group: A/D
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Typical profile

0 to 17 inches: Fine sand
17 to 80 inches: Fine sand

Minor Components

Pickney

Percent of map unit: 10 percent
Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G133AP800FL)

Hurricane

Percent of map unit: 5 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Chipley

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Leon

Percent of map unit: 5 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Pamlico

Percent of map unit: 5 percent

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

28—Tifton fine sandy loam, 0 to 2 percent slopes**Map Unit Setting**

Elevation: 30 to 500 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Tifton and similar soils: 80 percent

Minor components: 20 percent

Description of Tifton

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 42 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 1
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Typical profile

0 to 11 inches: Fine sandy loam
11 to 18 inches: Gravelly sandy loam
18 to 29 inches: Gravelly sandy clay loam
29 to 55 inches: Sandy clay loam
55 to 80 inches: Sandy loam

Minor Components

Dothan

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Floral

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Fuquay

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Malbis

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Orangeburg

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

29—Tifton fine sandy loam, 2 to 5 percent slopes**Map Unit Setting**

Elevation: 100 to 600 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Tifton and similar soils: 80 percent
Minor components: 20 percent

Description of Tifton**Setting**

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 2 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 42 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.7 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 2e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Typical profile

0 to 9 inches: Fine sandy loam
9 to 13 inches: Gravelly fine sandy loam
13 to 57 inches: Gravelly sandy clay loam
57 to 80 inches: Gravelly sandy clay loam

Minor Components**Dothan**

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Orangeburg

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Malbis

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Angie

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Floral

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Escambia

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

30—Tifton fine sandy loam, 5 to 8 percent slopes

Map Unit Setting

Elevation: 100 to 600 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Tifton and similar soils: 75 percent
Minor components: 25 percent

Description of Tifton

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 5 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 42 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.8 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 3e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Typical profile

0 to 9 inches: Fine sandy loam
9 to 56 inches: Sandy clay loam
56 to 80 inches: Sandy clay loam

Minor Components

Dothan

Percent of map unit: 5 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Orangeburg

Percent of map unit: 5 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on ridges and side slopes of mesic uplands (G133AA312FL), Unnamed (G133AP140FL)

Angie

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 5 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Escambia

Percent of map unit: 2 percent

Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP328FL)

31—Troup sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 350 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Troup and similar soils: 85 percent
Minor components: 15 percent

Description of Troup

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.1 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Typical profile

0 to 7 inches: Sand
7 to 51 inches: Loamy sand
51 to 80 inches: Sandy clay loam

Minor Components

Bonifay

Percent of map unit: 4 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Lucy

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Blanton

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Albany

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Lakeland

Percent of map unit: 2 percent

Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

32—Troup sand, 5 to 8 percent slopes

Map Unit Setting

Elevation: 20 to 350 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Troup and similar soils: 80 percent
Minor components: 20 percent

Description of Troup

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 5 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.1 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 4s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Typical profile

0 to 4 inches: Sand
4 to 53 inches: Loamy sand
53 to 65 inches: Sandy loam
65 to 80 inches: Sandy clay loam

Minor Components

Lucy

Percent of map unit: 5 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Bonifay

Percent of map unit: 4 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Lakeland

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP195FL)

Fuquay

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Albany

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Blanton

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Chipley

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP080FL)

33—Troup sand, 8 to 12 percent slopes

Map Unit Setting

Elevation: 20 to 700 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Troup and similar soils: 75 percent
Minor components: 25 percent

Description of Troup

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 8 to 12 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 4.8 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 6s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP144FL)

Typical profile

0 to 2 inches: Sand

2 to 62 inches: Loamy sand

62 to 80 inches: Sandy loam

Minor Components

Lucy

Percent of map unit: 5 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Bonifay

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP141FL)

Cowarts

Percent of map unit: 4 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Lakeland

Percent of map unit: 4 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP195FL)

Orangeburg

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Albany

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Blanton

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

34—Troup sand, 12 to 25 percent slopes**Map Unit Setting**

Elevation: 40 to 700 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Troup and similar soils: 75 percent

Minor components: 25 percent

Description of Troup**Setting**

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 12 to 25 percent

Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 4.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP147FL)

Typical profile

0 to 2 inches: Sand
2 to 64 inches: Loamy sand
64 to 80 inches: Sandy loam

Minor Components

Lakeland

Percent of map unit: 13 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP195FL)

Cowarts

Percent of map unit: 12 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

35—Troup-Orangeburg-Cowarts loamy sands, 5 to 12 percent slopes

Map Unit Setting

Elevation: 30 to 700 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Troup and similar soils: 40 percent
Orangeburg and similar soils: 20 percent
Cowarts and similar soils: 15 percent
Minor components: 25 percent

Description of Troup

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 5 to 12 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.5 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 6s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP144FL)

Typical profile

0 to 2 inches: Sand
2 to 42 inches: Loamy sand
42 to 80 inches: Sandy loam

Description of Orangeburg

Setting

Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 5 to 12 percent

Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.3 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 4e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Typical profile

0 to 6 inches: Sandy loam
6 to 17 inches: Sandy loam
17 to 33 inches: Sandy loam
33 to 80 inches: Sandy clay loam

Description of Cowarts

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 5 to 12 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Farmland classification: Farmland of local importance
Land capability (nonirrigated): 6e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Typical profile

0 to 4 inches: Loamy sand
4 to 40 inches: Sandy clay loam
40 to 80 inches: Loamy sand

Minor Components**Bonifay**

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP141FL)

Lucy

Percent of map unit: 5 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 3 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Angie

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Convex

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP144FL)

Fuquay

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Tifton

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

36—Pits

Map Unit Composition

Pits: 100 percent

Description of Pits

Setting

Landform: Marine terraces

Landform position (three-dimensional): Interfluve, dip

Down-slope shape: Linear

Across-slope shape: Linear

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 8

Other vegetative classification: Forage suitability group not assigned (G133AA999FL), Unnamed (G133AP900FL)

37—Angie sandy loam, 2 to 5 percent slopes

Map Unit Setting

Elevation: 30 to 600 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Angie and similar soils: 80 percent

Minor components: 20 percent

Description of Angie

Setting

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 2 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 36 to 60 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 8.9 inches)

Interpretive groups

Farmland classification: All areas are prime farmland

Land capability (nonirrigated): 3e

Hydrologic Soil Group: C

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Typical profile

0 to 4 inches: Sandy loam

4 to 6 inches: Sandy loam

6 to 35 inches: Clay

35 to 80 inches: Clay

Minor Components

Shubuta

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Dothan

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

38—Bonneau-Norfolk-Angie complex, 5 to 12 percent slopes**Map Unit Setting**

Elevation: 30 to 700 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Bonneau and similar soils: 35 percent
Norfolk and similar soils: 30 percent
Angie and similar soils: 20 percent
Minor components: 15 percent

Description of Bonneau

Setting

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Convex

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 5 to 12 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 42 to 60 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 6.5 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 3s

Hydrologic Soil Group: B

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP144FL)

Typical profile

0 to 5 inches: Loamy sand

5 to 25 inches: Loamy sand

25 to 68 inches: Sandy clay loam

68 to 80 inches: Fine sandy loam

Description of Norfolk

Setting

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy marine deposits

Properties and qualities

Slope: 5 to 10 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 48 to 72 inches

Frequency of flooding: None

Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3e
Hydrologic Soil Group: B
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Typical profile

0 to 5 inches: Loamy sand
5 to 15 inches: Loamy sand
15 to 62 inches: Sandy clay loam
62 to 80 inches: Sandy loam

Description of Angie

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy and clayey marine deposits

Properties and qualities

Slope: 5 to 12 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.9 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 6e
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Typical profile

0 to 4 inches: Sandy loam
4 to 6 inches: Sandy loam
6 to 35 inches: Clay
35 to 80 inches: Clay

Minor Components

Cowarts

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Bonifay

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of mesic uplands (G133AA123FL), Unnamed (G133AP141FL)

Shubuta

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on ridges and side slopes of mesic uplands (G133AA312FL), Unnamed (G133AP143FL)

Orangeburg

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP144FL)

39—Pantego loam, depressional

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Pantego and similar soils: 75 percent

Minor components: 25 percent

Description of Pantego

Setting

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 0 to 18 inches

Frequency of flooding: None

Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: High (about 9.9 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 6w

Hydrologic Soil Group: B/D

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

Typical profile

0 to 17 inches: Loam

17 to 35 inches: Sandy clay loam

35 to 80 inches: Sandy clay loam

Minor Components

Escambia

Percent of map unit: 5 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Leefield

Percent of map unit: 5 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Pamlico

Percent of map unit: 4 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Kinston

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Johnston

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Bibb

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Albany

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

40—Escambia sandy loam, 0 to 2 percent slopes

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Escambia and similar soils: 80 percent

Minor components: 20 percent

Description of Escambia

Setting

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately low to moderately high (0.06 to 0.57 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: High (about 9.9 inches)

Interpretive groups

Farmland classification: Prime farmland if drained

Land capability (nonirrigated): 2w

Hydrologic Soil Group: C

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Typical profile

0 to 5 inches: Sandy loam
5 to 17 inches: Sandy loam
17 to 67 inches: Sandy loam
67 to 80 inches: Loam

Minor Components**Albany**

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Floralia

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Malbis

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Leefield

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Pantego

Percent of map unit: 2 percent
Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

Dothan

Percent of map unit: 2 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Fuquay

Percent of map unit: 2 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Stilson

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

41—Maurepas muck, frequently flooded**Map Unit Setting**

Elevation: 0 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Maurepas and similar soils: 75 percent
Minor components: 25 percent

Description of Maurepas**Setting**

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Woody organic material

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (2.00 to 19.98 in/hr)
Depth to water table: About 0 inches
Frequency of flooding: Frequent
Frequency of ponding: Frequent
Maximum salinity: Nonsaline to very slightly saline (0.0 to 4.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very high (about 20.9 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 8
Hydrologic Soil Group: A/D
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

Typical profile

0 to 65 inches: Muck

Minor Components

Bibb

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Kinston

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Chipley

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Foxworth

Percent of map unit: 4 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Pamlico

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Concave

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Rutlege

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave

Across-slope shape: Linear, concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

42—Blanton sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Blanton and similar soils: 80 percent

Minor components: 20 percent

Description of Blanton

Setting

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 48 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very low (about 3.0 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Typical profile

0 to 6 inches: Sand
6 to 65 inches: Fine sand
65 to 70 inches: Loamy fine sand
70 to 80 inches: Fine sandy loam

Minor Components

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Bonifay

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Troup

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Lakeland

Percent of map unit: 3 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

Albany

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Chipley

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Fuquay

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Leefield

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

43—Kinston-Bibb association, frequently flooded

Map Unit Setting

Elevation: 30 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Kinston and similar soils: 45 percent

Bibb and similar soils: 40 percent

Minor components: 15 percent

Description of Kinston

Setting

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Loamy alluvium

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: High (about 9.7 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 6w

Hydrologic Soil Group: B/D

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Typical profile

0 to 10 inches: Loam

10 to 50 inches: Sandy clay loam

50 to 80 inches: Clay loam

Description of Bibb

Setting

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Loamy and sandy alluvium

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 6 to 18 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 5w
Hydrologic Soil Group: B/D
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Typical profile

0 to 12 inches: Loam
12 to 37 inches: Sandy loam
37 to 65 inches: Loamy sand

Minor Components

Pantego

Percent of map unit: 3 percent
Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

Pamlico

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Concave

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Garcon

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Bigbee

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces, stream terraces on marine terraces

Landform position (three-dimensional): Tread, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Johnston

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

44—Lakeland-Troup-Urban land complex, 0 to 5 percent slopes

Map Unit Setting

Elevation: 40 to 300 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Lakeland and similar soils: 40 percent

Troup and similar soils: 25 percent

Urban land: 20 percent

Minor components: 15 percent

Description of Lakeland

Setting

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Eolian or sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 3.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 4s
Hydrologic Soil Group: A
Other vegetative classification: Forage suitability group not assigned (G133AA999FL), Unnamed (G133AP192FL)

Typical profile

0 to 4 inches: Sand
4 to 80 inches: Sand

Description of Troup

Setting

Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 5.1 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 3s
Hydrologic Soil Group: A

Other vegetative classification: Forage suitability group not assigned
(G133AA999FL), Unnamed (G133AP141FL)

Typical profile

0 to 7 inches: Sand
7 to 51 inches: Loamy sand
51 to 55 inches: Sandy loam
55 to 80 inches: Sandy clay loam

Description of Urban Land

Setting

Landform: Marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: No parent material

Interpretive groups

Farmland classification: Not prime farmland
Other vegetative classification: Forage suitability group not assigned
(G133AA999FL), Unnamed (G133AP900FL)

Minor Components

Bonifay

Percent of map unit: 10 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Forage suitability group not assigned
(G133AA999FL), Unnamed (G133AP141FL)

Fuquay

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Forage suitability group not assigned
(G133AA999FL), Unnamed (G133AP140FL)

Lucy

Percent of map unit: 2 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Forage suitability group not assigned
(G133AA999FL), Unnamed (G133AP140FL)

45—Dirego muck, frequently flooded

Map Unit Setting

Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Dirego and similar soils: 85 percent
Minor components: 15 percent

Description of Dirego

Setting

Landform: Tidal marshes on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Herbaceous organic material over sandy marine deposits

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Very poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: Frequent
Frequency of ponding: None
Maximum salinity: Moderately saline to strongly saline (16.0 to 32.0 mmhos/cm)
Sodium adsorption ratio, maximum: 55.0
Available water capacity: Very low (about 1.2 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 8
Hydrologic Soil Group: A/D
Other vegetative classification: Forage suitability group not assigned (G133AA999FL), Unnamed (G152AP850FL)

Typical profile

0 to 48 inches: Muck
48 to 65 inches: Fine sand

Minor Components

Maurepas

Percent of map unit: 12 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf

Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

Bigbee

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

46—Norfolk loamy sand, 2 to 5 percent slopes**Map Unit Setting**

Elevation: 30 to 700 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Norfolk and similar soils: 80 percent
Minor components: 20 percent

Description of Norfolk**Setting**

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 2 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 48 to 72 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.1 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 2e
Hydrologic Soil Group: B

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Typical profile

0 to 3 inches: Loamy sand
3 to 15 inches: Loamy sand
15 to 62 inches: Sandy clay loam
62 to 80 inches: Sandy loam

Minor Components**Angie**

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Cowarts

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP143FL)

Bibb

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Pamlico

Percent of map unit: 2 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Bonifay

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Floralia

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

47—Bonneau loamy sand, 0 to 5 percent slopes**Map Unit Setting**

Elevation: 40 to 700 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Bonneau and similar soils: 80 percent
Minor components: 20 percent

Description of Bonneau**Setting**

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 42 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 6.5 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 2s
Hydrologic Soil Group: B
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Typical profile

0 to 4 inches: Loamy sand
4 to 25 inches: Loamy sand
25 to 68 inches: Sandy clay loam
68 to 80 inches: Sandy clay loam

Minor Components

Angie

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Bonifay

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP141FL)

Fuquay

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 3 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Floral

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Cowarts

Percent of map unit: 2 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP143FL)

Troup

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

Leefield

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

48—Yemassee-Garcon-Bigbee complex, occasionally flooded

Map Unit Setting

Elevation: 0 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Yemassee and similar soils: 34 percent
Garcon and similar soils: 25 percent
Bigbee and similar soils: 20 percent
Minor components: 21 percent

Description of Yemassee

Setting

Landform: Stream terraces on marine terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 12 to 18 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.4 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 2w
Hydrologic Soil Group: B/D
Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Typical profile

0 to 8 inches: Loamy sand
8 to 17 inches: Loamy sand

17 to 50 inches: Sandy clay loam
50 to 80 inches: Sand

Description of Garcon

Setting

Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 18 to 36 inches
Frequency of flooding: Occasional
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 6.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 2w
Hydrologic Soil Group: A
Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Typical profile

0 to 18 inches: Loamy fine sand
18 to 28 inches: Loamy fine sand
28 to 51 inches: Sandy clay loam
51 to 80 inches: Sand

Description of Bigbee

Setting

Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy fluviomarine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: About 42 to 72 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 4.3 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 3s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Typical profile

0 to 8 inches: Loamy sand

8 to 80 inches: Sand

Minor Components

Mandarin

Percent of map unit: 3 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Kinston

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Johnston

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Bibb

Percent of map unit: 3 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

Pamlico

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Leon

Percent of map unit: 2 percent
Landform: Flats on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 2 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Pantego

Percent of map unit: 2 percent
Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

49—Eglin sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 300 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Eglin and similar soils: 80 percent

Minor components: 20 percent

Description of Eglin**Setting**

Landform: Marine terraces, rises

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat excessively drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 5.95 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 4.8 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 4s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

Typical profile

0 to 2 inches: Sand

2 to 68 inches: Sand

68 to 80 inches: Sand

Minor Components**Lakeland**

Percent of map unit: 7 percent

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP192FL)

Hurricane

Percent of map unit: 5 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Foxworth

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Chipley

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

50—Mandarin sand

Map Unit Setting

Elevation: 0 to 300 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Mandarin and similar soils: 80 percent
Minor components: 20 percent

Description of Mandarin

Setting

Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.57 to 1.98 in/hr)
Depth to water table: About 18 to 42 inches

Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 4.3 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 6s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Typical profile

0 to 8 inches: Sand
8 to 21 inches: Sand
21 to 38 inches: Sand
38 to 80 inches: Sand

Minor Components

Hurricane

Percent of map unit: 5 percent
Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Chipley

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Foxworth

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Resota

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G152AP142FL)

Leon

Percent of map unit: 2 percent
Landform: Flats on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 2 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

51—Bigbee loamy sand, 0 to 5 percent slopes, occasionally flooded**Map Unit Setting**

Elevation: 0 to 250 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Bigbee and similar soils: 80 percent
Minor components: 20 percent

Description of Bigbee**Setting**

Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy fluviomarine deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: About 42 to 72 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 4.4 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 3s

Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Typical profile

0 to 8 inches: Loamy sand

8 to 23 inches: Loamy sand

23 to 80 inches: Sand

Minor Components

Garcon

Percent of map unit: 7 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Yemassee

Percent of map unit: 5 percent

Landform: Stream terraces on marine terraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Blanton

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluvial

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Mandarin

Percent of map unit: 3 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

52—Yemassee fine sandy loam, occasionally flooded**Map Unit Setting**

Elevation: 0 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Yemassee and similar soils: 75 percent

Minor components: 25 percent

Description of Yemassee**Setting**

Landform: Stream terraces on marine terraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 12 to 18 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 8.7 inches)

Interpretive groups

Farmland classification: Prime farmland if drained

Land capability (nonirrigated): 2w

Hydrologic Soil Group: B/D

Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Typical profile

0 to 6 inches: Fine sandy loam
6 to 47 inches: Sandy clay loam
47 to 80 inches: Fine sandy loam

Minor Components**Garcon**

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Bigbee

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Kinston

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Pantego

Percent of map unit: 3 percent
Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

Floral

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Stilson

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Leefield

Percent of map unit: 3 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

53—Arents, 2 to 8 percent slopes

Map Unit Setting

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Arents and similar soils: 100 percent

Description of Arents

Setting

Landform: Rises on marine terraces

Landform position (three-dimensional): Rise

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Altered marine deposits

Properties and qualities

Slope: 2 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Capacity of the most limiting layer to transmit water (Ksat): Very high (19.98 to 50.02 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very low (about 2.4 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7s
Hydrologic Soil Group: A
Other vegetative classification: Forage suitability group not assigned
 (G133AA999FL), Unnamed (G152AP900FL)

Typical profile

0 to 80 inches: Sand

54—Newhan-Corolla sands, rolling

Map Unit Setting

Elevation: 0 to 300 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Newhan and similar soils: 45 percent
Corolla and similar soils: 35 percent
Minor components: 20 percent

Description of Newhan

Setting

Landform: Dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy eolian deposits

Properties and qualities

Slope: 2 to 30 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): Very high
 (19.98 to 50.02 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Very slightly saline to slightly saline (4.0 to 8.0
 mmhos/cm)
Sodium adsorption ratio, maximum: 20.0
Available water capacity: Very low (about 1.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7s
Hydrologic Soil Group: A

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G152AP198FL)

Typical profile

0 to 5 inches: Sand
5 to 64 inches: Sand

Description of Corolla**Setting**

Landform: Rises on dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy marine deposits

Properties and qualities

Slope: 2 to 6 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Capacity of the most limiting layer to transmit water (Ksat): Very high (19.98 to 50.02 in/hr)
Depth to water table: About 18 to 36 inches
Frequency of flooding: Rare
Frequency of ponding: None
Maximum salinity: Slightly saline to moderately saline (8.0 to 16.0 mmhos/cm)
Sodium adsorption ratio, maximum: 20.0
Available water capacity: Very low (about 1.2 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7s
Hydrologic Soil Group: A/D
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP180FL)

Typical profile

0 to 5 inches: Sand
5 to 57 inches: Sand
57 to 80 inches: Sand

Minor Components**Kureb**

Percent of map unit: 5 percent
Landform: Dunes on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G152AP192FL)

Resota

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G152AP142FL)

Mandarin

Percent of map unit: 5 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Leon

Percent of map unit: 3 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave

Across-slope shape: Linear, concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

55—Beaches**Map Unit Setting**

Elevation: 0 to 40 feet

Mean annual precipitation: 42 to 70 inches

Mean annual air temperature: 52 to 70 degrees F

Frost-free period: 190 to 262 days

Map Unit Composition

Beaches: 90 percent

Minor components: 10 percent

Description of Beaches

Setting

Landform: Beaches on marine terraces
Landform position (three-dimensional): Rise
Down-slope shape: Convex
Across-slope shape: Linear

Properties and qualities

Slope: 1 to 5 percent
Drainage class: Poorly drained
Depth to water table: About 0 to 72 inches
Frequency of flooding: Very frequent

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 8
Other vegetative classification: Forage suitability group not assigned
 (G133AA999FL), Unnamed (G133AP900FL)

Minor Components

Newhan

Percent of map unit: 5 percent
Landform: Dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of
 xeric uplands (G133AA111FL), Unnamed (G152AP198FL)

Corolla

Percent of map unit: 5 percent
Landform: Rises on dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of
 mesic uplands (G133AA131FL), Unnamed (G152AP180FL)

56—Kureb sand, hilly

Map Unit Setting

Elevation: 10 to 300 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Kureb and similar soils: 80 percent
Minor components: 20 percent

Description of Kureb

Setting

Landform: Dunes on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Eolian deposits or sandy fluvial or marine deposits

Properties and qualities

Slope: 8 to 20 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Excessively drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very low (about 1.8 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 7s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G152AP195FL)

Typical profile

0 to 2 inches: Sand
2 to 45 inches: Sand
45 to 80 inches: Sand

Minor Components

Lakeland

Percent of map unit: 10 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G133AP195FL)

Newhan

Percent of map unit: 5 percent
Landform: Dunes on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Sandy soils on strongly sloping to steep side slopes of xeric uplands (G133AA113FL), Unnamed (G152AP198FL)

Resota

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G152AP142FL)

57—Hurricane sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 10 to 350 feet

Mean annual precipitation: 59 to 70 inches

Mean annual air temperature: 63 to 72 degrees F

Frost-free period: 223 to 306 days

Map Unit Composition

Hurricane and similar soils: 90 percent

Minor components: 10 percent

Description of Hurricane

Setting

Landform: Flats on marine terraces, rises on marine terraces

Landform position (two-dimensional): Footslope, toeslope

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): High (2.00 to 6.00 in/hr)

Depth to water table: About 24 to 42 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 3.4 inches)

Interpretive groups

Farmland classification: Farmland of local importance

Land capability classification (irrigated): 3w

Land capability (nonirrigated): 3w

Hydrologic Soil Group: A

Ecological site: North Florida Flatwoods (R138XY004FL)

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G152AA131FL), Unnamed (G152AT080FL)

Typical profile

0 to 5 inches: Sand

5 to 55 inches: Sand

55 to 80 inches: Sand

Minor Components

Leon

Percent of map unit: 3 percent

Landform: Marine terraces, flatwoods

Landform position (three-dimensional): Tread, talf

Down-slope shape: Linear, convex

Across-slope shape: Linear

Ecological site: North Florida Flatwoods (R152AY004FL)

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AT013FL)

Albany

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (two-dimensional): Shoulder, summit

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Linear, convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G133AP068FL)

Pottsburg, non-hydric

Percent of map unit: 2 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on flats of mesic or hydric lowlands (G152AA141FL), Unnamed (G152AT003FL)

Lutterloh

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G152AA131FL), Unnamed (G152AT080FL)

58—Duckston muck, frequently flooded

Map Unit Setting

Elevation: 0 to 300 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Duckston and similar soils: 80 percent

Minor components: 20 percent

Description of Duckston

Setting

Landform: Depressions on marine terraces, swales on marine terraces, flats on marine terraces

Landform position (three-dimensional): Dip, talf

Down-slope shape: Concave, linear

Across-slope shape: Concave, linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: Frequent

Frequency of ponding: None

Calcium carbonate, maximum content: 5 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 4.0 mmhos/cm)

Sodium adsorption ratio, maximum: 20.0

Available water capacity: Low (about 3.4 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 7w

Hydrologic Soil Group: A/D

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP003FL)

Typical profile

0 to 4 inches: Muck

4 to 21 inches: Sand

21 to 80 inches: Sand

Minor Components

Dirego

Percent of map unit: 10 percent

Landform: Tidal marshes on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

Leon

Percent of map unit: 5 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 5 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave

Across-slope shape: Linear, concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

59—Malbis fine sandy loam, 0 to 2 percent slopes

Map Unit Setting

Elevation: 100 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Malbis and similar soils: 75 percent

Minor components: 25 percent

Description of Malbis

Setting

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 30 to 48 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.0 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 1
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Typical profile

0 to 7 inches: Fine sandy loam
7 to 13 inches: Fine sandy loam
13 to 45 inches: Sandy clay loam
45 to 80 inches: Sandy clay loam

Minor Components**Dothan**

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Fuquay

Percent of map unit: 4 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Tifton

Percent of map unit: 4 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Stilson

Percent of map unit: 4 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Escambia

Percent of map unit: 4 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Leefield

Percent of map unit: 4 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

60—Malbis fine sandy loam, 2 to 5 percent slopes

Map Unit Setting

Elevation: 100 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Malbis and similar soils: 75 percent
Minor components: 25 percent

Description of Malbis

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve

Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 2 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 30 to 48 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.5 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 2e
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP328FL)

Typical profile

0 to 6 inches: Fine sandy loam
6 to 38 inches: Fine sandy loam
38 to 48 inches: Sandy clay loam
48 to 80 inches: Fine sandy loam

Minor Components

Dothan

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Tifton

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 5 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Escambia

Percent of map unit: 5 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Stilson

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Leefield

Percent of map unit: 2 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

61—Malbis fine sandy loam, 5 to 8 percent slopes**Map Unit Setting**

Elevation: 100 to 400 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Malbis and similar soils: 75 percent

Minor components: 25 percent

Description of Malbis

Setting

Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Loamy marine deposits

Properties and qualities

Slope: 5 to 8 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 30 to 48 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.9 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 3e
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP328FL)

Typical profile

0 to 7 inches: Fine sandy loam
7 to 17 inches: Fine sandy loam
17 to 60 inches: Sandy clay loam
60 to 80 inches: Sandy clay loam

Minor Components

Dothan

Percent of map unit: 12 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Tifton

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Fuquay

Percent of map unit: 5 percent

Landform: Ridges on marine terraces

Landform position (three-dimensional): Side slope, interflue

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Escambia

Percent of map unit: 3 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interflue, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

62—Resota sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 20 to 250 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Resota and similar soils: 85 percent

Minor components: 15 percent

Description of Resota

Setting

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interflue

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy marine deposits

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very high (19.98 to 50.02 in/hr)

Depth to water table: About 42 to 60 inches

Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Very low (about 2.4 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 6s
Hydrologic Soil Group: A
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G152AP142FL)

Typical profile

0 to 3 inches: Sand
3 to 13 inches: Sand
13 to 53 inches: Sand
53 to 80 inches: Sand

Minor Components**Foxworth**

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Mandarin

Percent of map unit: 5 percent
Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP077FL)

Kureb

Percent of map unit: 5 percent
Landform: Dunes on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G152AP192FL)

63—Pickney sand, depressional

Map Unit Setting

Elevation: 0 to 300 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Pickney and similar soils: 80 percent

Minor components: 20 percent

Description of Pickney

Setting

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Sandy marine deposits and/or fluvio-marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: None

Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 4.2 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 6w

Hydrologic Soil Group: A/D

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G133AP800FL)

Typical profile

0 to 37 inches: Sand

37 to 80 inches: Sand

Minor Components

Leon

Percent of map unit: 5 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

Rutlege

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear, concave
Across-slope shape: Linear, concave
Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Hurricane

Percent of map unit: 5 percent
Landform: Flats on marine terraces, rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Pamlico

Percent of map unit: 5 percent
Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

64—Pamlico muck**Map Unit Setting**

Elevation: 0 to 300 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Pamlico and similar soils: 80 percent
Minor components: 20 percent

Description of Pamlico**Setting**

Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave

Parent material: Herbaceous organic material over sandy marine deposits

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 5.95 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: Rare

Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Very high (about 13.3 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 7w

Hydrologic Soil Group: A/D

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G152AP850FL)

Typical profile

0 to 25 inches: Muck

25 to 65 inches: Sand

Minor Components

Dorovan

Percent of map unit: 7 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Organic soils in depressions and on flood plains (G133AA645FL), Unnamed (G133AP850FL)

Pickney

Percent of map unit: 5 percent

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G133AP800FL)

Rutlege

Percent of map unit: 5 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave

Across-slope shape: Linear, concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Leon

Percent of map unit: 3 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP013FL)

65—Garcon loamy fine sand, occasionally flooded

Map Unit Setting

Elevation: 0 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Garcon and similar soils: 70 percent

Minor components: 30 percent

Description of Garcon

Setting

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.57 to 1.98 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: Occasional

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 6.0 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 2w

Hydrologic Soil Group: A

Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Typical profile

0 to 6 inches: Loamy fine sand
6 to 25 inches: Loamy fine sand
25 to 46 inches: Fine sandy loam
46 to 80 inches: Fine sand

Minor Components**Yemassee**

Percent of map unit: 10 percent
Landform: Stream terraces on marine terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Bigbee

Percent of map unit: 10 percent
Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Blanton

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Kinston

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Pantego

Percent of map unit: 2 percent

Landform: Depressions on marine terraces
Landform position (three-dimensional): Dip
Down-slope shape: Concave
Across-slope shape: Concave
Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

66—Kenansville loamy fine sand, 0 to 5 percent slopes

Map Unit Setting

Elevation: 0 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Kenansville and similar soils: 80 percent
Minor components: 20 percent

Description of Kenansville

Setting

Landform: Stream terraces on marine terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Sandy and loamy marine or fluvial deposits

Properties and qualities

Slope: 0 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 4.4 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 2s
Hydrologic Soil Group: A
Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

Typical profile

0 to 10 inches: Loamy fine sand
10 to 31 inches: Loamy fine sand

31 to 57 inches: Fine sandy loam
57 to 80 inches: Fine sand

Minor Components

Garcon

Percent of map unit: 5 percent
Landform: Flood plains on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Other vegetative classification: Sandy or sandy over loamy soils on stream terraces or flood plains (G133AA134FL), Unnamed (G133AP078FL)

Yemassee

Percent of map unit: 5 percent
Landform: Stream terraces on marine terraces
Landform position (three-dimensional): Tread
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Bonneau

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Bigbee

Percent of map unit: 3 percent
Landform: Flood plains on marine terraces, stream terraces on marine terraces
Landform position (three-dimensional): Tread, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy soils on stream terraces or flood plains (G133AA114FL), Unnamed (G133AP092FL)

Norfolk

Percent of map unit: 2 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Troup

Percent of map unit: 2 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on ridges and dunes of xeric uplands (G133AA111FL), Unnamed (G133AP141FL)

68—Floralia loamy fine sand, 0 to 2 percent slopes**Map Unit Setting**

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Floralia and similar soils: 80 percent

Minor components: 20 percent

Description of Floralia**Setting**

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy fluviomarine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Moderate (about 7.5 inches)

Interpretive groups

Farmland classification: Prime farmland if drained

Land capability (nonirrigated): 2w

Hydrologic Soil Group: C

Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP075FL)

Typical profile

0 to 6 inches: Loamy fine sand
6 to 10 inches: Loamy fine sand
10 to 30 inches: Fine sandy loam
30 to 80 inches: Sandy clay loam

Minor Components**Escambia**

Percent of map unit: 3 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Fuquay

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP137FL)

Stilson

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Albany

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP065FL)

Leefield

Percent of map unit: 3 percent

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Pantego

Percent of map unit: 2 percent

Landform: Depressions on marine terraces

Landform position (three-dimensional): Dip

Down-slope shape: Concave

Across-slope shape: Concave

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP800FL)

69—Floralia loamy fine sand, 2 to 5 percent slopes

Map Unit Setting

Elevation: 50 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Floralia and similar soils: 75 percent

Minor components: 25 percent

Description of Floralia

Setting

Landform: Rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Loamy fluviomarine deposits

Properties and qualities

Slope: 2 to 5 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 18 to 30 inches

Frequency of flooding: None

Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 7.0 inches)

Interpretive groups

Farmland classification: Prime farmland if drained
Land capability (nonirrigated): 2e
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Typical profile

0 to 8 inches: Loamy fine sand
8 to 17 inches: Loamy fine sand
17 to 39 inches: Fine sandy loam
39 to 80 inches: Fine sandy loam

Minor Components

Escambia

Percent of map unit: 5 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve, talf
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP325FL)

Dothan

Percent of map unit: 5 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Leefield

Percent of map unit: 5 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Fuquay

Percent of map unit: 3 percent
Landform: Ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP140FL)

Stilson

Percent of map unit: 3 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on rises and knolls of mesic uplands (G133AA231FL), Unnamed (G133AP078FL)

Kinston

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP201FL)

Bibb

Percent of map unit: 2 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces, flood plains, or in depressions (G133AA345FL), Unnamed (G133AP015FL)

70—Shubuta fine sandy loam, 2 to 5 percent slopes

Map Unit Setting

Elevation: 30 to 600 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Shubuta and similar soils: 75 percent

Minor components: 20 percent

Description of Shubuta

Setting

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Parent material: Clayey marine deposits

Properties and qualities

Slope: 2 to 5 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: High (about 9.4 inches)

Interpretive groups

Farmland classification: All areas are prime farmland
Land capability (nonirrigated): 2e
Hydrologic Soil Group: C
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Typical profile

0 to 6 inches: Fine sandy loam
6 to 11 inches: Fine sandy loam
11 to 34 inches: Clay
34 to 80 inches: Sandy clay

Minor Components

Angie

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Orangeburg

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on knolls and ridges of mesic uplands (G133AA311FL), Unnamed (G133AP140FL)

Dothan

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 3 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises and knolls of mesic uplands (G133AA321FL), Unnamed (G133AP140FL)

Floral

Percent of map unit: 2 percent
Landform: Rises on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on flats of hydric or mesic lowlands (G133AA331FL), Unnamed (G133AP078FL)

Bonneau

Percent of map unit: 2 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP141FL)

Lucy

Percent of map unit: 2 percent
Landform: Ridges on marine terraces
Landform position (three-dimensional): Interfluve
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

71—Shubuta fine sandy loam, 5 to 12 percent slopes

Map Unit Setting

Elevation: 30 to 700 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Shubuta and similar soils: 70 percent

Minor components: 30 percent

Description of Shubuta

Setting

Landform: Ridges on marine terraces, hills on marine terraces

Landform position (three-dimensional): Interfluve, side slope

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Clayey marine deposits

Properties and qualities

Slope: 5 to 12 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: High (about 9.4 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 4e

Hydrologic Soil Group: C

Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Typical profile

0 to 5 inches: Fine sandy loam

5 to 11 inches: Fine sandy loam

11 to 46 inches: Clay

46 to 80 inches: Sandy clay

Minor Components

Cowarts

Percent of map unit: 10 percent

Landform: Ridges on marine terraces
Landform position (three-dimensional): Side slope, interfluvium
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP143FL)

Orangeburg

Percent of map unit: 5 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on strongly sloping to steep side slopes of mesic uplands (G133AA313FL), Unnamed (G133AP140FL)

Angie

Percent of map unit: 5 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Norfolk

Percent of map unit: 4 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear
Other vegetative classification: Loamy and clayey soils on rises, knolls, and ridges of mesic uplands (G133AA322FL), Unnamed (G133AP140FL)

Bonneau

Percent of map unit: 3 percent
Landform: Knolls on marine terraces, ridges on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Convex
Other vegetative classification: Sandy over loamy soils on rises, knolls, and ridges of mesic uplands (G133AA221FL), Unnamed (G133AP144FL)

Lucy

Percent of map unit: 3 percent
Landform: Ridges on marine terraces, hills on marine terraces
Landform position (three-dimensional): Interfluvium, side slope
Down-slope shape: Convex
Across-slope shape: Linear

Other vegetative classification: Sandy over loamy soils on knolls and ridges of mesic uplands (G133AA211FL), Unnamed (G133AP140FL)

72—Osier fine sand

Map Unit Setting

Elevation: 0 to 450 feet
Mean annual precipitation: 62 to 70 inches
Mean annual air temperature: 63 to 70 degrees F
Frost-free period: 232 to 262 days

Map Unit Composition

Osier and similar soils: 70 percent
Minor components: 30 percent

Description of Osier

Setting

Landform: Flats on marine terraces
Landform position (three-dimensional): Talf
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Sandy alluvium

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: About 0 to 12 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Low (about 4.2 inches)

Interpretive groups

Farmland classification: Not prime farmland
Land capability (nonirrigated): 5w
Hydrologic Soil Group: A/D
Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP002FL)

Typical profile

0 to 4 inches: Fine sand
4 to 80 inches: Fine sand

Minor Components

Chipley

Percent of map unit: 10 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Albany

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP065FL)

Yemassee

Percent of map unit: 5 percent

Landform: Stream terraces on marine terraces

Landform position (three-dimensional): Tread

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Loamy and clayey soils on stream terraces and flood plains (G133AA334FL), Unnamed (G133AP263FL)

Rutlege

Percent of map unit: 5 percent

Landform: Flood plains on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear, concave

Across-slope shape: Linear, concave

Other vegetative classification: Sandy soils on stream terraces, flood plains, or in depressions (G133AA145FL), Unnamed (G152AP800FL)

Hurricane

Percent of map unit: 5 percent

Landform: Flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

73—Albany loamy sand

Map Unit Setting

Elevation: 20 to 450 feet

Mean annual precipitation: 62 to 70 inches

Mean annual air temperature: 63 to 70 degrees F

Frost-free period: 232 to 262 days

Map Unit Composition

Albany and similar soils: 85 percent

Minor components: 15 percent

Description of Albany**Setting**

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Interfluve, talf

Down-slope shape: Convex

Across-slope shape: Linear

Parent material: Sandy and loamy marine deposits

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water

(Ksat): Moderately high to high (0.60 to 2.00 in/hr)

Depth to water table: About 12 to 30 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 2.0 mmhos/cm)

Sodium adsorption ratio, maximum: 4.0

Available water capacity: Low (about 3.3 inches)

Interpretive groups

Farmland classification: Not prime farmland

Land capability (nonirrigated): 3w

Hydrologic Soil Group: A/D

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP065FL)

Typical profile

0 to 11 inches: Loamy sand

11 to 45 inches: Loamy sand

45 to 80 inches: Sandy loam

Minor Components**Blanton**

Percent of map unit: 8 percent

Landform: Knolls on marine terraces, ridges on marine terraces

Landform position (three-dimensional): Side slope, interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises, knolls, and ridges of mesic uplands (G133AA121FL), Unnamed (G133AP142FL)

Chipley

Percent of map unit: 5 percent

Landform: Knolls on marine terraces, flats on marine terraces, rises on marine terraces

Landform position (three-dimensional): Interfluve

Down-slope shape: Convex

Across-slope shape: Linear

Other vegetative classification: Sandy soils on rises and knolls of mesic uplands (G133AA131FL), Unnamed (G152AP080FL)

Osier

Percent of map unit: 2 percent

Landform: Flats on marine terraces

Landform position (three-dimensional): Talf

Down-slope shape: Linear

Across-slope shape: Linear

Other vegetative classification: sandy soils on flats of mesic or hydric lowlands (G133AA141FL), Unnamed (G152AP002FL)

99—Water

Map Unit Composition

Water: 100 percent

Description of Water

Interpretive groups

Farmland classification: Not prime farmland

Other vegetative classification: Forage suitability group not assigned (G133AA999FL)

100—Waters of the Gulf of Mexico

Map Unit Composition

Waters of the gulf of mexico: 100 percent

Description of Waters Of The Gulf Of Mexico

Interpretive groups

Farmland classification: Not prime farmland

Other vegetative classification: Forage suitability group not assigned (G133AA999FL)

Data Source Information

Soil Survey Area: Walton County, Florida

Survey Area Data: Version 12, Dec 27, 2013