

Renville County, Minnesota

musym	muname	RUSLE2 slope factors	
		slope_%	slope_length_ft.
27A	Dickinson loam, 0 to 2 percent slopes	1	102
27B	Dickinson loam, 2 to 6 percent slopes	4	125
35	Blue Earth mucky silt loam, 0 to 1 percent slopes	1	102
39A	Wadena loam, 0 to 2 percent slopes	2	102
39B	Wadena loam, 2 to 6 percent slopes	4	125
85	Calco silty clay loam, 0 to 2 percent slopes, occasionally flooded	1	151
86	Canisteo clay loam, moderately fine substratum, 0 to 2 percent slopes	1	102
94C	Terril loam, 6 to 12 percent slopes	10	151
102B	Clarion loam, 2 to 6 percent slopes	3	98
112	Harps clay loam, 0 to 2 percent slopes	1	102
113	Webster clay loam, 0 to 2 percent slopes	1	102
118	Crippin loam, moderately fine substratum, 1 to 3 percent slopes	1	102
128C2	Grogan silt loam, 6 to 15 percent slopes, eroded	9	125
130	Nicollet clay loam, 1 to 3 percent slopes	2	151
134	Okoboji silty clay loam, depressionnal, 0 to 1 percent slopes	1	102
156	Fairhaven silt loam, 0 to 2 percent slopes	1	203
227	Lemond loam, 0 to 2 percent slopes	1	151
247	Linder loam, 0 to 2 percent slopes	1	151
255	Mayer loam, 0 to 2 percent slopes	1	161
282	Hanska loam, 0 to 2 percent slopes	1	151
318	Mayer clay loam, depressionnal, 0 to 1 percent slopes	1	49
327A	Dickman sandy loam, 0 to 2 percent slopes	1	203
327B	Dickman sandy loam, 2 to 6 percent slopes	4	151
327C	Dickman sandy loam, 6 to 12 percent slopes	9	125
336	Delft loam, 1 to 3 percent slopes	2	102
386	Okoboji mucky silty clay loam, depressionnal, 0 to 1 percent slopes	1	102
392	Biscay loam, 0 to 2 percent slopes	2	151
423	Seaforth loam, 1 to 3 percent slopes	2	102
446	Normania loam, 1 to 3 percent slopes	2	125
463A	Minneiska silt loam, 0 to 2 percent slopes, occasionally flooded	1	151
463B	Minneiska loam, 1 to 4 percent slopes, rarely flooded	3	151
519	Klossner muck, depressionnal, calcareous, 0 to 1 percent slopes	1	102

Renville County, Minnesota

musym	muname	RUSLE2 slope factors	
		slope_%	slope_length_ft.
525	Muskego soils, 0 to 1 percent slopes	0	689
574	Du Page loam, 0 to 2 percent slopes, occasionally flooded	1	151
575	Nishna silty clay, 0 to 2 percent slopes, occasionally flooded	1	102
595F	Swanlake loam, 18 to 50 percent slopes	35	125
610	Calco silty clay loam, 0 to 1 percent slopes, frequently flooded	1	102
770C2	Ves-Terril complex, 6 to 15 percent slopes, eroded	12	125
810	Coriff-Fieldon complex, 0 to 2 percent slopes	1	102
817	Canisteo-Seaforth complex, 0 to 3 percent slopes	1	102
875C	Hawick-Estherville complex, 6 to 12 percent slopes	9	125
887B	Clarion-Swanlake complex, 2 to 6 percent slopes	4	151
899	Harps-Okobojo, depressional, complex, 0 to 2 percent slopes	1	102
920B	Clarion-Storden-Hawick complex, 2 to 6 percent slopes	4	128
927	Harps-Seaforth-Okobojo, depressional, complex, 0 to 3 percent slopes	1	102
954C2	Ves-Storden complex, 6 to 12 percent slopes, eroded	9	151
956	Canisteo-Glencoe, depressional, complex, 0 to 2 percent slopes	1	102
960D2	Storden-Omsrud complex, 12 to 18 percent slopes, eroded	14	125
960F	Storden-Omsrud complex, 18 to 50 percent slopes	25	102
978	Cordova-Rolfe, depressional, complex, 0 to 2 percent slopes	1	102
999C2	Ves-Storden-Hawick complex, 6 to 12 percent slopes, eroded	9	151
1030	Pits, gravel-Udipsamments complex	10	26
1080	Klossner, okobojo, and glencoe soils, ponded, 0 to 1 percent slopes	1	102
1100	Nicollet silty clay loam, 1 to 3 percent slopes	2	102
1101	Webster silty clay loam, moderately fine substratum, 0 to 2 percent slopes	2	102
1205	Leen-Okobojo, depressional, complex, 0 to 2 percent slopes	1	102
1242F	Swanlake-Terril complex, 18 to 50 percent slopes	35	125
1261B	Bechyn loam, 2 to 6 percent slopes	4	125
1262	Seaforth silt loam, 1 to 3 percent slopes	2	102
1267	Cedarrock silty clay loam, 0 to 2 percent slopes, frequently flooded	1	151
1268	Hanlon loam, 1 to 3 percent slopes, rarely flooded	1	151
1269	Lowlein silt loam, 0 to 2 percent slopes	1	102
1270D	Bechyn-Rock outcrop complex, 0 to 40 percent slopes	20	102
1285	Chetomba silty clay loam, 0 to 2 percent slopes	1	102

Renville County, Minnesota

musym	muname	RUSLE2 slope factors	
		slope_%	slope_length_ft.
1286	Prinsburg silty clay loam, 0 to 2 percent slopes	1	102
1287	Calco silty clay loam, ponded, 0 to 1 percent slopes, frequently flooded	1	102
1355B	Amiret-Swanlake complex, 2 to 6 percent slopes	4	125
1369A	Crooksford silt loam, 1 to 3 percent slopes	2	151
1369B	Crooksford silt loam, 3 to 5 percent slopes	4	203
1370B	Amiret loam, 2 to 5 percent slopes	4	203
1371B	Crooksford-Swanlake complex, 3 to 6 percent slopes	4	151
1373C	Omsrud-Storden-Hawick complex, 6 to 12 percent slopes, eroded	9	125
1374	Havelock clay loam, 0 to 2 percent slopes, occasionally flooded	1	151
1375D	Storden-Ves complex, 12 to 18 percent slopes, eroded	14	125
1376C	Omsrud-Storden complex, 6 to 12 percent slopes, eroded	9	125
1382	Louris silt loam, 1 to 3 percent slopes	2	102
1386B	Amiret-Swanlake-Hawick complex, 2 to 6 percent slopes	4	151
1388B	Terril loam, moderately wet, 2 to 6 percent slopes	5	151
1389	Havelock silt loam, 0 to 2 percent slopes, frequently flooded	1	102
1390	Leen silty clay loam, 0 to 2 percent slopes	1	102
1392B	Grogan silt loam, moderately wet, 1 to 4 percent slopes	4	151
1802	Calcousta-Okoboji complex, depressional, 0 to 1 percent slopes	1	102
1833	Coland clay loam, 0 to 2 percent slopes, occasionally flooded	1	151
1834	Coland clay loam, 0 to 2 percent slopes, frequently flooded	1	102
1845A	Estherville loam, 0 to 2 percent slopes	2	102
1845B	Estherville loam, 2 to 6 percent slopes	4	125
1900	Okoboji-Canisteo complex, depressional, 0 to 1 percent slopes	1	102
1917	Nishna silty clay loam, 0 to 1 percent slopes, frequently flooded	1	102
1958	Danube silty clay, 0 to 2 percent slopes	1	203
1999	Minneiska-Rushriver complex, 0 to 2 percent slopes, frequently flooded	1	102
L13A	Klossner muck, 0 to 1 percent slopes	0	689
L33A	Kandiyohi clay, 0 to 2 percent slopes	1	151
L33B	Kandiyohi clay, 2 to 5 percent slopes	3	151
L34A	Cosmos silty clay, 0 to 2 percent slopes	1	102
L83A	Webster clay loam, 0 to 2 percent slopes	1	125
L85A	Nicollet clay loam, 1 to 3 percent slopes	2	125

Renville County, Minnesota

musym	muname	RUSLE2 slope factors	
		slope_%	slope_length_ft.
L107A	Canisteo-Glencoe, depressional complex, 0 to 2 percent slopes	1	125
L163A	Okoboji silty clay loam, depressional, 0 to 1 percent slopes	1	125
L164A	Lura silty clay, depressional, firm substratum, 0 to 1 percent slopes	1	102
L166C2	Newlondon-Strout complex, 6 to 12 percent slopes, moderately eroded	9	125
L179A	Corvuso-Lura, depressional, firm substratum complex, 0 to 2 percent slopes	1	102
L184A	Corvuso silty clay loam, 0 to 2 percent slopes	1	102
L185B	Strout-Arkton complex, 2 to 6 percent slopes	4	151
L187A	Klossner and Lura soils, ponded, firm substratum, 0 to 1 percent slopes	1	102
L192A	Okoboji mucky silty clay loam, depressional, firm substratum, 0 to 1 percent slopes	1	102
L200A	Klossner muck, depressional, firm substratum, 0 to 1 percent slopes	1	125
L201A	Normania loam, 1 to 3 percent slopes	2	125
L204C2	Newlondon-Strout-Hawick complex, 6 to 12 percent slopes, moderately eroded	9	125
L206B	Strout-Arkton-Estherville complex, 2 to 6 percent slopes	4	125
M-W	Water, miscellaneous		0
W	Water		0