

**Wright County Crop Productivity Index (CPI) by Map Unit Name**

| Mapunit Symbol | Map Unit Name   | CPI |
|----------------|---|-----|
| 441            | Almora loam, 0 to 2 percent slopes                                    | 78  |
| 1036B          | Angus fine sandy loam, 2 to 5 percent slopes                          | 90  |
| 1362B          | Angus loam, 2 to 5 percent slopes                                     | 90  |
| 1094B          | Angus-Cordova complex, 0 to 5 percent slopes                          | 89  |
| 1408B          | Angus-Kilkenny complex, 2 to 6 percent slopes                         | 83  |
| 1901B          | Angus-Le Sueur complex, 1 to 5 percent slopes                         | 94  |
| 1087B          | Angus-Malardi complex, 2 to 6 percent slopes                          | 75  |
| 1000           | Arvilla sandy loam, map >25, 0 to 2 percent slopes                    | 44  |
| 1161           | Barry loam, 0 to 2 percent slopes                                     | 87  |
| L317A          | Barry loam, 0 to 2 percent slopes                                     | 87  |
| 1443           | Belleville sandy loam, 0 to 2 percent slopes                          | 30  |
| 392            | Biscay loam, 0 to 2 percent slopes                                    | 69  |
| 35             | Blue Earth mucky silty clay loam, depressional, 0 to 1 percent slopes | 77  |
| 169C           | Braham loamy fine sand, 6 to 12 percent slopes                        | 52  |
| 1438B          | Braham loamy fine sand, moderately wet, 2 to 5 percent slopes         | 60  |
| 1380A          | Bygland silt loam, map >25, 0 to 2 percent slopes                     | 86  |
| 1380B          | Bygland silt loam, map >25, 2 to 6 percent slopes                     | 86  |
| 1380C          | Bygland silt loam, map >25, 6 to 12 percent slopes, eroded            | 71  |
| 86             | Canisteo clay loam, moderately fine substratum, 0 to 2 percent slopes | 93  |
| 956            | Canisteo-Glencoe, depressional, complex, 0 to 2 percent slopes        | 91  |
| 1256           | Cantlin loamy fine sand, 0 to 3 percent slopes                        | 43  |
| 1213D          | Cokato-Storden complex, 12 to 18 percent slopes, eroded               | 71  |
| 1213C          | Cokato-Storden complex, 6 to 12 percent slopes, eroded                | 84  |
| 1220C          | Cokato-Storden-Hawick complex, 6 to 12 percent slopes, eroded         | 77  |
| 109            | Cordova clay loam, 0 to 2 percent slopes                              | 87  |
| 1156           | Cordova loam, 0 to 2 percent slopes                                   | 87  |
| 978            | Cordova-Rolfe, depressional, complex, 0 to 2 percent slopes           | 86  |
| 1035B          | Crowfork loamy sand, 1 to 6 percent slopes                            | 43  |
| 1035D          | Crowfork loamy sand, 12 to 18 percent slopes                          | 32  |
| 1035C          | Crowfork loamy sand, 6 to 12 percent slopes                           | 41  |
| 74B            | Dickinson fine sandy loam, 1 to 6 percent slopes                      | 60  |
| 327A           | Dickman sandy loam, 0 to 2 percent slopes                             | 49  |
| 327B           | Dickman sandy loam, 2 to 6 percent slopes                             | 49  |
| 406            | Dorset sandy loam, 0 to 2 percent slopes                              | 46  |
| 1379B          | Dorset-Almora complex, 1 to 4 percent slopes                          | 55  |
| 1377D          | Dorset-Two Inlets complex, 12 to 20 percent slopes                    | 34  |
| 1377B          | Dorset-Two Inlets complex, 2 to 6 percent slopes                      | 45  |
| 1377E          | Dorset-Two Inlets complex, 20 to 35 percent slopes                    | 13  |
| 1377C          | Dorset-Two Inlets complex, 6 to 12 percent slopes                     | 44  |
| 260            | Duelm loamy sand, 0 to 2 percent slopes                               | 42  |
| 1037           | Eckvoll loamy fine sand, map >25, 0 to 3 percent slopes               | 60  |
| 1255           | Elkriver fine sandy loam, 0 to 2 percent slopes, occasionally flooded | 62  |
| 771            | Elkriver fine sandy loam, 0 to 2 percent slopes, rarely flooded       | 72  |
| 1257           | Elkriver-Mosford complex, 0 to 6 percent slopes, rarely flooded       | 69  |
| 1931           | Essexville sandy loam, 0 to 2 percent slopes                          | 30  |

|        |   |    |
|--------|---|----|
| 875B   | Estherville-Hawick complex, 2 to 6 percent slopes   | 43 |
| 375    | Forada loam, 0 to 2 percent slopes  | 65 |
| 1942   | Forada mucky loam, depressional, 0 to 1 percent slopes                                      | 15 |
| 792    | Fordum fine sandy loam, 0 to 2 percent slopes, frequently flooded                           | 20 |
| 1378   | Fordum loam, 0 to 2 percent slopes, occasionally flooded                                    | 63 |
| 1946   | Fordum-Winterfield complex, 0 to 2 percent slopes, frequently flooded                       | 20 |
| L324A  | Forestcity, overwash-Forestcity complex, 1 to 4 percent slopes                              | 90 |
| 1186   | Forestcity-Lundlake, depressional, complex, 0 to 3 percent slopes                           | 88 |
| 114    | Glencoe clay loam, depressional, 0 to 1 percent slopes                                      | 86 |
| 1099   | Granby loamy fine sand, very wet, 0 to 1 percent slopes                                     | 15 |
| 414    | Hamel loam, 1 to 3 percent slopes   | 94 |
| 740    | Hamel-Glencoe, depressional, complex, 0 to 3 percent slopes                                 | 93 |
| 603    | Hanlon fine sandy loam, 0 to 2 percent slopes, occasionally flooded                         | 82 |
| 611D   | Hawick gravelly sandy loam, 12 to 25 percent slopes   | 31 |
| 875C   | Hawick-Estherville complex, 6 to 12 percent slopes  | 39 |
| L334A  | Houghton and Muskego soils, lundlake catena, 0 to 1 percent slopes                          | 15 |
| 523    | Houghton muck, depressional, 0 to 1 percent slopes  | 74 |
| L351A  | Houghton muck, lundlake catena, 0 to 1 percent slopes                                       | 74 |
| 7A     | Hubbard loamy sand, 0 to 2 percent slopes   | 38 |
| 7B     | Hubbard loamy sand, 2 to 6 percent slopes   | 36 |
| 7C     | Hubbard loamy sand, 6 to 12 percent slopes  | 34 |
| D67A   | Hubbard loamy sand, Mississippi River Valley, 0 to 2 percent slopes                         | 38 |
| D67B   | Hubbard loamy sand, Mississippi River Valley, 2 to 6 percent slopes                         | 36 |
| D67C   | Hubbard loamy sand, Mississippi River Valley, 6 to 12 percent slopes                        | 34 |
| 1231   | Hubbard-Mosford complex, 0 to 3 percent slopes  | 42 |
| D62A   | Hubbard-Mosford complex, Mississippi River Valley, 0 to 3 percent slopes                    | 42 |
| 1224   | Hubbard-Verndale, acid substratum, complex, 0 to 3 percent slopes                           | 42 |
| D61A   | Hubbard-Verndale, acid substratum, complex, Mississippi River Valley, 0 to 3 percent slopes | 42 |
| 1110   | Isan sandy loam, 0 to 2 percent slopes  | 36 |
| 261    | Isan sandy loam, depressional, 0 to 1 percent slopes  | 15 |
| 896B   | Kingsley-Gotham complex, 2 to 6 percent slopes  | 73 |
| 896C   | Kingsley-Gotham complex, 6 to 12 percent slopes   | 60 |
| L347A  | Klossner and Lundlake soils, 0 to 1 percent slopes, ponded                                  | 5  |
| 1199   | Klossner and Lundlake soils, ponded, 0 to 1 percent slopes                                  | 5  |
| 1075   | Klossner and Muskego soils, ponded, 0 to 1 percent slopes                                   | 5  |
| 539    | Klossner muck, depressional, 0 to 1 percent slopes  | 77 |
| L331A  | Klossner muck, lundlake catena, 0 to 1 percent slopes                                       | 77 |
| L335A  | Klossner soils, lundlake catena, 0 to 1 percent slopes                                      | 15 |
| 1080   | Klossner, Okoboji, and Glencoe soils, ponded, 0 to 1 percent slopes                         | 5  |
| 461E   | Koronis loam, 18 to 40 percent slopes   | 29 |
| 461B   | Koronis loam, 2 to 6 percent slopes   | 88 |
| L307B  | Koronis loam, 2 to 6 percent slopes   | 88 |
| 461C2  | Koronis loam, 6 to 12 percent slopes, eroded  | 77 |
| L307C2 | Koronis loam, 6 to 12 percent slopes, moderatley eroded                                     | 77 |
| 807D2  | Koronis-Sunburg complex, 12 to 18 percent slopes, eroded                                    | 61 |
| 804D2  | Koronis-Sunburg-Hawick complex, 12 to 18 percent slopes, eroded                             | 52 |
| 804E   | Koronis-Sunburg-Hawick complex, 18 to 40 percent slopes                                     | 23 |

|        |  |    |
|--------|--|----|
| 804B   | Koronis-Sunburg-Hawick complex, 2 to 6 percent slopes                                  | 76 |
| 804C2  | Koronis-Sunburg-Hawick complex, 6 to 12 percent slopes, eroded                         | 66 |
| L357D2 | Koronis-Sundburg complex, 12 to 18 percent slopes, moderately eroded                   | 61 |
| L355D2 | Koronis-Sundburg-Hawick complex, 12 to 18 percent slopes, moderately eroded            | 52 |
| L355E  | Koronis-Sundburg-Hawick complex, 18 to 40 percent slopes                               | 23 |
| L355B  | Koronis-Sundburg-Hawick complex, 2 to 6 percent slopes                                 | 76 |
| L355C2 | Koronis-Sundburg-Hawick complex, 6 to 12 percent slopes, moderately eroded             | 66 |
| 239    | Le Sueur clay loam, 1 to 3 percent slopes  | 98 |
| 138    | Lerdal silty clay loam, 1 to 3 percent slopes  | 80 |
| 231D   | Lester fine sandy loam, 12 to 18 percent slopes  | 67 |
| 231C   | Lester fine sandy loam, 6 to 12 percent slopes   | 76 |
| 106D2  | Lester loam, 12 to 18 percent slopes, eroded   | 67 |
| 106E   | Lester loam, 18 to 25 percent slopes   | 34 |
| 106C2  | Lester loam, 6 to 12 percent slopes, eroded  | 82 |
| 783C2  | Lester-Kilkenny complex, 6 to 12 percent slopes, eroded                                | 78 |
| 1023D  | Lester-Malardi complex, 12 to 18 percent slopes, eroded                                | 61 |
| 1023C  | Lester-Malardi complex, 6 to 12 percent slopes, eroded                                 | 70 |
| 945D2  | Lester-Storden complex, 12 to 18 percent slopes, eroded                                | 68 |
| 945C2  | Lester-Storden complex, 6 to 12 percent slopes, eroded                                 | 79 |
| 1196B  | Lida-Two Inlets complex, 1 to 8 percent slopes   | 45 |
| 1196E  | Lida-Two Inlets complex, 15 to 30 percent slopes                                       | 14 |
| 1196C  | Lida-Two Inlets complex, 8 to 15 percent slopes  | 38 |
| 1381   | Lindaas silt loam, morainic, 0 to 2 percent slopes                                     | 85 |
| 247    | Linder loam, 0 to 2 percent slopes   | 60 |
| 181    | Litchfield loamy fine sand, 0 to 2 percent slopes                                      | 67 |
| 1026B  | Lizzie silt loam, moderately wet, 1 to 5 percent slopes                                | 95 |
| L318A  | Lundlake silty clay loam, 0 to 1 percent slopes  | 86 |
| 1165   | Lundlake silty clay loam, depressional, 0 to 1 percent slopes                          | 86 |
| 1066B  | Malardi-Hawick complex, 1 to 6 percent slopes  | 39 |
| 1066E  | Malardi-Hawick complex, 18 to 35 percent slopes  | 22 |
| 1066C  | Malardi-Hawick complex, 6 to 12 percent slopes   | 37 |
| L350A  | Marcellon loam, 0 to 3 percent slopes  | 90 |
| 511    | Marcellon loam, map <30, 0 to 3 percent slopes   | 90 |
| 543    | Markey muck, depressional, 0 to 1 percent slopes                                       | 15 |
| 255    | Mayer loam, 0 to 2 percent slopes  | 72 |
| 256    | Mazaska silty clay loam, 0 to 2 percent slopes   | 81 |
| 548    | Medo muck, depressional, 0 to 1 percent slopes   | 84 |
| 1406   | Medo, Dassel, and Biscay soils, ponded, 0 to 1 percent slopes                          | 5  |
| 768    | Mosford sandy loam, 0 to 2 percent slopes  | 50 |
| 1173   | Muskego and Klossner soils, 0 to 1 percent slopes, frequently flooded                  | 20 |
| L320A  | Muskego and Klossner soils, lundlake catena, 0 to 1 percent slopes, frequently flooded | 20 |
| 525    | Muskego muck, depressional, 0 to 1 percent slopes                                      | 15 |
| L330A  | Muskego, Blue Earth and Houghton soils, lundlake catena, 0 to 1 percent slopes, ponded | 5  |
| 1203   | Muskego, Blue Earth, and Houghton soils, ponded, 0 to 1 percent slopes                 | 5  |
| 235    | Nessel loam, 1 to 3 percent slopes   | 92 |
| 1975   | Oylen sandy loam, 0 to 2 percent slopes  | 50 |
| 1030   | Pits, gravel-Udipsammments complex   | 0  |

|       |  |    |
|-------|--|----|
| 294A  | Rasset sandy loam, 0 to 2 percent slopes                                   | 61 |
| 1204B | Reedslake loam, 2 to 5 percent slopes                                      | 98 |
| 1207B | Reedslake-Le Sueur complex, 1 to 5 percent slopes                          | 99 |
| 708   | Rushlake coarse sand, 1 to 4 percent slopes                                | 39 |
| 258E  | Sandberg loamy sand, 12 to 35 percent slopes                               | 17 |
| 258B  | Sandberg loamy sand, 2 to 6 percent slopes                                 | 32 |
| 258C  | Sandberg loamy sand, 6 to 12 percent slopes                                | 30 |
| 1223  | Sandberg-Arvilla complex, map >25, 0 to 3 percent slopes                   | 40 |
| 799   | Seelyeville and Bowstring soils, 0 to 1 percent slopes, frequently flooded | 5  |
| 1288  | Seelyeville and Markey soils, ponded, 0 to 1 percent slopes                | 5  |
| 540   | Seelyeville muck, depressional, 0 to 1 percent slopes                      | 15 |
| 323   | Shields silty clay loam, 0 to 3 percent slopes                             | 84 |
| 286B  | Shorewood silty clay loam, 3 to 6 percent slopes                           | 95 |
| 1383A | Shorewood silty clay loam, moderately wet, 0 to 3 percent slopes           | 95 |
| 1368  | Southhaven loam, 0 to 2 percent slopes                                     | 95 |
| 8B    | Sparta loamy sand, 1 to 6 percent slopes                                   | 39 |
| 1260E | Stonelake-Nebish complex, 12 to 25 percent slopes                          | 17 |
| 1260C | Stonelake-Nebish complex, 6 to 12 percent slopes                           | 38 |
| 1260B | Stonelake-Nebish, moderately wet, complex, 2 to 6 percent slopes           | 47 |
| 1197  | Suckercreek fine sandy loam, 0 to 2 percent slopes, occasionally flooded   | 64 |
| 1163  | Suckercreek loam, 0 to 2 percent slopes, frequently flooded                | 20 |
| 1388B | Terril loam, moderately wet, 2 to 6 percent slopes                         | 99 |
| 1015  | Udipsammments (cut and fill land)  | 0  |
| 1016  | Udorthents, loamy (cut and fill land)                                      | 0  |
| 1072  | Udorthents, shallow (sanitary landfill)                                    | 0  |
| 1027  | Udorthents, wet substratum (fill land)                                     | 0  |
| 1038A | Verndale sandy loam, acid substratum, 0 to 2 percent slopes                | 49 |
| 1038B | Verndale sandy loam, acid substratum, 2 to 6 percent slopes                | 49 |
| 229   | Waldorf silty clay loam, 0 to 2 percent slopes                             | 85 |
| W     | Water  | 0  |
| 1356  | Water, miscellaneous   | 0  |
| 158A  | Zimmerman fine sand, 0 to 3 percent slopes                                 | 39 |
| 158E  | Zimmerman fine sand, 12 to 25 percent slopes                               | 15 |
| 158B  | Zimmerman fine sand, 3 to 6 percent slopes                                 | 39 |
| 158C  | Zimmerman fine sand, 6 to 12 percent slopes                                | 37 |