

Selected Soil Interpretations

This report allows the customer to produce a report showing the results of the soil interpretation(s) of his or her choice. It is useful when a standard report that displays the results of the selected interpretation(s) is not available.

When customers select this report, they are presented with a list of interpretations with results for the selected map units. The customer may select up to three interpretations to be presented in table format.

For a description of the particular interpretations and their criteria, use the "Selected Survey Area Interpretation Descriptions" report.

Report—Selected Soil Interpretations

| Selected Soil Interpretations—Lake County, Ohio | | | |
|--|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| Ad—Adrian muck | | | |
| Adrian | 90 | D-1 | |
| | | D-1 | 0.30 |
| As—Allis silt loam | | | |
| Allis | 95 | C-2 | |
| | | C-2 | 0.12 |
| Bs—Beaches | | | |
| Beaches | 100 | Not rated | |
| Cg—Carlisle muck | | | |
| Carlisle | 95 | D-1 | |
| | | D-1 | 0.30 |
| CoB—Colonie loamy fine sand, 2 to 6 percent slopes | | | |
| Colonie | 95 | B-1 | |
| | | B-1 | 0.07 |
| CoD—Colonie loamy fine sand, 6 to 18 percent slopes | | | |
| Colonie | 95 | B-1 | |
| | | B-1 | 0.07 |
| CoF—Colonie loamy fine sand, 25 to 50 percent slopes | | | |
| Colonie | 85 | B-2 | |
| | | B-2 | 0.08 |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|---|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| CtA—Conneaut silt loam, 0 to 1 percent slopes | | | |
| Conneaut | 95 | C-1 | |
| | | C-1 | 0.11 |
| CtB—Conneaut silt loam, 1 to 4 percent slopes | | | |
| Conneaut | 90 | C-1 | |
| | | C-1 | 0.11 |
| CwA—Conneaut silt loam, shale substratum, 0 to 2 percent slopes | | | |
| Conneaut | 100 | C-1 | |
| | | C-1 | 0.11 |
| CxA—Conotton loam, 0 to 2 percent slopes | | | |
| Conotton | 90 | B-1 | |
| | | B-1 | 0.07 |
| CyB—Conotton gravelly loam, 2 to 6 percent slopes | | | |
| Conotton | 90 | B-1 | |
| | | B-1 | 0.07 |
| CyC—Conotton gravelly loam, 6 to 15 percent slopes | | | |
| Conotton | 95 | B-1 | |
| | | B-1 | 0.07 |
| DaA—Darrien silt loam, 0 to 1 percent slopes | | | |
| Darrien | 95 | C-1 | |
| | | C-1 | 0.11 |
| DaB—Darrien silt loam, 1 to 4 percent slopes | | | |
| Darrien | 90 | C-1 | |
| | | C-1 | 0.11 |
| DaC—Darrien silt loam, 6 to 12 percent slopes | | | |
| Darrien | 90 | C-1 | |
| | | C-1 | 0.11 |
| Dc—Dumps, covered | | | |
| Dumps | 100 | Not rated | |
| DhB—Darrien-Hornell silt loams, 2 to 6 percent slopes | | | |
| Darrien | 48 | C-1 | |
| | | C-1 | 0.11 |
| Hornell | 42 | C-2 | |
| | | C-2 | 0.12 |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|--|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| Du—Dumps, chemical waste | | | |
| Dumps | 100 | Not rated | |
| EIB—Ellsworth silt loam, 2 to 6 percent slopes | | | |
| Ellsworth | 85 | A-1 | |
| | | A-1 | 0.01 |
| EIC—Ellsworth silt loam, 6 to 12 percent slopes | | | |
| Ellsworth | 90 | A-1 | |
| | | A-1 | 0.01 |
| EID—Ellsworth silt loam, 12 to 18 percent slopes | | | |
| Ellsworth | 90 | A-1 | |
| | | A-1 | 0.01 |
| EIF—Ellsworth silt loam, 25 to 70 percent slopes | | | |
| Ellsworth | 85 | H-1 | |
| | | H-1 | 0.71 |
| EmC—Ellsworth silt loam, shale substratum, 6 to 12 percent slopes | | | |
| Ellsworth, shale substratum | 85 | B-1 | |
| | | B-1 | 0.07 |
| EmD—Ellsworth silt loam, shale substratum, 12 to 18 percent slopes | | | |
| Ellsworth, shale substratum | 85 | B-1 | |
| | | B-1 | 0.07 |
| EnB—Elnora loamy fine sand, 1 to 5 percent slopes | | | |
| Elnora | 90 | B-1 | |
| | | B-1 | 0.07 |
| EuA—Euclid silt loam, 0 to 2 percent slopes | | | |
| Euclid | 90 | C-1 | |
| | | C-1 | 0.11 |
| FcA—Fitchville silt loam, 1 to 4 percent slopes | | | |
| Fitchville | 85 | C-1 | |
| | | C-1 | 0.11 |
| GfA—Glenford silt loam, 0 to 2 percent slopes | | | |
| Glenford | 90 | A-6 | |
| | | A-6 | 0.06 |
| GfB—Glenford silt loam, 2 to 6 percent slopes | | | |
| Glenford | 85 | A-6 | |
| | | A-6 | 0.06 |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|---|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| GoF—Gosport silty clay loam, 25 to 70 percent slopes | | | |
| Gosport | 90 | H-1 | |
| | | H-1 | 0.71 |
| Gr—Granby sandy loam | | | |
| Granby | 90 | C-1 | |
| | | C-1 | 0.11 |
| HaA—Harbor fine sandy loam, 0 to 3 percent slopes | | | |
| Harbor | 87 | A-1 | |
| | | A-1 | 0.01 |
| Ho—Holly silt loam, frequently flooded | | | |
| Holly | 85 | C-3 | |
| | | C-3 | 0.13 |
| HrA—Hornell silt loam, 0 to 2 percent slopes | | | |
| Hornell | 95 | C-2 | |
| | | C-2 | 0.12 |
| HrB—Hornell silt loam, 2 to 6 percent slopes | | | |
| Hornell | 90 | C-2 | |
| | | C-2 | 0.12 |
| Kf—Kingsville fine sand | | | |
| Kingsville | 95 | C-1 | |
| | | C-1 | 0.11 |
| Lb—Lobdell silt loam | | | |
| Lobdell | 85 | A-5 | |
| | | A-5 | 0.05 |
| LrB—Lordstown channery silt loam, 2 to 6 percent slopes | | | |
| Lordstown | 85 | F-1 | |
| | | F-1 | 0.51 |
| LrC—Lordstown channery silt loam, 6 to 12 percent slopes | | | |
| Lordstown | 85 | F-1 | |
| | | F-1 | 0.51 |
| LxF—Lordstown-Rock outcrop complex, 25 to 70 percent slopes | | | |
| Lordstown | 50 | H-1 | |
| | | H-1 | 0.71 |
| Rock outcrop | 30 | Not rated | |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|---|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| MgA—Mahoning silt loam, 0 to 2 percent slopes | | | |
| Mahoning | 85 | C-2 | |
| | | C-2 | 0.12 |
| MgB—Mahoning silt loam, 2 to 6 percent slopes | | | |
| Mahoning | 85 | C-2 | |
| | | C-2 | 0.12 |
| MhB—Mahoning silt loam, shale substratum, 2 to 6 percent slopes | | | |
| Mahoning, shale substratum | 85 | C-2 | |
| | | C-2 | 0.12 |
| Mk—Otego silt loam, frequently flooded | | | |
| Otego | 95 | A-5 | |
| | | A-5 | 0.05 |
| Mm—Mill silt loam | | | |
| Mill | 86 | C-1 | |
| | | C-1 | 0.11 |
| Mo—Minoa fine sandy loam | | | |
| Minoa | 90 | C-1 | |
| | | C-1 | 0.11 |
| MtA—Mitiwanga silt loam, 0 to 2 percent slopes | | | |
| Mitiwanga | 90 | C-1 | |
| | | C-1 | 0.11 |
| Or—Orrville silt loam | | | |
| Orrville | 85 | C-3 | |
| | | C-3 | 0.13 |
| OsA—Oshtemo sandy loam, 0 to 2 percent slopes | | | |
| Oshtemo | 95 | B-1 | |
| | | B-1 | 0.07 |
| OsB—Oshtemo sandy loam, 2 to 6 percent slopes | | | |
| Oshtemo | 90 | B-1 | |
| | | B-1 | 0.07 |
| OtB—Otisville gravelly loamy sand, 1 to 6 percent slopes | | | |
| Otisville | 95 | B-1 | |
| | | B-1 | 0.07 |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|---|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| Pa—Painesville fine sandy loam | | | |
| Painesville | 90 | C-1 | |
| | | C-1 | 0.11 |
| PeB—Pierpont silt loam, 2 to 6 percent slopes | | | |
| Pierpont | 85 | F-3 | |
| | | F-3 | 0.53 |
| PeB2—Pierpont silt loam, 2 to 6 percent slopes, moderately eroded | | | |
| Pierpont | 90 | F-3 | |
| | | F-3 | 0.53 |
| PeC2—Pierpont silt loam, 6 to 12 percent slopes, moderately eroded | | | |
| Pierpont | 85 | F-3 | |
| | | F-3 | 0.53 |
| PeD2—Pierpont silt loam, 12 to 18 percent slopes, moderately eroded | | | |
| Pierpont | 85 | F-3 | |
| | | F-3 | 0.53 |
| Po—Pits, gravel | | | |
| Pits | 100 | Not rated | |
| PsA—Platea silt loam, 0 to 2 percent slopes | | | |
| Platea | 95 | C-2 | |
| | | C-2 | 0.12 |
| PsB—Platea silt loam, 2 to 6 percent slopes | | | |
| Platea | 90 | C-2 | |
| | | C-2 | 0.12 |
| PtB—Platea-Darien silt loams, 2 to 6 percent slopes | | | |
| Platea | 50 | C-2 | |
| | | C-2 | 0.12 |
| Darien | 39 | C-1 | |
| | | C-1 | 0.11 |
| RhA—Red Hook sandy loam, 0 to 2 percent slopes | | | |
| Red hook | 85 | C-1 | |
| | | C-1 | 0.11 |
| RkB—Red Hook silt loam, 2 to 6 percent slopes | | | |
| Red hook | 85 | C-1 | |
| | | C-1 | 0.11 |

| Selected Soil Interpretations—Lake County, Ohio | | | |
|---|------------------|------------------------------------|-------|
| Map symbol and soil name | Pct. of map unit | Grl - pasture and hayland sg (oh) | |
| | | Rating class and limiting features | Value |
| Rv—Riverwash | | | |
| Riverwash | 100 | Not rated | |
| St—Stafford loamy fine sand | | | |
| Stafford | 90 | C-1 | |
| | | C-1 | 0.11 |
| Sw—Swanton fine sandy loam | | | |
| Swanton | 95 | C-2 | |
| | | C-2 | 0.12 |
| Tg—Tioga loam | | | |
| Tioga | 90 | A-5 | |
| | | A-5 | 0.05 |
| Th—Tioga Variant silt loam | | | |
| Tioga variant | 90 | A-1 | |
| | | A-1 | 0.01 |
| ToD—Towerville silt loam, 12 to 18 percent slopes | | | |
| Towerville | 100 | F-1 | |
| | | F-1 | 0.51 |
| TyB—Tyner loamy sand, 1 to 6 percent slopes | | | |
| Tyner | 85 | B-1 | |
| | | B-1 | 0.07 |
| TyC—Tyner loamy sand, 6 to 12 percent slopes | | | |
| Tyner | 95 | B-1 | |
| | | B-1 | 0.07 |
| TzA—Tyner Variant sandy loam | | | |
| Tyner variant | 90 | B-1 | |
| | | B-1 | 0.07 |
| UdB—Udorthents, gently sloping | | | |
| Udorthents | 95 | Not rated | |
| UdD—Udorthents, moderately steep | | | |
| Udorthents | 100 | Not rated | |
| Ur—Urban land | | | |
| Urban land | 100 | Not rated | |
| W—Water | | | |
| Water | 100 | Not rated | |

Data Source Information

Soil Survey Area: Lake County, Ohio
Survey Area Data: Version 12, Sep 19, 2014