

Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description (Brief, Generated)

Charlotte County, Florida

Map Unit: 2—Canaveral fine sand

Component: Canaveral (95%)

The Canaveral component makes up 95 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 18 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Captiva (3%)

Generated brief soil descriptions are created for major components. The Captiva soil is a minor component.

Component: Kesson, tidal (2%)

Generated brief soil descriptions are created for major components. The Kesson soil is a minor component.

Map Unit: 4—Canaveral-Urban land complex

Component: Canaveral (60%)

The Canaveral component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 18 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Urban land (30%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Component: Captiva (10%)

Generated brief soil descriptions are created for major components. The Captiva soil is a minor component.

Map Unit: 5—Captiva fine sand

Component: Captiva (92%)

The Captiva component makes up 92 percent of the map unit. Slopes are 0 to 1 percent. This component is on drainageways on marine terraces. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 10 percent. The soil has a slightly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 15 within 30 inches of the soil surface.

Component: Canaveral (4%)

Generated brief soil descriptions are created for major components. The Canaveral soil is a minor component.

Component: Kesson, tidal (4%)

Generated brief soil descriptions are created for major components. The Kesson soil is a minor component.

Map Unit: 6—Hallandale fine sand

Component: Hallandale (90%)

The Hallandale component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 7 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (5%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Rock outcrop (5%)

Generated brief soil descriptions are created for major components. The Rock outcrop soil is a minor component.

Map Unit: 7—Matlacha-Urban land complex

Component: Matlacha (55%)

The Matlacha component makes up 55 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats, coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 30 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Urban land (45%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Map Unit: 8—Hallandale fine sand, tidal

Component: Hallandale, tidal (90%)

The Hallandale, tidal component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 7 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 50 within 30 inches of the soil surface.

Component: Rock Outcrop (10%)

Generated brief soil descriptions are created for major components. The Rock Outcrop soil is a minor component.

Map Unit: 9—EauGallie sand, 0 to 2 percent slopes

Component: EauGallie (85%)

The EauGallie component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wabasso (6%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Delray (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Pinellas (2%)

Generated brief soil descriptions are created for major components. The Pinellas soil is a minor component.

Map Unit: 10—Pompano fine sand, 0 to 2 percent slopes

Component: Pompano (85%)

The Pompano component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Anclote (5%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Malabar (3%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Basinger (2%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Map Unit: 11—Myakka fine sand, 0 to 2 percent slopes

Component: Myakka (90%)

The Myakka component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods, coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: EauGallie, non-hydric (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Placid, depressional (1%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 12—Felda fine sand, 0 to 2 percent slopes

Component: Felda (90%)

The Felda component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on coastal plains, marine terraces on coastal plains, flatwoods on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pinellas (4%)

Generated brief soil descriptions are created for major components. The Pinellas soil is a minor component.

Component: Wabasso (2%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Oldsmar (2%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Map Unit: 13—Boca fine sand, 0 to 2 percent slopes

Component: Boca (85%)

The Boca component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits over marl derived from limestone. Depth to a root restrictive layer, bedrock, lithic, is 8 to 40 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 7 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hallandale (7%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Wabasso (6%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Ft. Drum (2%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Map Unit: 14—Valkaria fine sand, 0 to 2 percent slopes

Component: Valkaria (85%)

The Valkaria component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Satellite (2%)

Generated brief soil descriptions are created for major components. The Satellite soil is a minor component.

Map Unit: 15—Estero muck

Component: Estero, tidal (96%)

The Estero, tidal component makes up 96 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 46 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: Hallandale, tidal (4%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 16—Peckish mucky fine sand

Component: Peckish, tidal (88%)

The Peckish, tidal component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 10 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 65 within 30 inches of the soil surface.

Component: Hallandale, tidal (4%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Boca, tidal (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Estero, tidal (4%)

Generated brief soil descriptions are created for major components. The Estero soil is a minor component.

Map Unit: 17—Daytona sand

Component: Daytona (92%)

The Daytona component makes up 92 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 51 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Immokalee (2%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Pompano (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Orsino (2%)

Generated brief soil descriptions are created for major components. The Orsino soil is a minor component.

Map Unit: 18—Matlacha gravelly fine sand, limestone substratum

Component: Matlacha (83%)

The Matlacha component makes up 83 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats, coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer, bedrock, lithic, is 40 to 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 30 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (9%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (8%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 19—Gator muck

Component: Gator (85%)

The Gator component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 70 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Terra Ceia (15%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Map Unit: 20—Terra Ceia muck

Component: Terra Ceia, drained (85%)

The Terra Ceia, drained component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on marshes on marine terraces on coastal plains. The parent material consists of herbaceous organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator, drained (15%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Map Unit: 22—Beaches

Component: Beaches (95%)

Generated brief soil descriptions are created for major soil components. The Beaches is a miscellaneous area.

Component: Canaveral (5%)

Generated brief soil descriptions are created for major components. The Canaveral soil is a minor component.

Map Unit: 23—Wulfert muck

Component: Wulfert, tidal (90%)

The Wulfert, tidal component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: Kesson, tidal (10%)

Generated brief soil descriptions are created for major components. The Kesson soil is a minor component.

Map Unit: 24—Kesson fine sand

Component: Kesson, tidal (88%)

The Kesson, tidal component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy marine deposits with shells. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 10 percent. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 20 within 30 inches of the soil surface.

Component: Captiva (6%)

Generated brief soil descriptions are created for major components. The Captiva soil is a minor component.

Component: Wulfert, tidal (6%)

Generated brief soil descriptions are created for major components. The Wulfert soil is a minor component.

Map Unit: 25—St. Augustine, organic substratum-Urban land complex

Component: St. Augustine (55%)

The St. Augustine component makes up 55 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Urban land (35%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Component: Kesson, tidal (10%)

Generated brief soil descriptions are created for major components. The Kesson soil is a minor component.

Map Unit: 26—Pineda fine sand, 0 to 2 percent slopes

Component: Pineda (93%)

The Pineda component makes up 93 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 27—Pompano fine sand, depressional

Component: Pompano (92%)

The Pompano component makes up 92 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Anclote (2%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Valkaria (2%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Malabar (2%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Map Unit: 28—Immokalee sand, 0 to 2 percent slopes

Component: Immokalee (90%)

The Immokalee component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Valkaria (2%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Map Unit: 29—Punta fine sand

Component: Punta (94%)

The Punta component makes up 94 percent of the map unit. Slopes are 1 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Immokalee (2%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Smyrna (2%)

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

Map Unit: 33—Oldsmar sand, 0 to 2 percent slopes

Component: Oldsmar (85%)

The Oldsmar component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods, coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Immokalee (7%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Basinger (4%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 34—Malabar fine sand, 0 to 2 percent slopes

Component: Malabar (85%)

The Malabar component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 35—Wabasso sand, 0 to 2 percent slopes

Component: Wabasso (89%)

The Wabasso component makes up 89 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer, strongly contrasting textural stratification, is 9 to 50 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hallandale (6%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Boca (5%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 36—Immokalee-Urban land complex

Component: Immokalee (60%)

The Immokalee component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Urban land (30%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Component: Myakka (4%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Smyrna (3%)

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

Map Unit: 37—Satellite fine sand, 0 to 2 percent slopes

Component: Satellite (85%)

The Satellite component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 18 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (6%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Immokalee (5%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Basinger (3%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 38—Isles fine sand, slough

Component: Isles (83%)

The Isles component makes up 83 percent of the map unit. Slopes are 0 to 1 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 40 to 72 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. The soil has a moderately saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Oldsmar (3%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Wabasso (3%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Map Unit: 39—Isles fine sand, depressional

Component: Isles (80%)

The Isles component makes up 80 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 40 to 72 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. The soil has a moderately saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: Malabar (5%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Felda (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Pineda (5%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Pompano (5%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 40—Anclote sand, depressional

Component: Anclote (88%)

The Anclote component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana (6%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Pompano (6%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 41—Valkaria fine sand, depressional

Component: Valkaria (93%)

The Valkaria component makes up 93 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Anclote (3%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Malabar (2%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pompano (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 42—Wabasso sand, limestone substratum

Component: Wabasso, limestone substratum (85%)

The Wabasso, limestone substratum component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 40 to 80 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Oldsmar, limestone substratum (4%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Myakka (4%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Wabasso (3%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 43—Smyrna fine sand, 0 to 2 percent slopes

Component: Smyrna (85%)

The Smyrna component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (6%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Oldsmar (3%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 44—Malabar fine sand, depressional, 0 to 1 percent slopes

Component: Malabar, depressional (85%)

The Malabar, depressional component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 45—Copeland sandy loam, depressional

Component: Copeland (88%)

The Copeland component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer, bedrock, lithic, is 20 to 50 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Pompano (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Anclote (2%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Boca (2%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Chobee (2%)

Generated brief soil descriptions are created for major components. The Chobee soil is a minor component.

Map Unit: 48—St. Augustine sand

Component: St. Augustine (95%)

The St. Augustine component makes up 95 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: St. Augustine, organic substratum (5%)

Generated brief soil descriptions are created for major components. The St. Augustine soil is a minor component.

Map Unit: 49—Felda fine sand, depressional

Component: Felda (86%)

The Felda component makes up 86 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Winder (2%)

Generated brief soil descriptions are created for major components. The Winder soil is a minor component.

Component: Pineda (2%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Floridana (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Malabar (2%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pompano (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Anclote (2%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Boca (2%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 50—Oldsmar fine sand, limestone substratum

Component: Oldsmar, limestone substratum (88%)

The Oldsmar, limestone substratum component makes up 88 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 60 to 72 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Oldsmar (4%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Wabasso (4%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Immokalee (4%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Map Unit: 51—Floridana sand, depressional

Component: Floridana (88%)

The Floridana component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 8 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (4%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Anclote (4%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Winder (4%)

Generated brief soil descriptions are created for major components. The Winder soil is a minor component.

Map Unit: 53—Myakka fine sand, depressional

Component: Myakka (90%)

The Myakka component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Anclote (2%)

Generated brief soil descriptions are created for major components. The Anclote soil is a minor component.

Component: Immokalee (2%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Wabasso (1%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Oldsmar (1%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Valkaria (1%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 55—Cocoa fine sand

Component: Cocoa (85%)

The Cocoa component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (8%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (7%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 56—Isles muck

Component: Isles, tidal (85%)

The Isles, tidal component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 42 to 72 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 35 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: Wulfert, tidal (5%)

Generated brief soil descriptions are created for major components. The Wulfert soil is a minor component.

Component: Kesson, tidal (5%)

Generated brief soil descriptions are created for major components. The Kesson soil is a minor component.

Component: Boca, tidal (5%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 57—Boca fine sand, tidal

Component: Boca, tidal (85%)

The Boca, tidal component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 24 to 40 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is very frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. The soil has a moderately saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 11 within 30 inches of the soil surface.

Component: Isles, tidal (5%)

Generated brief soil descriptions are created for major components. The Isles soil is a minor component.

Component: Hallandale, tidal (5%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 59—Urban land

Component: Urban land (90%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Component: Matlacha (2%)

Generated brief soil descriptions are created for major components. The Matlacha soil is a minor component.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Immokalee (2%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Smyrna (2%)

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

Component: Hallandale (1%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Boca (1%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 61—Orsino fine sand

Component: Orsino (90%)

The Orsino component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 54 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Daytona (4%)

Generated brief soil descriptions are created for major components. The Daytona soil is a minor component.

Component: Electra (3%)

Generated brief soil descriptions are created for major components. The Electra soil is a minor component.

Component: Satellite (3%)

Generated brief soil descriptions are created for major components. The Satellite soil is a minor component.

Map Unit: 62—Winder sand, depressional

Component: Winder (85%)

The Winder component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Copeland (4%)

Generated brief soil descriptions are created for major components. The Copeland soil is a minor component.

Component: Felda (4%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Rock outcrop (1%)

Generated brief soil descriptions are created for major components. The Rock outcrop soil is a minor component.

Map Unit: 63—Malabar fine sand, high, 0 to 2 percent slopes

Component: Malabar, high (85%)

The Malabar, high component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 66—Caloosa fine sand

Component: Caloosa (85%)

The Caloosa component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on dredge spoil banks on marine terraces on coastal plains. The parent material consists of sandy and clayey dredge spoils. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is moderate. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 36 inches during July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Matlacha (8%)

Generated brief soil descriptions are created for major components. The Matlacha soil is a minor component.

Component: St. Augustine (7%)

Generated brief soil descriptions are created for major components. The St. Augustine soil is a minor component.

Map Unit: 67—Smyrna-Urban land complex

Component: Smyrna (60%)

The Smyrna component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Urban land (35%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

Component: EauGallie (2%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Myakka (1%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Immokalee (1%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Map Unit: 69—Matlacha gravelly fine sand

Component: Matlacha (100%)

The Matlacha component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats, coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 30 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 70—Heights fine sand

Component: Heights (87%)

The Heights component makes up 87 percent of the map unit. Slopes are 0 to 1 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (7%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Wabasso (6%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 72—Bradenton fine sand, 0 to 2 percent slopes

Component: Bradenton (85%)

The Bradenton component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (6%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Parkwood (3%)

Generated brief soil descriptions are created for major components. The Parkwood soil is a minor component.

Component: Copeland (1%)

Generated brief soil descriptions are created for major components. The Copeland soil is a minor component.

Map Unit: 73—Pineda fine sand, depressional, 0 to 1 percent slopes

Component: Pineda, depressional (93%)

The Pineda, depressional component makes up 93 percent of the map unit. Slopes are 0 to 1 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 74—Boca fine sand, slough

Component: Boca (85%)

The Boca component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 24 to 40 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Felda (3%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Pompano (2%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Valkaria (2%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Wabasso (2%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 75—Hallandale fine sand, slough

Component: Hallandale (88%)

The Hallandale component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 2 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Pompano (4%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Map Unit: 76—Electra fine sand

Component: Electra (83%)

The Electra component makes up 83 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during July, August, September, October. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (5%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Immokalee (4%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Bradenton (4%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Daytona (4%)

Generated brief soil descriptions are created for major components. The Daytona soil is a minor component.

Map Unit: 77—Pineda fine sand, limestone substratum

Component: Pineda (88%)

The Pineda component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 40 to 80 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Wabasso (4%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Hallandale (4%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 78—Chobee muck

Component: Chobee (90%)

The Chobee component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 35 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Copeland (3%)

Generated brief soil descriptions are created for major components. The Copeland soil is a minor component.

Component: Floridana (3%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Winder (2%)

Generated brief soil descriptions are created for major components. The Winder soil is a minor component.

Component: Gator (2%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Map Unit: 99—Water

Component: Water (100%)

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

Map Unit: 100—Waters of the Gulf of Mexico

Component: Water of the Gulf of Mexico (100%)

Generated brief soil descriptions are created for major soil components. The Water of the Gulf of Mexico is a miscellaneous area.

Data Source Information

Soil Survey Area: Charlotte County, Florida
Survey Area Data: Version 11, Sep 9, 2014