

Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description (Brief, Generated)

Glades County, Florida

Map Unit: 2—Hallandale fine sand

Component: Hallandale (85%)

The Hallandale component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 7 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Malabar (3%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Boca (3%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Ft. Drum (3%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Component: Pople (3%)

Generated brief soil descriptions are created for major components. The Pople soil is a minor component.

Map Unit: 4—Valkaria fine sand, 0 to 2 percent slopes

Component: Valkaria (85%)

The Valkaria component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Satellite (2%)

Generated brief soil descriptions are created for major components. The Satellite soil is a minor component.

Map Unit: 5—Smyrna fine sand, 0 to 2 percent slopes

Component: Smyrna (85%)

The Smyrna component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (6%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Oldsmar (3%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 6—Malabar fine sand, 0 to 2 percent slopes

Component: Malabar (85%)

The Malabar component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 7—Pople fine sand

Component: Pople (85%)

The Pople component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 15 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Ft. Drum (4%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Valkaria (3%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Map Unit: 8—Gator muck, depressional

Component: Gator (85%)

The Gator component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 70 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (3%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Tequesta, drained (3%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Component: Terra Ceia, drained (3%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Component: Floridana, depressional (3%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Chobee (3%)

Generated brief soil descriptions are created for major components. The Chobee soil is a minor component.

Map Unit: 9—Sanibel muck, depressional

Component: Sanibel (88%)

The Sanibel component makes up 88 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of thin organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 35 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Dania, drained (3%)

Generated brief soil descriptions are created for major components. The Dania soil is a minor component.

Component: Lauderhill, drained (3%)

Generated brief soil descriptions are created for major components. The Lauderhill soil is a minor component.

Component: Plantation, drained (2%)

Generated brief soil descriptions are created for major components. The Plantation soil is a minor component.

Component: Pahokee, drained (2%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Component: Terra Ceia, drained (2%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Map Unit: 10—Felda fine sand, 0 to 2 percent slopes

Component: Felda (90%)

The Felda component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on coastal plains, marine terraces on coastal plains, flatwoods on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pinellas (4%)

Generated brief soil descriptions are created for major components. The Pinellas soil is a minor component.

Component: Wabasso (2%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Oldsmar (2%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Myakka (2%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Map Unit: 11—Tequesta muck, drained

Component: Tequesta, drained (86%)

The Tequesta, drained component makes up 86 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of stratified sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 55 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 1 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger, depressional (4%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Floridana, depressional (4%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Gator (3%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Component: Sanibel (3%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Map Unit: 12—Chobee loamy fine sand, depressional

Component: Chobee (86%)

The Chobee component makes up 86 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 15 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, depressional (3%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Component: Felda (3%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Tequesta, drained (2%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Component: Sanibel (2%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Component: Floridana, depressional (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Gator (2%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Map Unit: 13—Boca fine sand, 0 to 2 percent slopes

Component: Boca (85%)

The Boca component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits over marl derived from limestone. Depth to a root restrictive layer, bedrock, lithic, is 8 to 40 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 7 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hallandale (7%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Component: Wabasso (6%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Ft. Drum (2%)

Generated brief soil descriptions are created for major components. The Ft. Drum soil is a minor component.

Map Unit: 14—Basinger fine sand, 0 to 2 percent slopes

Component: Basinger (90%)

The Basinger component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Placid, depressional (3%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Component: Margate (3%)

Generated brief soil descriptions are created for major components. The Margate soil is a minor component.

Map Unit: 15—Pineda fine sand, 0 to 2 percent slopes

Component: Pineda (93%)

The Pineda component makes up 93 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 16—Floridana fine sand, depressional

Component: Floridana, depressional (85%)

The Floridana, depressional component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 8 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator (3%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Component: Felda (3%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Tequesta, drained (3%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Component: Sanibel (3%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Component: Astor, depressional (3%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Map Unit: 17—Okeelanta muck, 0 to 1 percent slopes

Component: Okeelanta (90%)

The Okeelanta component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, depressional (3%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Component: Sanibel (3%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Component: Tequesta (2%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Component: Floridana, depressional (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Map Unit: 19—Terra Ceia muck, drained

Component: Terra Ceia, drained (88%)

The Terra Ceia, drained component makes up 88 percent of the map unit. Slopes are 0 to 1 percent. This component is on marshes on marine terraces on coastal plains. The parent material consists of herbaceous organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelanta (4%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Pahokee, drained (4%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Component: Lauderhill, drained (4%)

Generated brief soil descriptions are created for major components. The Lauderhill soil is a minor component.

Map Unit: 20—EauGallie fine sand, 0 to 2 percent slopes

Component: EauGallie (85%)

The EauGallie component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wabasso (6%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Delray (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Pinellas (2%)

Generated brief soil descriptions are created for major components. The Pinellas soil is a minor component.

Map Unit: 22—Astor fine sand, depressional

Component: Astor, depressional (87%)

The Astor, depressional component makes up 87 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger, depressional (2%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Floridana, depressional (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Gator (2%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Okeelanta (2%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Sanibel (2%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Component: Tequesta, drained (1%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Map Unit: 23—Oldsmar sand, 0 to 2 percent slopes

Component: Oldsmar (85%)

The Oldsmar component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods, coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Immokalee (7%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Basinger (4%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 24—Hallandale-Pople complex

Component: Hallandale (45%)

The Hallandale component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 7 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pople (35%)

The Pople component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 15 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (10%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Malabar (10%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Map Unit: 26—Immokalee sand, 0 to 2 percent slopes

Component: Immokalee (90%)

The Immokalee component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Felda (2%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Valkaria (2%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Map Unit: 27—Ft. Drum fine sand

Component: Ft. Drum (85%)

The Ft. Drum component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 20 within 30 inches of the soil surface.

Component: Pople (4%)

Generated brief soil descriptions are created for major components. The Pople soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pineda (4%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Valkaria (3%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Map Unit: 28—Pomello fine sand, 0 to 2 percent slopes

Component: Pomello (85%)

The Pomello component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY001FL Sand Pine Scrub ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Immokalee (5%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Duette (5%)

Generated brief soil descriptions are created for major components. The Duette soil is a minor component.

Component: Jonathan (3%)

Generated brief soil descriptions are created for major components. The Jonathan soil is a minor component.

Component: Tavares (2%)

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

Map Unit: 29—Myakka fine sand, 0 to 2 percent slopes

Component: Myakka (90%)

The Myakka component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods, coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: EauGallie, non-hydric (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Placid, depressional (1%)

Generated brief soil descriptions are created for major components. The Placid soil is a minor component.

Map Unit: 32—Floridana, Astor, and Felda soils, frequently flooded

Component: Floridana, flooded (32%)

The Floridana, flooded component makes up 32 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 11 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, flooded (31%)

The Astor, flooded component makes up 31 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda, flooded (31%)

The Felda, flooded component makes up 31 percent of the map unit. Slopes are 0 to 1 percent. This component is on swamps on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (2%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Okeelanta (1%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Chobee (1%)

Generated brief soil descriptions are created for major components. The Chobee soil is a minor component.

Component: Gator (1%)

Generated brief soil descriptions are created for major components. The Gator soil is a minor component.

Component: Terra Ceia, drained (1%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Map Unit: 34—Basinger fine sand, depressional, 0 to 1 percent slopes

Component: Basinger, depressional (90%)

The Basinger, depressional component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Smyrna, hydric (5%)

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

Component: Samsula, muck (3%)

Generated brief soil descriptions are created for major components. The Samsula soil is a minor component.

Component: Floridana, hydric (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Map Unit: 35—Arents, very steep

Component: Arents (100%)

The Arents component makes up 100 percent of the map unit. Slopes are 45 to 60 percent. This component is on fills, rises on marine terraces on coastal plains. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 36—Malabar fine sand, high, 0 to 2 percent slopes

Component: Malabar, high (85%)

The Malabar, high component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY003FL South Florida Flatwoods ecological site. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 37—Lauderhill muck, drained

Component: Lauderhill, drained (92%)

The Lauderhill, drained component makes up 92 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over limestone. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, June, July, August, September, October, November. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Terra Ceia, drained (2%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Component: Plantation, drained (2%)

Generated brief soil descriptions are created for major components. The Plantation soil is a minor component.

Component: Dania, drained (2%)

Generated brief soil descriptions are created for major components. The Dania soil is a minor component.

Component: Pahokee, drained (2%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Map Unit: 38—Pahokee muck, drained, 0 to 1 percent slopes

Component: Pahokee, drained (90%)

The Pahokee, drained component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on marine terraces on coastal plains. The parent material consists of herbaceous organic material over limestone. Depth to a root restrictive layer, bedrock, lithic, is 36 to 51 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 83 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca, depressional (6%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Dania, drained (2%)

Generated brief soil descriptions are created for major components. The Dania soil is a minor component.

Component: Lauderhill, drained (2%)

Generated brief soil descriptions are created for major components. The Lauderhill soil is a minor component.

Map Unit: 40—Plantation muck, drained

Component: Plantation, drained (94%)

The Plantation, drained component makes up 94 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, June, July, August, September, October, November. Organic matter content in the surface horizon is about 45 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lauderhill, drained (2%)

Generated brief soil descriptions are created for major components. The Lauderhill soil is a minor component.

Component: Dania, drained (2%)

Generated brief soil descriptions are created for major components. The Dania soil is a minor component.

Component: Sanibel, drained (1%)

Generated brief soil descriptions are created for major components. The Sanibel soil is a minor component.

Component: Pahokee, drained (1%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Map Unit: 41—Dania muck, drained

Component: Dania, drained (94%)

The Dania, drained component makes up 94 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over limestone. Depth to a root restrictive layer, bedrock, lithic, is 8 to 20 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, June, July, August, September, October, November. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Plantation, drained (2%)

Generated brief soil descriptions are created for major components. The Plantation soil is a minor component.

Component: Lauderhill, drained (2%)

Generated brief soil descriptions are created for major components. The Lauderhill soil is a minor component.

Component: Pahokee, drained (2%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Map Unit: 42—Okeelanta and Dania soils, depressional

Component: Okeelanta (55%)

The Okeelanta component makes up 55 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 65 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Dania (35%)

The Dania component makes up 35 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over limestone. Depth to a root restrictive layer, bedrock, lithic, is 8 to 20 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, depressional (2%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Component: Floridana, depressional (2%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Pahokee, drained (2%)

Generated brief soil descriptions are created for major components. The Pahokee soil is a minor component.

Component: Lauderhill, drained (2%)

The Lauderhill, drained component makes up 2 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over limestone. Depth to a root restrictive layer, bedrock, lithic, is 20 to 40 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, June, July, August, September, October, November. Organic matter content in the surface horizon is about 75 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Terra Ceia, drained (1%)

Generated brief soil descriptions are created for major components. The Terra Ceia soil is a minor component.

Component: Tequesta, drained (1%)

Generated brief soil descriptions are created for major components. The Tequesta soil is a minor component.

Map Unit: 43—Sanibel muck, drained

Component: Sanibel, drained (88%)

The Sanibel, drained component makes up 88 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of thin organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, June, July, August, September, October, November. Organic matter content in the surface horizon is about 55 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelanta (4%)

Generated brief soil descriptions are created for major components. The Okeelanta soil is a minor component.

Component: Astor, depressional (4%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Component: Floridana, depressional (4%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Map Unit: 99—Water

Component: Water (100%)

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

Data Source Information

Soil Survey Area: Glades County, Florida
Survey Area Data: Version 11, Sep 9, 2014