

## Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

## Report—Map Unit Description (Brief, Generated)

### Hamilton County, Florida

**Map Unit:** 2—Albany fine sand, 0 to 5 percent slopes

**Component:** Albany (90%)

The Albany component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 21 inches during January, February, March, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Plummer, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Component:** Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit:** 3—Alpin sand, 0 to 5 percent slopes

**Component:** Alpin (90%)

The Alpin component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of eolian deposits or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Albany (4%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Component:** Blanton (3%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component:** Chipley (3%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

**Map Unit:** 4—Alpin sand, 5 to 8 percent slopes

**Component:** Alpin (90%)

The Alpin component makes up 90 percent of the map unit. Slopes are 5 to 8 percent. This component is on marine terraces on coastal plains, hillslopes. The parent material consists of eolian deposits or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (5%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component: Albany (5%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit: 5—Blanton sand, 0 to 5 percent slopes**

**Component: Blanton (90%)**

The Blanton component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Albany (10%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit: 6—Blanton sand, 5 to 8 percent slopes**

**Component: Blanton (85%)**

The Blanton component makes up 85 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Albany (8%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Component: Alpin (7%)**

Generated brief soil descriptions are created for major components. The Alpin soil is a minor component.

**Map Unit: 7—Kenansville fine sand, 0 to 5 percent slopes, occasionally flooded**

**Component: Kenansville (90%)**

The Kenansville component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine or fluvial deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Ocilla (5%)**

Generated brief soil descriptions are created for major components. The Ocilla soil is a minor component.

**Component: Blanton (5%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit: 8—Chipley sand, 0 to 5 percent slopes**

**Component:** Chipley (90%)

The Chipley component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on flats on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 30 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Mascotte, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component:** Pottsburg, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Map Unit:** 9—Foxworth sand, 0 to 5 percent slopes

**Component:** Foxworth (95%)

The Foxworth component makes up 95 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges, coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, March, April, May, June, July, August, September, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R133AY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Lakeland (4%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

**Component:** Chipley (1%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

**Map Unit: 10**—Lowndes sand, 0 to 5 percent slopes

**Component: Lowndes (90%)**

The Lowndes component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Norfolk (10%)**

Generated brief soil descriptions are created for major components. The Norfolk soil is a minor component.

**Map Unit: 11**—Lowndes sand, 5 to 8 percent slopes

**Component: Lowndes (90%)**

The Lowndes component makes up 90 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Valdosta (10%)**

Generated brief soil descriptions are created for major components. The Valdosta soil is a minor component.

**Map Unit: 12**—Lowndes and Norfolk soils, 8 to 12 percent slopes

**Component: Lowndes (40%)**

The Lowndes component makes up 40 percent of the map unit. Slopes are 8 to 12 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Norfolk (30%)**

The Norfolk component makes up 30 percent of the map unit. Slopes are 8 to 10 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Wampee (15%)**

Generated brief soil descriptions are created for major components. The Wampee soil is a minor component.

**Component: Valdosta (15%)**

Generated brief soil descriptions are created for major components. The Valdosta soil is a minor component.

**Map Unit: 13—Mascotte sand**

**Component: Mascotte, non-hydric (70%)**

The Mascotte, non-hydric component makes up 70 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on stream terraces on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 15 inches during March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Mascotte, hydric (20%)

The Mascotte, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on stream terraces on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Sapelo, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Sapelo soil is a minor component.

**Component:** Pottsburg, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Map Unit:** 14—Pottsburg sand

**Component:** Pottsburg, non-hydric (70%)

The Pottsburg, non-hydric component makes up 70 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Pottsburg, hydric (20%)

The Pottsburg, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Albany (4%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Component:** Mascotte, non-hydric (3%)

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component:** Plummer, depressional (3%)

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Map Unit:** 15—Valdosta sand, 0 to 5 percent slopes

**Component:** Valdosta (90%)

The Valdosta component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component:** Lowndes (5%)

Generated brief soil descriptions are created for major components. The Lowndes soil is a minor component.

**Map Unit:** 16—Valdosta sand, 5 to 8 percent slopes

**Component:** Valdosta (90%)

The Valdosta component makes up 90 percent of the map unit. Slopes are 5 to 8 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Lowndes (10%)

Generated brief soil descriptions are created for major components. The Lowndes soil is a minor component.

**Map Unit:** 17—Wadley sand, 5 to 12 percent slopes

**Component:** Wadley (87%)

The Wadley component makes up 87 percent of the map unit. Slopes are 5 to 12 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Foxworth (13%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

**Map Unit:** 18—Wadley sand, 0 to 5 percent slopes

**Component:** Wadley (90%)

The Wadley component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Alpin (5%)

Generated brief soil descriptions are created for major components. The Alpin soil is a minor component.

**Component:** Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit:** 19—Valdosta-Lowndes complex, 12 to 20 percent slopes

**Component:** Valdosta (67%)

The Valdosta component makes up 67 percent of the map unit. Slopes are 12 to 20 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Lowndes (28%)**

The Lowndes component makes up 28 percent of the map unit. Slopes are 12 to 20 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (3%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component: Wampee (2%)**

Generated brief soil descriptions are created for major components. The Wampee soil is a minor component.

**Map Unit: 20—Pamlico muck, depressional**

**Component: Pamlico (90%)**

The Pamlico component makes up 90 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Pelham, hydric (3%)

Generated brief soil descriptions are created for major components. The Pelham soil is a minor component.

**Component:** Mascotte, non-hydric (3%)

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component:** Plummer, depressional (2%)

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Component:** Pottsburg, hydric (2%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Map Unit:** 21—Plummer and Surrency soils, depressional

**Component:** Plummer, depressional (50%)

The Plummer, depressional component makes up 50 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Surrency, depressional (33%)

The Surrency, depressional component makes up 33 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 15 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Mascotte, hydric (10%)

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component:** Pottsburg, non-hydric (7%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Map Unit:** 22—Alpin fine sand, 0 to 5 percent slopes, occasionally flooded

**Component:** Alpin (95%)

The Alpin component makes up 95 percent of the map unit. Slopes are 0 to 5 percent. This component is on stream terraces on marine terraces on coastal plains. The parent material consists of eolian deposits or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit:** 23—Blanton loamy sand, 0 to 5 percent slopes

**Component:** Blanton (90%)

The Blanton component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Valdosta (5%)**

Generated brief soil descriptions are created for major components. The Valdosta soil is a minor component.

**Component: Kenansville (5%)**

Generated brief soil descriptions are created for major components. The Kenansville soil is a minor component.

**Map Unit: 24—Ocilla loamy fine sand, 0 to 5 percent slopes**

**Component: Ocilla (90%)**

The Ocilla component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 21 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (5%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component: Pelham, non-hydric (5%)**

Generated brief soil descriptions are created for major components. The Pelham soil is a minor component.

**Map Unit: 25—Wampee-Blanton complex, 8 to 12 percent slopes**

**Component: Wampee (54%)**

The Wampee component makes up 54 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (36%)**

The Blanton component makes up 36 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Albany (4%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Component: Mascotte, non-hydric (3%)**

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component: Plummer, non-hydric (3%)**

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Map Unit: 26—Mascotte and Plummer soils, occasionally flooded**

**Component: Mascotte (53%)**

The Mascotte component makes up 53 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Plummer (36%)

The Plummer component makes up 36 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Stockade (11%)

Generated brief soil descriptions are created for major components. The Stockade soil is a minor component.

**Map Unit:** 27—Kenansville loamy sand, 0 to 5 percent slopes

**Component:** Kenansville (90%)

The Kenansville component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine or fluvial deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Norfolk (10%)

Generated brief soil descriptions are created for major components. The Norfolk soil is a minor component.

**Map Unit: 28—Wampee loamy sand, 5 to 8 percent slopes**

**Component: Wampee (87%)**

The Wampee component makes up 87 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Albany (7%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Component: Blanton (6%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit: 29—Bonneau sand, 0 to 5 percent slopes**

**Component: Bonneau (90%)**

The Bonneau component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 51 inches during January, February, March, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Norfolk (10%)**

Generated brief soil descriptions are created for major components. The Norfolk soil is a minor component.

**Map Unit: 31—Wampee-Blanton complex, 12 to 20 percent slopes**

**Component: Wampee (50%)**

The Wampee component makes up 50 percent of the map unit. Slopes are 12 to 15 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (37%)**

The Blanton component makes up 37 percent of the map unit. Slopes are 12 to 20 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Albany (13%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit: 32—Norfolk loamy fine sand, 2 to 5 percent slopes**

**Component: Norfolk (90%)**

The Norfolk component makes up 90 percent of the map unit. Slopes are 2 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Lowndes (4%)**

Generated brief soil descriptions are created for major components. The Lowndes soil is a minor component.

**Component: Ocilla (3%)**

Generated brief soil descriptions are created for major components. The Ocilla soil is a minor component.

**Component: Valdosta (3%)**

Generated brief soil descriptions are created for major components. The Valdosta soil is a minor component.

**Map Unit: 33—Pelham sand**

**Component: Pelham, non-hydric (60%)**

The Pelham, non-hydric component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, April. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Pelham, hydric (30%)**

The Pelham, hydric component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Albany (10%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit:** 34—Plummer sand

**Component:** Plummer, non-hydric (60%)

The Plummer, non-hydric component makes up 60 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, April, May, June, July, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Plummer, hydric (30%)

The Plummer, hydric component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Sapelo, non-hydric (10%)

Generated brief soil descriptions are created for major components. The Sapelo soil is a minor component.

**Map Unit:** 35—Wahee fine sandy loam, 0 to 4 percent slopes, occasionally flooded

**Component:** Wahee (90%)

The Wahee component makes up 90 percent of the map unit. Slopes are 0 to 4 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is moderate. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Ocilla (5%)

Generated brief soil descriptions are created for major components. The Ocilla soil is a minor component.

**Component:** Eunola (5%)

Generated brief soil descriptions are created for major components. The Eunola soil is a minor component.

**Map Unit:** 36—Blanton fine sand, 0 to 5 percent slopes, occasionally flooded

**Component:** Blanton (90%)

The Blanton component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on stream terraces on marine terraces, coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 66 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Kenansville (5%)

Generated brief soil descriptions are created for major components. The Kenansville soil is a minor component.

**Component: Alpin (5%)**

Generated brief soil descriptions are created for major components. The Alpin soil is a minor component.

**Map Unit: 37—Eunola loamy fine sand, 0 to 5 percent slopes, occasionally flooded**

**Component: Eunola (90%)**

The Eunola component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during January, February, March, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (4%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component: Ocilla (3%)**

Generated brief soil descriptions are created for major components. The Ocilla soil is a minor component.

**Component: Wahee (3%)**

Generated brief soil descriptions are created for major components. The Wahee soil is a minor component.

**Map Unit: 46—Stockade fine sandy loam**

**Component: Stockade (90%)**

The Stockade component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Surrency, depressional (5%)

Generated brief soil descriptions are created for major components. The Surrency soil is a minor component.

**Component:** Pelham, hydric (5%)

Generated brief soil descriptions are created for major components. The Pelham soil is a minor component.

**Map Unit:** 47—Goldhead fine sand, 0 to 5 percent slopes

**Component:** Goldhead, non-hydric (65%)

The Goldhead, non-hydric component makes up 65 percent of the map unit. Slopes are 0 to 5 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Goldhead, hydric (25%)

The Goldhead, hydric component makes up 25 percent of the map unit. Slopes are 0 to 5 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Wampee (5%)**

Generated brief soil descriptions are created for major components. The Wampee soil is a minor component.

**Component: Albany (5%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit: 48—Bivans loamy sand, 8 to 12 percent slopes**

**Component: Bivans (85%)**

The Bivans component makes up 85 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes, marine terraces on coastal plains. The parent material consists of sandy and clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is very low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is high. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 15 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Wampee (5%)**

Generated brief soil descriptions are created for major components. The Wampee soil is a minor component.

**Component: Pelham, non-hydric (5%)**

Generated brief soil descriptions are created for major components. The Pelham soil is a minor component.

**Component: Plummer, non-hydric (5%)**

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Map Unit: 49—Otela-Alpin complex, 0 to 5 percent slopes**

**Component: Otela (59%)**

The Otela component makes up 59 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Alpin (28%)**

The Alpin component makes up 28 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of eolian deposits or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Ocilla (13%)**

Generated brief soil descriptions are created for major components. The Ocilla soil is a minor component.

**Map Unit: 51—Bigbee fine sand, undulating, occasionally flooded**

**Component: Bigbee (90%)**

The Bigbee component makes up 90 percent of the map unit. Slopes are 0 to 5 percent. This component is on stream terraces on marine terraces on coastal plains. The parent material consists of sandy fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Blanton (10%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit:** 52—Pelham fine sand, occasionally flooded

**Component:** Pelham, non-hydric (70%)

The Pelham, non-hydric component makes up 70 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, April. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Pelham, hydric (20%)

The Pelham, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Stockade (5%)

Generated brief soil descriptions are created for major components. The Stockade soil is a minor component.

**Component: Albany (5%)**

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit: 54—Pits**

**Component: Pits (100%)**

Generated brief soil descriptions are created for major soil components. The Pits is a miscellaneous area.

**Map Unit: 56—Bibb-Bigbee complex, undulating, occasionally flooded**

**Component: Bibb (50%)**

The Bibb component makes up 50 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy and sandy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Bigbee (40%)**

The Bigbee component makes up 40 percent of the map unit. Slopes are 0 to 5 percent. This component is on stream terraces on marine terraces on coastal plains. The parent material consists of sandy fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Eunola (5%)**

Generated brief soil descriptions are created for major components. The Eunola soil is a minor component.

**Component: Blanton (5%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Map Unit: 57—Osier sand, occasionally flooded**

**Component: Osier (90%)**

The Osier component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Pottsburg, non-hydric (5%)**

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Component: Plummer (5%)**

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Map Unit: 58—Sapelo sand**

**Component: Sapelo, non-hydric (75%)**

The Sapelo, non-hydric component makes up 75 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, April, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Sapelo, hydric (15%)**

The Sapelo, hydric component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Albany (10%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

**Map Unit:** 59—Dorovan muck, depressional

**Component:** Dorovan (90%)

The Dorovan component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Pelham, hydric (3%)

Generated brief soil descriptions are created for major components. The Pelham soil is a minor component.

**Component:** Mascotte, hydric (3%)

Generated brief soil descriptions are created for major components. The Mascotte soil is a minor component.

**Component:** Plummer, depressional (2%)

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

**Component:** Pottsburg, hydric (2%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

**Map Unit: 60—Alpin-Shadeville complex, karst**

**Component: Alpin (50%)**

The Alpin component makes up 50 percent of the map unit. Slopes are 0 to 8 percent. This component is on ridges, karstic marine terraces on coastal plains. The parent material consists of eolian deposits or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Shadeville (40%)**

The Shadeville component makes up 40 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on karstic marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 40 to 72 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 60 inches during July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Bivans (4%)**

Generated brief soil descriptions are created for major components. The Bivans soil is a minor component.

**Component: Blanton (3%)**

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

**Component: Wampee (3%)**

Generated brief soil descriptions are created for major components. The Wampee soil is a minor component.

**Map Unit: 61—Arents, 0 to 5 percent slopes**

**Component: Arents (95%)**

The Arents component makes up 95 percent of the map unit. Slopes are 0 to 5 percent. This component is on fills on marine terraces on coastal plains. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Pits (5%)**

Generated brief soil descriptions are created for major components. The Pits soil is a minor component.

**Map Unit: 62—Resota-Blanton-Bigbee complex, undulating, occasionally flooded**

**Component: Resota (35%)**

The Resota component makes up 35 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 51 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Blanton (33%)**

The Blanton component makes up 33 percent of the map unit. Slopes are 0 to 8 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 66 inches during March, April, May, June, July, August. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Bigbee (25%)**

The Bigbee component makes up 25 percent of the map unit. Slopes are 0 to 5 percent. This component is on stream terraces on marine terraces on coastal plains, flood plains on marine terraces on coastal plains. The parent material consists of sandy fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Otela (7%)**

Generated brief soil descriptions are created for major components. The Otela soil is a minor component.

**Map Unit: 63—Arents-Water complex**

**Component: Arents (55%)**

The Arents component makes up 55 percent of the map unit. Slopes are 2 to 5 percent. This component is on fills on marine terraces on coastal plains. The parent material consists of altered marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Water (45%)**

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

**Map Unit: 64—Hydraquents, clayey**

**Component: Hydraquents (95%)**

The Hydraquents component makes up 95 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is high. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 7 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component:** Arents (3%)

Generated brief soil descriptions are created for major components. The Arents soil is a minor component.

**Component:** Pits (2%)

Generated brief soil descriptions are created for major components. The Pits soil is a minor component.

**Map Unit:** 65—Gypsum land

**Component:** Gypsum land (100%)

Generated brief soil descriptions are created for major soil components. The Gypsum land is a miscellaneous area.

**Map Unit:** 66—Urban land

**Component:** Urban land (85%)

Generated brief soil descriptions are created for major soil components. The Urban land is a miscellaneous area.

**Component:** Arents (15%)

Generated brief soil descriptions are created for major components. The Arents soil is a minor component.

**Map Unit:** 67—Quartzipsamments, 1 to 5 percent slopes

**Component:** Quartzipsamments (90%)

The Quartzipsamments component makes up 90 percent of the map unit. Slopes are 1 to 5 percent. This component is on fills on rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

**Component: Arents (4%)**

Generated brief soil descriptions are created for major components. The Arents soil is a minor component.

**Component: Pits (3%)**

Generated brief soil descriptions are created for major components. The Pits soil is a minor component.

**Component: Hydraquents (3%)**

Generated brief soil descriptions are created for major components. The Hydraquents soil is a minor component.

**Map Unit: 99—Water**

**Component: Water (100%)**

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

## Data Source Information

Soil Survey Area: Hamilton County, Florida  
Survey Area Data: Version 13, Sep 24, 2014