Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description (Brief, Generated)

Liberty County, Florida

Map Unit: 2—Albany sand, 0 to 5 percent slopes

Component: Albany (85%)

The Albany component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 21 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Plummer, non-hydric (10%)

Generated brief soil descriptions are created for major components. The Plummer soil is a minor component.

Component: Blanton (3%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Component: Leefield (2%)

Generated brief soil descriptions are created for major components. The Leefield soil is a minor component.

Map Unit: 4—Alpin-Foxworth complex, 5 to 12 percent slopes

Component: Alpin (45%)

The Alpin component makes up 45 percent of the map unit. Slopes are 5 to 12 percent. This component is on ridges on marine terraces on coastal plains, hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (40%)

The Foxworth component makes up 40 percent of the map unit. Slopes are 5 to 8 percent. This component is on ridges on marine terraces on coastal plains, hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Centenary (8%)

Generated brief soil descriptions are created for major components. The Centenary soil is a minor component.

Component: Chipley (7%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 5—Rains and Bladen soils

Component: Rains (50%)

The Rains component makes up 50 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on fluviomarine terraces on coastal plains, drainageways on fluviomarine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bladen (45%)

The Bladen component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on fluviomarine terraces on coastal plains, drainageways on fluviomarine terraces on coastal plains. The parent material consists of loamy and clayey alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is moderate. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynchburg (3%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Component: Leefield (2%)

Generated brief soil descriptions are created for major components. The Leefield soil is a minor component.

Map Unit: 6—Blanton sand, 0 to 5 percent slopes

Component: Blanton (85%)

The Blanton component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on hillslopes on marine terraces on coastal plains, ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Fuquay (10%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Albany (5%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Map Unit: 7—Blanton sand, 5 to 8 percent slopes

Component: Blanton (80%)

The Blanton component makes up 80 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Albany (10%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Component: Fuquay (5%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Foxworth (5%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Map Unit: 8—Brickyard clay loam, frequently flooded

Component: Brickyard (75%)

The Brickyard component makes up 75 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of loamy and clayey alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is moderate. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wahee (18%)

Generated brief soil descriptions are created for major components. The Wahee soil is a minor component.

Component: Ochlockonee (3%)

Generated brief soil descriptions are created for major components. The Ochlockonee soil is a minor component.

Component: Chowan (2%)

Generated brief soil descriptions are created for major components. The Chowan soil is a minor component.

Component: Ousley (2%)

Generated brief soil descriptions are created for major components. The Ousley soil is a minor component.

Map Unit: 9—Centenary sand, 0 to 5 percent slopes

Component: Centenary (85%)

The Centenary component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 51 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lakeland (8%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

Component: Alpin (5%)

Generated brief soil descriptions are created for major components. The Alpin soil is a minor component.

Component: Chipley (2%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 11—Chipley-Foxworth complex, 0 to 5 percent slopes

Component: Chipley (50%)

The Chipley component makes up 50 percent of the map unit. Slopes are 0 to 5 percent. This component is on knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (40%)

The Foxworth component makes up 40 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lakeland (5%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

Component: Leon, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Map Unit: 12—Rutlege and Plummer soils, depressional

Component: Rutlege (45%)

The Rutlege component makes up 45 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Plummer (40%)

The Plummer component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bibb (15%)

Generated brief soil descriptions are created for major components. The Bibb soil is a minor component.

Map Unit: 13—Dorovan-Pamlico complex, depressional

Component: Dorovan (50%)

The Dorovan component makes up 50 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pamlico (45%)

The Pamlico component makes up 45 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynn Haven (5%)

Generated brief soil descriptions are created for major components. The Lynn Haven soil is a minor component.

Map Unit: 14—Dothan loamy sand, 0 to 2 percent slopes

Component: Dothan (80%)

The Dothan component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on interfluves on coastal plains. The parent material consists of marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 34 inches during January, February, March. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 1. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface.

Component: Cowarts (5%)

Generated brief soil descriptions are created for major components. The Cowarts soil is a minor component.

Component: Fuquay (5%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Norfolk (5%)

Generated brief soil descriptions are created for major components. The Norfolk soil is a minor component.

Component: Clarendon (5%)

Generated brief soil descriptions are created for major components. The Clarendon soil is a minor component.

Map Unit: 15—Dothan loamy sand, 2 to 5 percent slopes

Component: Dothan (80%)

The Dothan component makes up 80 percent of the map unit. Slopes are 2 to 5 percent. This component is on interfluves on coastal plains. The parent material consists of marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 34 inches during January, February, March. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface.

Component: Nankin (5%)

Generated brief soil descriptions are created for major components. The Nankin soil is a minor component.

Component: Clarendon (5%)

Generated brief soil descriptions are created for major components. The Clarendon soil is a minor component.

Component: Fuquay (5%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Cowarts (5%)

Generated brief soil descriptions are created for major components. The Cowarts soil is a minor component.

Map Unit: 17—Dothan-Fuquay complex, 8 to 12 percent slopes

Component: Dothan (40%)

The Dothan component makes up 40 percent of the map unit. Slopes are 8 to 12 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of loamy and clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 42 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Fuquay (35%)

The Fuquay component makes up 35 percent of the map unit. Slopes are 8 to 10 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Goldsboro (10%)

Generated brief soil descriptions are created for major components. The Goldsboro soil is a minor component.

Component: Blanton (10%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Component: Stilson (5%)

Generated brief soil descriptions are created for major components. The Stilson soil is a minor component.

Map Unit: 24—Goldsboro loamy sand, 0 to 2 percent slopes

Component: Goldsboro (90%)

The Goldsboro component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on rises on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 27 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynchburg (5%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Component: Fuquay (3%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Albany (2%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Map Unit: 25—Goldsboro loamy sand, 2 to 5 percent slopes

Component: Goldsboro (90%)

The Goldsboro component makes up 90 percent of the map unit. Slopes are 2 to 5 percent. This component is on knolls on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 27 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Fuquay (3%)

Generated brief soil descriptions are created for major components. The Fuquay soil is a minor component.

Component: Albany (3%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Component: Dothan (2%)

Generated brief soil descriptions are created for major components. The Dothan soil is a minor component.

Component: Lynchburg (2%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Map Unit: 26—Foxworth sand, 0 to 5 percent slopes

Component: Foxworth (95%)

The Foxworth component makes up 95 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges, coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during January, February, March, April, May, June, July, August, September, December. Organic matter content in the surface horizon is about 1 percent. This component is in the R133AY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lakeland (4%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

Component: Chipley (1%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 27—Fuquay loamy sand, 0 to 5 percent slopes

Component: Fuquay (85%)

The Fuquay component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on sand sheets on marine terraces on coastal plains, hillslopes on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Blanton (8%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Component: Goldsboro (4%)

Generated brief soil descriptions are created for major components. The Goldsboro soil is a minor component.

Component: Stilson (3%)

Generated brief soil descriptions are created for major components. The Stilson soil is a minor component.

Map Unit: 30—Elloree, Bibb, and Meggett soils, 0 to 3 percent slopes, frequently flooded

Component: Elloree (35%)

The Elloree component makes up 35 percent of the map unit. Slopes are 0 to 3 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria.

Component: Bibb (30%)

The Bibb component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Meggett (25%)

The Meggett component makes up 25 percent of the map unit. Slopes are 0 to 3 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of clayey. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is high. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rutlege (10%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Map Unit: 31—Hurricane and Chipley soils, 0 to 3 percent slopes

Component: Hurricane (45%)

The Hurricane component makes up 45 percent of the map unit. Slopes are 0 to 3 percent. This component is on rises on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (35%)

The Chipley component makes up 35 percent of the map unit. Slopes are 0 to 3 percent. This component is on knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon, non-hydric (7%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Component: Centenary (7%)

Generated brief soil descriptions are created for major components. The Centenary soil is a minor component.

Component: Pottsburg, non-hydric (6%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

Map Unit: 32—Plummer and Pelham soils

Component: Plummer (45%)

The Plummer component makes up 45 percent of the map unit. Slopes are 0 to 1 percent. This component is on open depressions on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pelham (40%)

The Pelham component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on open depressions on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pottsburg, non-hydric (5%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

Component: Pottsburg, hydric (5%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

Component: Sapelo, hydric (5%)

Generated brief soil descriptions are created for major components. The Sapelo soil is a minor component.

Map Unit: 34—Lakeland sand, 0 to 5 percent slopes

Component: Lakeland (77%)

The Lakeland component makes up 77 percent of the map unit. Slopes are 0 to 5 percent. This component is on hills on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. Irrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Troup (14%)

Generated brief soil descriptions are created for major components. The Troup soil is a minor component.

Component: Bonifay (9%)

Generated brief soil descriptions are created for major components. The Bonifay soil is a minor component.

Map Unit: 35—Lakeland sand, 5 to 8 percent slopes

Component: Lakeland (85%)

The Lakeland component makes up 85 percent of the map unit. Slopes are 5 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (10%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Component: Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Map Unit: 36—Lakeland sand, 8 to 15 percent slopes

Component: Lakeland (90%)

The Lakeland component makes up 90 percent of the map unit. Slopes are 8 to 15 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (10%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Map Unit: 37—Lakeland-Foxworth complex, 15 to 30 percent slopes

Component: Lakeland (50%)

The Lakeland component makes up 50 percent of the map unit. Slopes are 15 to 30 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (40%)

The Foxworth component makes up 40 percent of the map unit. Slopes are 15 to 25 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 38—Leefield loamy sand, 0 to 5 percent slopes

Component: Leefield (85%)

The Leefield component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Albany (5%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Component: Stilson (5%)

Generated brief soil descriptions are created for major components. The Stilson soil is a minor component.

Component: Dothan (5%)

Generated brief soil descriptions are created for major components. The Dothan soil is a minor component.

Map Unit: 39—Leon sand, 0 to 2 percent slopes

Component: Leon (80%)

The Leon component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on marine terraces, coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 5 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. Irrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon, hydric (5%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Component: Pottsburg (4%)

Generated brief soil descriptions are created for major components. The Pottsburg soil is a minor component.

Component: Hurricane (4%)

Generated brief soil descriptions are created for major components. The Hurricane soil is a minor component.

Component: Mandarin (3%)

Generated brief soil descriptions are created for major components. The Mandarin soil is a minor component.

Component: Rutlege (2%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Component: Pickney (2%)

Generated brief soil descriptions are created for major components. The Pickney soil is a minor component.

Map Unit: 42—Lucy sand, 0 to 5 percent slopes

Component: Lucy (85%)

The Lucy component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on sand sheets on marine terraces on coastal plains, hillslopes on sand sheets on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Stilson (10%)

Generated brief soil descriptions are created for major components. The Stilson soil is a minor component.

Component: Blanton (5%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Map Unit: 44—Lynchburg loamy sand

Component: Lynchburg (80%)

The Lynchburg component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rains (10%)

Generated brief soil descriptions are created for major components. The Rains soil is a minor component.

Component: Goldsboro (10%)

Generated brief soil descriptions are created for major components. The Goldsboro soil is a minor component.

Map Unit: 45—Lynn Haven sand

Component: Lynn Haven (90%)

The Lynn Haven component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon, hydric (5%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Component: Pickney, frequently flooded (3%)

Generated brief soil descriptions are created for major components. The Pickney soil is a minor component.

Component: Pamlico, frequently flooded (2%)

Generated brief soil descriptions are created for major components. The Pamlico soil is a minor component.

Map Unit: 46—Hurricane, Leon, and Albany soils

Component: Hurricane (35%)

The Hurricane component makes up 35 percent of the map unit. Slopes are 0 to 3 percent. This component is on knolls on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon (30%)

The Leon component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on swales on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Albany (25%)

The Albany component makes up 25 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 21 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (5%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Component: Foxworth (5%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Map Unit: 47—Torhunta-Lynn Haven-Croatan complex, frequently flooded

Component: Torhunta (35%)

The Torhunta component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on marine terraces, coastal plains, flood plains. The parent material consists of loamy and sandy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 7 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynn Haven (30%)

The Lynn Haven component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Croatan (25%)

The Croatan component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of herbaceous organic material and/or loamy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is frequently flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, July, August, September, November. Organic matter content in the surface horizon is about 43 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon, non-hydric (10%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Map Unit: 48—Meadowbrook sand

Component: Meadowbrook, non-hydric (75%)

The Meadowbrook, non-hydric component makes up 75 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Meadowbrook, hydric (15%)

The Meadowbrook, hydric component makes up 15 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Goldhead, depressional (8%)

Generated brief soil descriptions are created for major components. The Goldhead soil is a minor component.

Component: Pantego, depressional (2%)

Generated brief soil descriptions are created for major components. The Pantego soil is a minor component.

Map Unit: 49—Meadowbrook sand, slough

Component: Meadowbrook, slough (85%)

The Meadowbrook, slough component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rutlege (8%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Component: Surrency, depressional (7%)

Generated brief soil descriptions are created for major components. The Surrency soil is a minor component.

Map Unit: 54—Pelham loamy sand

Component: Pelham, non-hydric (65%)

The Pelham, non-hydric component makes up 65 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pelham, hydric (20%)

The Pelham, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leefield (10%)

Generated brief soil descriptions are created for major components. The Leefield soil is a minor component.

Component: Lynchburg (5%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Map Unit: 55—Plummer sand, 0 to 5 percent slopes

Component: Plummer, non-hydric (60%)

The Plummer, non-hydric component makes up 60 percent of the map unit. Slopes are 0 to 5 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Plummer, hydric (20%)

The Plummer, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynchburg (10%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Component: Leefield (10%)

Generated brief soil descriptions are created for major components. The Leefield soil is a minor component.

Map Unit: 56—Pottsburg-Pottsburg, wet, sand, 0 to 2 percent slopes

Component: Pottsburg (65%)

The Pottsburg component makes up 65 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4w. Irrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pottsburg, wet (25%)

The Pottsburg, wet component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4w. Irrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Leon, non-hydric (6%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Component: Allanton (2%)

Generated brief soil descriptions are created for major components. The Allanton soil is a minor component.

Component: Chipley (2%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 57—Surrency, Pantego, and Croatan soils, depressional

Component: Surrency, depressional (35%)

The Surrency, depressional component makes up 35 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pantego, depressional (30%)

The Pantego, depressional component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 7 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Croatan, depressional (25%)

The Croatan, depressional component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material and/or loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 43 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Osier (10%)

Generated brief soil descriptions are created for major components. The Osier soil is a minor component.

Map Unit: 58—Rutlege, Bibb, and Surrency soils, frequently flooded

Component: Rutlege (35%)

The Rutlege component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bibb (30%)

The Bibb component makes up 30 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Surrency (25%)

The Surrency component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wahee (10%)

Generated brief soil descriptions are created for major components. The Wahee soil is a minor component.

Map Unit: 59—Hosford mucky coarse sand, 2 to 8 percent slopes

Component: Hosford (80%)

The Hosford component makes up 80 percent of the map unit. Slopes are 2 to 8 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 10 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Component: Hurricane (10%)

Generated brief soil descriptions are created for major components. The Hurricane soil is a minor component.

Map Unit: 60—Sapelo sand

Component: Sapelo, non-hydric (65%)

The Sapelo, non-hydric component makes up 65 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Sapelo, hydric (20%)

The Sapelo, hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Component: Rutlege (5%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Map Unit: 61—Osier sand

Component: Osier (80%)

The Osier component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rutlege (10%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Component: Hosford (10%)

Generated brief soil descriptions are created for major components. The Hosford soil is a minor component.

Map Unit: 62—Scranton loamy sand, slough

Component: Scranton, slough (90%)

The Scranton, slough component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rutlege (10%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Map Unit: 63—Stilson fine sand, 0 to 3 percent slopes

Component: Stilson (85%)

The Stilson component makes up 85 percent of the map unit. Slopes are 0 to 3 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Albany (10%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Component: Dothan (5%)

Generated brief soil descriptions are created for major components. The Dothan soil is a minor component.

Map Unit: 65—Pickney, Dorovan, and Bibb soils, frequently flooded

Component: Pickney (40%)

The Pickney component makes up 40 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 10 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bibb (25%)

The Bibb component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Dorovan (25%)

The Dorovan component makes up 25 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of herbaceous organic material and/or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wahee (10%)

Generated brief soil descriptions are created for major components. The Wahee soil is a minor component.

Map Unit: 66—Wahee and Ochlockonee soils, 0 to 3 percent slopes, occasionally flooded

Component: Wahee (45%)

The Wahee component makes up 45 percent of the map unit. Slopes are 0 to 3 percent. This component is on stream terraces on flood plains on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is moderate. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 24 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ochlockonee (35%)

The Ochlockonee component makes up 35 percent of the map unit. Slopes are 0 to 3 percent. This component is on stream terraces on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chowan (10%)

Generated brief soil descriptions are created for major components. The Chowan soil is a minor component.

Component: Brickyard (10%)

Generated brief soil descriptions are created for major components. The Brickyard soil is a minor component.

Map Unit: 67—Goldhead sand

Component: Goldhead (85%)

The Goldhead component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Surrency, depressional (10%)

Generated brief soil descriptions are created for major components. The Surrency soil is a minor component.

Component: Pantego, depressional (5%)

Generated brief soil descriptions are created for major components. The Pantego soil is a minor component.

Map Unit: 68—Goldhead-Meadowbrook complex, depressional

Component: Goldhead, depressional (45%)

The Goldhead, depressional component makes up 45 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Meadowbrook, depressional (40%)

The Meadowbrook, depressional component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pamlico, frequently flooded (8%)

Generated brief soil descriptions are created for major components. The Pamlico soil is a minor component.

Component: Surrency, depressional (7%)

Generated brief soil descriptions are created for major components. The Surrency soil is a minor component.

Map Unit: 69—Troup sand, 0 to 5 percent slopes

Component: Troup (80%)

The Troup component makes up 80 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. This component is in the R133AY002FL Longleaf Pine-turkey Oak Hills ecological site. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Blanton (10%)

Generated brief soil descriptions are created for major components. The Blanton soil is a minor component.

Component: Lakeland (5%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

Component: Foxworth (5%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Map Unit: 70—Troup sand, 5 to 8 percent slopes

Component: Troup (88%)

The Troup component makes up 88 percent of the map unit. Slopes are 5 to 8 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat excessively drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lucy (5%)

Generated brief soil descriptions are created for major components. The Lucy soil is a minor component.

Component: Bonifay (4%)

Generated brief soil descriptions are created for major components. The Bonifay soil is a minor component.

Component: Lakeland (3%)

Generated brief soil descriptions are created for major components. The Lakeland soil is a minor component.

Map Unit: 71—Pits

Component: Pits (98%)

Generated brief soil descriptions are created for major soil components. The Pits is a miscellaneous area.

Component: Troup (2%)

Generated brief soil descriptions are created for major components. The Troup soil is a minor component.

Map Unit: 72—Lakeland sand, 30 to 85 percent slopes

Component: Lakeland (75%)

The Lakeland component makes up 75 percent of the map unit. Slopes are 30 to 85 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (10%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Component: Gullied land (5%)

Generated brief soil descriptions are created for major components. The Gullied land soil is a minor component.

Map Unit: 73—Foxworth-Hosford-Lucy complex, 8 to 25 percent slopes

Component: Foxworth (45%)

The Foxworth component makes up 45 percent of the map unit. Slopes are 8 to 25 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hosford (25%)

The Hosford component makes up 25 percent of the map unit. Slopes are 3 to 12 percent. This component is on lower hillslopes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 10 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lucy (20%)

The Lucy component makes up 20 percent of the map unit. Slopes are 8 to 25 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 74—Garcon, Ochlockonee, and Ousley soils, occasionally flooded

Component: Garcon (40%)

The Garcon component makes up 40 percent of the map unit. Slopes are 0 to 3 percent. This component is on flood plains on marine terraces on coastal plains, stream terraces on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ousley (25%)

The Ousley component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on stream terraces on flood plains on marine terraces on coastal plains. The parent material consists of sandy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 0 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ochlockonee (25%)

The Ochlockonee component makes up 25 percent of the map unit. Slopes are 0 to 3 percent. This component is on flood plains on marine terraces on coastal plains, stream terraces on marine terraces on coastal plains. The parent material consists of sandy and loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Brickyard (5%)

Generated brief soil descriptions are created for major components. The Brickyard soil is a minor component.

Component: Chowan (5%)

Generated brief soil descriptions are created for major components. The Chowan soil is a minor component.

Map Unit: 75—Brantley-Okeelala-Lucy complex, 8 to 45 percent slopes

Component: Brantley (35%)

The Brantley component makes up 35 percent of the map unit. Slopes are 8 to 35 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of loamy and clayey marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 7e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Okeelala (30%)

The Okeelala component makes up 30 percent of the map unit. Slopes are 8 to 45 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lucy (25%)

The Lucy component makes up 25 percent of the map unit. Slopes are 8 to 25 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Troup (10%)

Generated brief soil descriptions are created for major components. The Troup soil is a minor component.

Map Unit: 78—Lucy-Blanton-Cowarts complex, 8 to 45 percent slopes

Component: Lucy (35%)

The Lucy component makes up 35 percent of the map unit. Slopes are 8 to 45 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Blanton (30%)

The Blanton component makes up 30 percent of the map unit. Slopes are 8 to 45 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Cowarts (25%)

The Cowarts component makes up 25 percent of the map unit. Slopes are 8 to 25 percent. This component is on hillslopes on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 6e. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Troup (10%)

Generated brief soil descriptions are created for major components. The Troup soil is a minor component.

Map Unit: 81—Scranton fine sand, 0 to 2 percent slopes

Component: Scranton (84%)

The Scranton component makes up 84 percent of the map unit. Slopes are 0 to 2 percent. This component is on coastal plains, flatwoods. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, April, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Scranton, slough (10%)

Generated brief soil descriptions are created for major components. The Scranton soil is a minor component.

Component: Rutlege (3%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Component: Leon (3%)

Generated brief soil descriptions are created for major components. The Leon soil is a minor component.

Map Unit: 82—Brickyard and Chowan soils, frequently flooded

Component: Brickyard (55%)

The Brickyard component makes up 55 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy and clayey alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is low. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is moderate. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chowan (35%)

The Chowan component makes up 35 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy marine deposits over organic material. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Wahee (10%)

Generated brief soil descriptions are created for major components. The Wahee soil is a minor component.

Map Unit: 83—Plummer, Sapelo, and Pottsburg soils

Component: Plummer (35%)

The Plummer component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Sapelo (30%)

The Sapelo component makes up 30 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pottsburg (25%)

The Pottsburg component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hurricane (6%)

Generated brief soil descriptions are created for major components. The Hurricane soil is a minor component.

Component: Foxworth (2%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Component: Centenary (2%)

Generated brief soil descriptions are created for major components. The Centenary soil is a minor component.

Map Unit: 91—Woodington loamy sand

Component: Woodington (90%)

The Woodington component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on marine terraces, flats, coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lynchburg (6%)

Generated brief soil descriptions are created for major components. The Lynchburg soil is a minor component.

Component: Leefield (4%)

Generated brief soil descriptions are created for major components. The Leefield soil is a minor component.

Map Unit: 92—Pamlico-Pickney complex, frequently flooded

Component: Pamlico, frequently flooded (50%)

The Pamlico, frequently flooded component makes up 50 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is frequently flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 50 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pickney, frequently flooded (35%)

The Pickney, frequently flooded component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits and/or fluviomarine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September. Organic matter content in the surface horizon is about 10 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Dorovan (8%)

Generated brief soil descriptions are created for major components. The Dorovan soil is a minor component.

Component: Rutlege (7%)

Generated brief soil descriptions are created for major components. The Rutlege soil is a minor component.

Map Unit: 95—Bibb, Rains, and Garcon Soils, occasionally flooded

Component: Bibb (40%)

The Bibb component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 9 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Rains (25%)

The Rains component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Garcon (25%)

The Garcon component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains, stream terraces on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Ochlockonee (10%)

Generated brief soil descriptions are created for major components. The Ochlockonee soil is a minor component.

Map Unit: 96—Foxworth-Lakeland complex, 0 to 5 percent slopes

Component: Foxworth (50%)

The Foxworth component makes up 50 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lakeland (40%)

The Lakeland component makes up 40 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 97—Foxworth-Lakeland complex, 5 to 15 percent slopes

Component: Foxworth (45%)

The Foxworth component makes up 45 percent of the map unit. Slopes are 5 to 10 percent. This component is on hillslopes on marine terraces on coastal plains, ridges on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during March. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Lakeland (40%)

The Lakeland component makes up 40 percent of the map unit. Slopes are 8 to 15 percent. This component is on ridges on marine terraces on coastal plains, hillslopes on marine terraces on coastal plains. The parent material consists of sandy eolian deposits and/or marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is excessively drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (15%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Map Unit: 98—Leon-Chipley complex

Component: Leon, non-hydric (45%)

The Leon, non-hydric component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (30%)

The Chipley component makes up 30 percent of the map unit. Slopes are 0 to 5 percent. This component is on knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 23 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Foxworth (10%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Component: Albany (10%)

Generated brief soil descriptions are created for major components. The Albany soil is a minor component.

Component: Leon, hydric (5%)

The Leon, hydric component makes up 5 percent of the map unit. Slopes are 0 to 2 percent. This component is on knolls on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 99—Water

Component: Water (100%)

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

Map Unit: 101—Albany-Blanton complex, 0 to 5 percent slopes

Component: Albany (55%)

The Albany component makes up 55 percent of the map unit. Slopes are 0 to 5 percent. This component is on rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 21 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Blanton (30%)

The Blanton component makes up 30 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 48 inches during January, February, March, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Chipley (10%)

Generated brief soil descriptions are created for major components. The Chipley soil is a minor component.

Component: Foxworth (5%)

Generated brief soil descriptions are created for major components. The Foxworth soil is a minor component.

Data Source Information

Soil Survey Area: Liberty County, Florida
Survey Area Data: Version 18, Sep 26, 2014