

Map Unit Description (Brief, Generated)

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions in this report, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

The Map Unit Description (Brief, Generated) report displays a generated description of the major soils that occur in a map unit. Descriptions of non-soil (miscellaneous areas) and minor map unit components are not included. This description is generated from the underlying soil attribute data.

Additional information about the map units described in this report is available in other Soil Data Mart reports, which give properties of the soils and the limitations, capabilities, and potentials for many uses. Also, the narratives that accompany the Soil Data Mart reports define some of the properties included in the map unit descriptions.

Report—Map Unit Description (Brief, Generated)

Sarasota County, Florida

Map Unit: 2—Beaches

Component: Beaches (100%)

Generated brief soil descriptions are created for major soil components. The Beaches is a miscellaneous area.

Map Unit: 3—Boca and Hallandale soils

Component: Boca (45%)

The Boca component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 24 to 40 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Hallandale (35%)

The Hallandale component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits over limestone. Depth to a root restrictive layer, bedrock, lithic, is 7 to 20 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pineda (5%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Rock outcrop (5%)

Generated brief soil descriptions are created for major components. The Rock outcrop soil is a minor component.

Component: Pompano (5%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Felda, hydric (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Map Unit: 4—Bradenton fine sand, 0 to 2 percent slopes

Component: Bradenton (85%)

The Bradenton component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 5 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (6%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Parkwood (3%)

Generated brief soil descriptions are created for major components. The Parkwood soil is a minor component.

Component: Copeland (1%)

Generated brief soil descriptions are created for major components. The Copeland soil is a minor component.

Map Unit: 5—Bradenton fine sand, frequently flooded

Component: Bradenton, flooded (85%)

The Bradenton, flooded component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pineda (5%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Component: Felda, flooded (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Astor, flooded (5%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Map Unit: 6—Canaveral fine sand, 0 to 5 percent slopes

Component: Canaveral (85%)

The Canaveral component makes up 85 percent of the map unit. Slopes are 0 to 5 percent. This component is on ridges on marine terraces on coastal plains, dunes on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 18 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: St. Augustine (10%)

Generated brief soil descriptions are created for major components. The St. Augustine soil is a minor component.

Component: Pompano (5%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 7—Cassia fine sand

Component: Cassia (85%)

The Cassia component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, rises on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 22 inches during July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (5%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pomello (5%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

Map Unit: 8—Delray fine sand, depressional

Component: Delray, depressional (80%)

The Delray, depressional component makes up 80 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator, depressional (5%)

The Gator, depressional component makes up 5 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 70 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pompano (5%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Felda (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Astor, flooded (5%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Map Unit: 9—Delray and Astor soils, frequently flooded

Component: Delray, flooded (45%)

The Delray, flooded component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 7 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, flooded (35%)

The Astor, flooded component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 3 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 14 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana, flooded (10%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Felda, flooded (10%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Map Unit: 10—EauGallie and Myakka fine sands

Component: EauGallie (45%)

The EauGallie component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (40%)

The Myakka component makes up 40 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Smyrna (5%)

Generated brief soil descriptions are created for major components. The Smyrna soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Component: Ona (5%)

Generated brief soil descriptions are created for major components. The Ona soil is a minor component.

Map Unit: 11—Felda fine sand

Component: Felda, hydric (65%)

The Felda, hydric component makes up 65 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda, non-hydric (20%)

The Felda, non-hydric component makes up 20 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 10 inches during July, August, September, October, November. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Bradenton (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Holopaw (5%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Map Unit: 12—Felda fine sand, depressional

Component: Felda (85%)

The Felda component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Bradenton (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Holopaw (5%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Component: Floridana, depressional (5%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Map Unit: 13—Felda and Pompano fine sands, frequently flooded

Component: Felda, flooded (45%)

The Felda, flooded component makes up 45 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pompano, flooded (35%)

The Pompano, flooded component makes up 35 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 6w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Astor, flooded (5%)

Generated brief soil descriptions are created for major components. The Astor soil is a minor component.

Component: Bradenton, flooded (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Holopaw (5%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Component: Delray, flooded (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 14—Floridana mucky fine sand

Component: Floridana, drained (85%)

The Floridana, drained component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains, flats on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 13 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda, hydric (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Manatee (5%)

Generated brief soil descriptions are created for major components. The Manatee soil is a minor component.

Component: Delray, depressional (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 15—Floridana and Gator soils, depressional

Component: Floridana, depressional (75%)

The Floridana, depressional component makes up 75 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 13 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator, depressional (25%)

The Gator, depressional component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 70 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 16—Floridana and Gator soils, frequently flooded

Component: Floridana, flooded (75%)

The Floridana, flooded component makes up 75 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 13 percent. Nonirrigated land capability classification is 5w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator, flooded (25%)

The Gator, flooded component makes up 25 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on marine terraces, coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 85 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 17—Gator muck

Component: Gator (45%)

The Gator component makes up 45 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 85 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Gator, drained (40%)

The Gator, drained component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of herbaceous organic material over loamy and sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is very high. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 18 inches during January, February, March, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 85 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Manatee (5%)

Generated brief soil descriptions are created for major components. The Manatee soil is a minor component.

Component: Delray, depressional (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Component: Floridana, depressional (5%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Map Unit: 21—Ft. Green fine sand

Component: Ft. Green (85%)

The Ft. Green component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 5 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Holopaw (4%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Component: EauGallie (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Wabasso (3%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 22—Holopaw fine sand, depressional

Component: Holopaw (85%)

The Holopaw component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Floridana, depressional (4%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Manatee (4%)

Generated brief soil descriptions are created for major components. The Manatee soil is a minor component.

Component: Malabar (4%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Map Unit: 24—Kesson and Wulfert mucks, frequently flooded

Component: Kesson, Tidal (50%)

The Kesson, Tidal component makes up 50 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of sandy marine deposits with shells. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 35 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 10 percent. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 20 within 30 inches of the soil surface.

Component: Wulfert, Tidal (40%)

The Wulfert, Tidal component makes up 40 percent of the map unit. Slopes are 0 to 1 percent. This component is on tidal marshes on marine terraces on coastal plains. The parent material consists of organic material over sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is high. Shrink-swell potential is low. This soil is frequently flooded. It is not ponded. A seasonal zone of water saturation is at 0 inches during January, February, March, April, May, June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 44 percent. Nonirrigated land capability classification is 8. This soil meets hydric criteria. The soil has a strongly saline horizon within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 60 within 30 inches of the soil surface.

Component: St. Augustine (5%)

Generated brief soil descriptions are created for major components. The St. Augustine soil is a minor component.

Component: Beaches (5%)

Generated brief soil descriptions are created for major components. The Beaches soil is a minor component.

Map Unit: 25—Malabar fine sand, 0 to 2 percent slopes

Component: Malabar (85%)

The Malabar component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. This component is in the R155XY011FL Slough ecological site. Nonirrigated land capability classification is 4w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Basinger (6%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Valkaria (5%)

Generated brief soil descriptions are created for major components. The Valkaria soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Component: Delray (1%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 26—Manatee loamy fine sand, depressional

Component: Manatee (85%)

The Manatee component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 6 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Holopaw (3%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Component: Floridana, depressional (3%)

Generated brief soil descriptions are created for major components. The Floridana soil is a minor component.

Component: Felda (3%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Malabar (3%)

Generated brief soil descriptions are created for major components. The Malabar soil is a minor component.

Component: Pineda (3%)

Generated brief soil descriptions are created for major components. The Pineda soil is a minor component.

Map Unit: 27—Matlacha gravelly sand

Component: Matlacha (100%)

The Matlacha component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats, coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 30 inches during June, July, August, September, October. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Map Unit: 29—Orsino fine sand

Component: Orsino (85%)

The Orsino component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is very high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 54 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 4s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (5%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Pomello (5%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

Component: Myakka (5%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Map Unit: 30—Ona fine sand

Component: Ona (85%)

The Ona component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pomello (4%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

Component: EauGallie (4%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Myakka (4%)

Generated brief soil descriptions are created for major components. The Myakka soil is a minor component.

Component: Pompano (3%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 31—Pineda fine sand, 0 to 2 percent slopes

Component: Pineda (93%)

The Pineda component makes up 93 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Boca (4%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Component: Hallandale (3%)

Generated brief soil descriptions are created for major components. The Hallandale soil is a minor component.

Map Unit: 32—Pits and Dumps

Component: Dumps (50%)

Generated brief soil descriptions are created for major soil components. The Dumps is a miscellaneous area.

Component: Pits (50%)

Generated brief soil descriptions are created for major soil components. The Pits is a miscellaneous area.

Map Unit: 33—Pomello fine sand, 0 to 2 percent slopes

Component: Pomello (85%)

The Pomello component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, flatwoods on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 33 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 1 percent. This component is in the R155XY001FL Sand Pine Scrub ecological site. Nonirrigated land capability classification is 6s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Duette (5%)

Generated brief soil descriptions are created for major components. The Duette soil is a minor component.

Component: Immokalee (5%)

Generated brief soil descriptions are created for major components. The Immokalee soil is a minor component.

Component: Jonathan (3%)

Generated brief soil descriptions are created for major components. The Jonathan soil is a minor component.

Component: Tavares (2%)

Generated brief soil descriptions are created for major components. The Tavares soil is a minor component.

Map Unit: 34—Pompano fine sand, depressional

Component: Pompano (85%)

The Pompano component makes up 85 percent of the map unit. Slopes are 0 to 1 percent. This component is on depressions on marine terraces on coastal plains. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is very poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is frequently ponded. A seasonal zone of water saturation is at 0 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 4 percent. Nonirrigated land capability classification is 7w. This soil meets hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (5%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Holopaw (5%)

Generated brief soil descriptions are created for major components. The Holopaw soil is a minor component.

Component: Delray, depressional (5%)

Generated brief soil descriptions are created for major components. The Delray soil is a minor component.

Map Unit: 36—Pople fine sand

Component: Pople (85%)

The Pople component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on drainageways on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 6 inches during June, July, August, September, October, November. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil meets hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 15 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (5%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Bradenton (5%)

Generated brief soil descriptions are created for major components. The Bradenton soil is a minor component.

Component: Wabasso (5%)

Generated brief soil descriptions are created for major components. The Wabasso soil is a minor component.

Map Unit: 38—Smyrna fine sand, 0 to 2 percent slopes

Component: Smyrna (85%)

The Smyrna component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on — Error in Exists On —. The parent material consists of sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches (or restricted depth) is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 3 percent. Nonirrigated land capability classification is 4w. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: EauGallie (6%)

Generated brief soil descriptions are created for major components. The EauGallie soil is a minor component.

Component: Basinger (5%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Oldsmar (3%)

Generated brief soil descriptions are created for major components. The Oldsmar soil is a minor component.

Component: Pompano (1%)

Generated brief soil descriptions are created for major components. The Pompano soil is a minor component.

Map Unit: 39—St. Augustine fine sand

Component: St. Augustine (90%)

The St. Augustine component makes up 90 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains. The parent material consists of sandy mine spoil or earthy fill. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is somewhat poorly drained. Water movement in the most restrictive layer is moderately low. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 22 inches during June, July, August, September. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 7s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Matlacha (10%)

Generated brief soil descriptions are created for major components. The Matlacha soil is a minor component.

Map Unit: 40—Tavares fine sand

Component: Tavares (85%)

The Tavares component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on ridges on marine terraces on coastal plains, knolls on marine terraces on coastal plains. The parent material consists of eolian or sandy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is very low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 57 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3s. This soil does not meet hydric criteria. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Pomello (5%)

Generated brief soil descriptions are created for major components. The Pomello soil is a minor component.

Component: Cassia (5%)

Generated brief soil descriptions are created for major components. The Cassia soil is a minor component.

Component: Orsino (5%)

Generated brief soil descriptions are created for major components. The Orsino soil is a minor component.

Map Unit: 41—Wabasso fine sand, 0 to 2 percent slopes

Component: Wabasso (85%)

The Wabasso component makes up 85 percent of the map unit. Slopes are 0 to 2 percent. This component is on flats on marine terraces on coastal plains, flatwoods on marine terraces on coastal plains. The parent material consists of sandy and loamy marine deposits. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is poorly drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches (or restricted depth) is low. Shrink-swell potential is low. This soil is not flooded. It is not ponded. A seasonal zone of water saturation is at 12 inches during June, July, August, September, October, November, December. Organic matter content in the surface horizon is about 2 percent. Nonirrigated land capability classification is 3w. This soil does not meet hydric criteria. The calcium carbonate equivalent within 40 inches, typically, does not exceed 2 percent. There are no saline horizons within 30 inches of the soil surface. The soil has a maximum sodium adsorption ratio of 1 within 30 inches of the soil surface.

Component: Felda (4%)

Generated brief soil descriptions are created for major components. The Felda soil is a minor component.

Component: Basinger (4%)

Generated brief soil descriptions are created for major components. The Basinger soil is a minor component.

Component: Riviera (4%)

Generated brief soil descriptions are created for major components. The Riviera soil is a minor component.

Component: Boca (3%)

Generated brief soil descriptions are created for major components. The Boca soil is a minor component.

Map Unit: 99—Water

Component: Water (100%)

Generated brief soil descriptions are created for major soil components. The Water is a miscellaneous area.

Map Unit: 100—Waters of the Gulf of Mexico

Component: Waters of the Gulf of Mexico (100%)

Generated brief soil descriptions are created for major soil components. The Waters of the Gulf of Mexico is a miscellaneous area.

Data Source Information

Soil Survey Area: Sarasota County, Florida
Survey Area Data: Version 10, Sep 9, 2014