

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT  
 Bullitt and Spencer Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
BeB	Beasley silt loam, 2 to 6 percent slopes	highly erodible
BeC	Beasley silt loam, 6 to 12 percent slopes, eroded	highly erodible
BeD	Beasley silt loam, 12 to 20 percent slopes, eroded	highly erodible
BfC3	Beasley silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
BfD3	Beasley silty clay loam, 12 to 20 percent slopes, severely eroded	highly erodible
Bo	Boonesboro silt loam, frequently flooded	not highly erodible
CaB	Caneyville silt loam, 2 to 6 percent slopes	highly erodible
CaC	Caneyville silt loam, 6 to 12 percent slopes, eroded	highly erodible
CbD	Caneyville-Beasley-Rock outcrop complex, 12 to 30 percent slopes	highly erodible
CnD	Caneyville-Rock outcrop complex, 6 to 20 percent slopes	highly erodible
CnE	Caneyville-Rock outcrop complex, 20 to 40 percent slopes	highly erodible
CrB	Crider silt loam, 2 to 6 percent slopes	not highly erodible
CrC	Crider silt loam, 6 to 12 percent slopes, eroded	highly erodible
CrD	Crider silt loam, 12 to 20 percent slopes, eroded	highly erodible
DAM	Dam, large	
EcC	Eden silty clay loam, 6 to 20 percent slopes, eroded	highly erodible
EdeE3	Eden flaggy silty clay, 20 to 30 percent slopes, severely eroded	highly erodible
EkA	Elk silt loam, 0 to 2 percent slopes	not highly erodible
EkB	Elk silt loam, 2 to 6 percent slopes	not highly erodible
EkC	Elk silt loam, 6 to 12 percent slopes	highly erodible
ElA	Elk silt loam, occasionally flooded, 0 to 2 percent slopes	not highly erodible
ElB	Elk silt loam, occasionally flooded, 2 to 6 percent slopes	not highly erodible
ElC	Elk silt loam, occasionally flooded, 6 to 12 percent slopes	highly erodible
ErA	Elk silt loam, 0 to 2 percent slopes, rarely flooded	not highly erodible
ErB	Elk silt loam, 2 to 6 percent slopes, rarely flooded	not highly erodible
ErC	Elk silt loam, 6 to 12 percent slopes, rarely flooded	highly erodible
FaC	Faywood silt loam, 6 to 12 percent slopes, eroded	highly erodible
FdD	Faywood silty clay loam, 12 to 20 percent slopes, eroded	highly erodible
FkF	Faywood-Beasley-Rock outcrop complex, 25 to 60 percent slopes	highly erodible
Fle	Faywood-Cynthiana complex, 12 to 30 percent slopes	highly erodible
FnF	Faywood-Fairmount-Woolper complex, 30 to 60 percent slopes	highly erodible
GmF	Garmon silt loam, 25 to 60 percent slopes	highly erodible
HaC	Hagerstown silt loam, 6 to 12 percent slopes, eroded	highly erodible
La	Lawrence silt loam, rarely flooded	not highly erodible
Le	Lawrence silt loam	not highly erodible
LfE	Lenberg-Carpenter complex, 20 to 40 percent slopes	highly erodible
LoB	Lowell silt loam, 2 to 6 percent slopes	highly erodible
LoC	Lowell silt loam, 6 to 12 percent slopes, eroded	highly erodible
LsC3	Lowell silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
MaB	Markland silt loam, rarely flooded, 2 to 6 percent slopes	highly erodible
MbD3	Markland silty clay, occasionally flooded, 10 to 30 percent slopes, severely eroded	highly erodible
Mc	McGary silt loam, rarely flooded	not highly erodible
Mo	Montgomery silty clay loam	not highly erodible
Mv	McGary variant silt loam, rarely flooded	not highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued  
 Bullitt and Spencer Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
Ne	Newark silt loam, frequently flooded	not highly erodible
NhA	Nicholson silt loam, 0 to 2 percent slopes	not highly erodible
NhB	Nicholson silt loam, 2 to 6 percent slopes	highly erodible
NhC	Nicholson silt loam, 6 to 12 percent slopes	highly erodible
No	Nolin silt loam, frequently flooded	not highly erodible
OtA	Otwell silt loam, 0 to 2 percent slopes	not highly erodible
OtB	Otwell silt loam, 2 to 6 percent slopes	highly erodible
OtC	Otwell silt loam, 6 to 12 percent slopes	highly erodible
OwB	Otwell silt loam, occasionally flooded, 2 to 6 percent slopes	highly erodible
Pt	Pits	
Sg	Sensabaugh gravelly loam, occasionally flooded	not highly erodible
ShB	Shelbyville silt loam, 2 to 6 percent slopes	not highly erodible
TrC	Trappist silt loam, 6 to 12 percent slopes, eroded	highly erodible
TrD	Trappist silt loam, 12 to 30 percent slopes, eroded	highly erodible
W	Water	
WoB	Woolper silty clay loam, 2 to 6 percent slopes	highly erodible
WoC	Woolper silty clay loam, 6 to 12 percent slopes	highly erodible
ZaB	Zanesville silt loam, 2 to 6 percent slopes	highly erodible
ZaC	Zanesville silt loam, 6 to 12 percent slopes	highly erodible