

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT  
 Garrard and Lincoln Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
AlB	Allegheny loam, 2 to 6 percent slopes, rarely flooded	not highly erodible
AlC2	Allegheny loam, 6 to 12 percent slopes, eroded	highly erodible
BaB	Beasley silt loam, 2 to 6 percent slopes	highly erodible
BbC2	Beasley silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
BeB	Berea silt loam, 2 to 6 percent slopes	highly erodible
uBlmB	Bluegrass-Maury silt loams, 2 to 6 percent slopes	not highly erodible
Bo	Boonesboro silt loam, frequently flooded	not highly erodible
CaE2	Caneyville silt loam, 12 to 30 percent slopes, eroded, rocky	highly erodible
CeB	Carpenter gravelly silt loam, 2 to 6 percent slopes	not highly erodible
CeC	Carpenter gravelly silt loam, 6 to 12 percent slopes	highly erodible
CgE2	Carpenter-Lenberg complex, 12 to 30 percent slopes, eroded	highly erodible
ChB	Chenault gravelly silt loam, 2 to 6 percent slopes	not highly erodible
ChC	Chenault gravelly silt loam, 6 to 12 percent slopes	highly erodible
CkC	Chenault-Lowell complex, phosphatic, 6 to 12 percent slopes	highly erodible
CldD2	Chenault-Faywood complex, phosphatic, 12 to 25 percent slopes, eroded, rocky	highly erodible
CmB	Christian silt loam, 2 to 6 percent slopes	highly erodible
CmC2	Christian silt loam, 6 to 12 percent slopes, eroded	highly erodible
CoD2	Christian silty clay loam, 12 to 25 percent slopes, eroded	highly erodible
CpF2	Colyer-Trappist complex, 25 to 60 percent slopes, eroded, very rocky	highly erodible
CrB	Crider silt loam, 2 to 6 percent slopes	not highly erodible
CrC	Crider silt loam, 6 to 12 percent slopes	highly erodible
CuB	Culleoka silt loam, 2 to 6 percent slopes	highly erodible
CuC2	Culleoka silt loam, 6 to 12 percent slopes, eroded	highly erodible
CuD2	Culleoka silt loam, 12 to 25 percent slopes, eroded	highly erodible
CyF2	Cynthiana-Faywood complex, 25 to 50 percent slopes, eroded, very rocky	highly erodible
DAM	Dam, large	
DoB	Donerail silt loam, 2 to 6 percent slopes	highly erodible
EddD2	Eden flaggy silty clay loam, 8 to 25 percent slopes, eroded	highly erodible
EffF2	Eden-Culleoka association, 25 to 50 percent slopes, eroded, stony	highly erodible
EkB	Elk silt loam, 2 to 6 percent slopes	highly erodible
EkC	Elk silt loam, 6 to 12 percent slopes	highly erodible
ErB	Elk silt loam, 2 to 6 percent slopes, rarely flooded	highly erodible
ErC	Elk silt loam, 6 to 12 percent slopes, rarely flooded	highly erodible
FaC2	Fairmount silty clay loam, 6 to 12 percent slopes, eroded, very rocky	highly erodible
FdF2	Fairmount-Faywood-Rock outcrop complex, 25 to 50 percent slopes, eroded	highly erodible
FecC2	Faywood-Cynthiana complex, 6 to 12 percent slopes, eroded, rocky	highly erodible
FedD2	Faywood-Cynthiana complex, 12 to 25 percent slopes, eroded, very rocky	highly erodible
FfcC2	Faywood-Fairmount complex, phosphatic, 6 to 12 percent slopes, eroded, rocky	highly erodible
FfdD2	Faywood-Fairmount complex, phosphatic, 12 to 25 percent slopes, eroded, very rocky	highly erodible
FoD2	Faywood-Shrouts complex, 12 to 25 percent slopes, eroded, rocky	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued  
 Garrard and Lincoln Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
FoF2	Faywood-Shrouths complex, 25 to 60 percent slopes, eroded, rocky	highly erodible
FrB	Frankstown gravelly silt loam, 2 to 6 percent slopes	highly erodible
FrC	Frankstown gravelly silt loam, 6 to 12 percent slopes	highly erodible
FrD2	Frankstown gravelly silt loam, 12 to 25 percent slopes, eroded	highly erodible
GaC2	Garlin-Shrouths complex, 6-12 percent slopes, eroded	highly erodible
GaD2	Garlin-Shrouths complex, 12 to 25 percent slopes, eroded, rocky	highly erodible
GmF	Garmon channery silt loam, 25 to 80 percent slopes, rocky	highly erodible
GnB	Gilpin silt loam, 2 to 6 percent slopes	highly erodible
GnC2	Gilpin silt loam, 6 to 12 percent slopes, eroded	highly erodible
GrB	Greenbriar silt loam, 2 to 6 percent slopes	highly erodible
HgC	Hagerstown silt loam, 6 to 12 percent slopes	highly erodible
JeB	Jessietown silt loam, 2 to 6 percent slopes	highly erodible
JeC	Jessietown silt loam, 6 to 12 percent slopes	highly erodible
Jm	Johnsburg-Mullins complex	not highly erodible
Jr	Johnsburg-Robertsville complex	not highly erodible
La	Lawrence silt loam, terrace, rarely flooded	not highly erodible
Le	Lawrence-Robertsville complex	not highly erodible
LgC2	Lenberg silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
LlB	Lily loam, 2 to 6 percent slopes	highly erodible
LlC	Lily loam, 6 to 12 percent slopes	highly erodible
LoB	Lowell silt loam, 2 to 6 percent slopes	highly erodible
LoC2	Lowell silt loam, 6 to 12 percent slopes, eroded	highly erodible
LpD2	Lowell-Faywood complex, 12 to 25 percent slopes, eroded, rocky	highly erodible
LsB	Lowell silt loam, phosphatic, 2 to 6 percent slopes	highly erodible
LsC2	Lowell silt loam, phosphatic, 6 to 12 percent slopes, eroded	highly erodible
LtD2	Lowell-Faywood complex, phosphatic, rocky, 12 to 25 percent slopes, eroded	highly erodible
uMlmc	Maury-Bluegrass silt loams, 6 to 12 percent slopes	highly erodible
Me	Melvin silt loam, frequently flooded	not highly erodible
MoB	Monongahela loam, 2 to 6 percent slopes	highly erodible
Ne	Newark silt loam, frequently flooded	not highly erodible
NhB	Nicholson silt loam, 2 to 6 percent slopes	highly erodible
NhC2	Nicholson silt loam, 6 to 12 percent slopes, eroded	highly erodible
No	Nolin silt loam, frequently flooded	not highly erodible
OtB	Otwell silt loam, 2 to 6 percent slopes	highly erodible
OwB	Otwell silt loam, 2 to 6 percent slopes, rarely flooded	highly erodible
PrB	Pricetown silt loam, 2 to 6 percent slopes	highly erodible
PrC	Pricetown silt loam, 6 to 12 percent slopes	highly erodible
Rb	Robertsville silt loam, terrace, rarely flooded	not highly erodible
RoF	Rock outcrop-Fairmount complex, 50 to 120 percent slopes	highly erodible
SaB	Sandview silt loam, 2 to 6 percent slopes	not highly erodible
SaC	Sandview silt loam, 6 to 12 percent slopes	highly erodible
SeC2	Shrouths silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
SfD3	Shrouths-Cynthiana complex, 12 to 25 percent slopes, severely eroded, rocky	highly erodible
SgF3	Shrouths-Garlin-Cynthiana complex, 25 to 50 percent slopes, severely eroded, very rocky	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued  
 Garrard and Lincoln Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
Sk	Skidmore very gravelly silt loam, frequently flooded	not highly erodible
TeB	Teddy silt loam, 2 to 6 percent slopes	highly erodible
TlB	Tilsit silt loam, 2 to 6 percent slopes	highly erodible
TlC	Tilsit silt loam, 6 to 12 percent slopes	highly erodible
TpB	Trappist silt loam, 2 to 6 percent slopes	highly erodible
TpC2	Trappist silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
TrD2	Trappist-Colyer complex, 12 to 25 percent slopes, eroded	highly erodible
W	Water	
Yo	Yosemite gravelly silt loam, frequently flooded	not highly erodible