

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT
 Henry and Trimble Counties, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
BaB	Beasley silt loam, 2 to 6 percent slopes	highly erodible
BeC2	Beasley silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
BeD2	Beasley silty clay loam, 12 to 20 percent slopes, eroded	highly erodible
BfC3	Beasley silty clay, 6 to 12 percent slopes, severely eroded	highly erodible
BfD3	Beasley silty clay, 12 to 20 percent slopes, severely eroded	highly erodible
Bo	Boonesboro silt loam, frequently flooded	not highly erodible
BsE2	Brassfield-Beasley complex, 20 to 40 percent slopes, eroded	highly erodible
CaB	Chenault silt loam, 2 to 6 percent slopes	not highly erodible
CaC	Chenault silt loam, 6 to 12 percent slopes	highly erodible
CcB	Cincinnati silt loam, 2 to 6 percent slopes	highly erodible
CcC	Cincinnati silt loam, 6 to 12 percent slopes	highly erodible
DAM	Dam, large	
EdC2	Eden silty clay loam, 6 to 20 percent slopes, eroded	highly erodible
EdE2	Eden silty clay loam, 20 to 35 percent slopes, eroded	highly erodible
EkA	Elk silt loam, occasionally flooded, 0 to 2 percent slopes	not highly erodible
EkB	Elk silt loam, 2 to 6 percent slopes	highly erodible
EkC	Elk silt loam, occasionally flooded, 6 to 12 percent slopes	highly erodible
En	Elk and Nolin silt loams, frequently flooded	highly erodible
ErB	Elk silt loam, 2 to 6 percent slopes, rarely flooded	highly erodible
FaE	Fairmount flaggy silty clay loam, 12 to 30 percent slopes, very rocky	highly erodible
FwF	Fairmount-Woolper complex, 30 to 65 percent slopes	highly erodible
FyC2	Faywood silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
FyD2	Faywood silty clay loam, 12 to 20 percent slopes, eroded	highly erodible
GbD2	Grayford-Beasley complex, 12 to 20 percent slopes, eroded	highly erodible
Hu	Huntington silt loam, occasionally flooded	not highly erodible
Lc	Lawrence silt loam, rarely flooded	not highly erodible
LoB	Lowell silt loam, 2 to 6 percent slopes	highly erodible
LoC	Lowell silt loam, 6 to 12 percent slopes	highly erodible
LsC3	Lowell silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
Mc	McGary silt loam	not highly erodible
Ne	Newark silt loam, frequently flooded	not highly erodible
NhB	Nicholson silt loam, 2 to 6 percent slopes	highly erodible
NhC	Nicholson silt loam, 6 to 12 percent slopes	highly erodible
No	Nolin silt loam, occasionally flooded	not highly erodible
OtA	Otwell silt loam, occasionally flooded, 0 to 2 percent slopes	not highly erodible
OtB	Otwell silt loam, rarely flooded, 2 to 6 percent slopes	highly erodible
Pt	Pits, quarries	
RoA	Rossmoyne silt loam, 0 to 2 percent slopes	not highly erodible
RoB	Rossmoyne silt loam, 2 to 6 percent slopes	highly erodible
RyB	Ryker silt loam, 2 to 6 percent slopes	not highly erodible
RyC	Ryker silt loam, 6 to 12 percent slopes	highly erodible
ShB	Shelbyville silt loam, 2 to 6 percent slopes	highly erodible
W	Water	
WeD	Wheeling loam, 6 to 20 percent slopes	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued
Henry and Trimble Counties, Kentucky: Detailed Soil Map Legend
(FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
WhB	Wheeling silt loam, 0 to 6 percent slopes	highly erodible
WoB	Woolper silty clay loam, 2 to 6 percent slopes	highly erodible
WoC	Woolper silty clay loam, 6 to 12 percent slopes	highly erodible
WoD	Woolper silty clay loam, 12 to 20 percent slopes	highly erodible