

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT  
 Hardin and Larue Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
A1D	Allegheny-Lenberg-Caneyville complex, 12 to 20 percent slopes	highly erodible
As	Ashton silt loam	not highly erodible
CnD	Caneyville-Rock outcrop complex, 6 to 20 percent slopes	highly erodible
CnE	Caneyville-Rock outcrop complex, 20 to 30 percent slopes	highly erodible
CrB	Crider silt loam, 2 to 6 percent slopes	not highly erodible
CrC	Crider silt loam, 6 to 12 percent slopes	highly erodible
CrD	Crider silt loam, 12 to 20 percent slopes	highly erodible
CsC	Cumberland silt loam, 6 to 12 percent slopes	highly erodible
CsD	Cumberland silt loam, 12 to 20 percent slopes	highly erodible
CtC3	Cumberland silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
CtD3	Cumberland silty clay loam, 12 to 20 percent slopes, severely eroded	highly erodible
DAM	Dam, large	
Dn	Dunning silty clay loam	not highly erodible
ElB	Elk silt loam, 2 to 6 percent slopes	not highly erodible
ElC	Elk silt loam, 6 to 12 percent slopes	highly erodible
FdC	Fredonia-Rock outcrop complex, 6 to 20 percent slopes	highly erodible
FrC	Frondorf-Lenberg silt loams, 6 to 12 percent slopes	highly erodible
FrD	Frondorf-Lenberg silt loams, 12 to 20 percent slopes	highly erodible
GmE	Garmon silt loam, 25 to 60 percent slopes	highly erodible
GnB	Gatton silt loam, 2 to 6 percent slopes	highly erodible
Gu	Gullied land (riny)	highly erodible
HnB	Hagerstown silt loam, 2 to 6 percent slopes	not highly erodible
HnC	Hagerstown silt loam, 6 to 12 percent slopes	highly erodible
HnD	Hagerstown silt loam, 12 to 20 percent slopes	highly erodible
Hu	Huntington silt loam	not highly erodible
Lc	Lawrence silt loam	not highly erodible
LfE	Lenberg-Frondorf complex, 20 to 30 percent slopes	highly erodible
Ln	Lindside silt loam	not highly erodible
MdC3	Markland silty clay, 6 to 12 percent slopes, severely eroded	highly erodible
Mr	McGary silt loam	not highly erodible
Mv	Melvin silt loam	not highly erodible
Nb	Newark silt loam	not highly erodible
NcA	Nicholson silt loam, 0 to 2 percent slopes	not highly erodible
NcB	Nicholson silt loam, 2 to 6 percent slopes	highly erodible
No	Nolin silt loam	not highly erodible
Nv	Nolin variant fine sandy loam (grigsby)	not highly erodible
OtA	Otwell silt loam, 0 to 2 percent slopes	not highly erodible
OtB	Otwell silt loam, 2 to 6 percent slopes	highly erodible
PmB	Pembroke silt loam, 2 to 6 percent slopes	not highly erodible
PmC	Pembroke silt loam, 6 to 12 percent slopes	highly erodible
Pt	Pits, quarries	
RaE	Ramsey-Steinsburg-Allegheny complex, 20 to 40 percent slopes	highly erodible
RbC	Riny loam, 6 to 12 percent slopes	highly erodible
RbD	Riny loam, 12 to 20 percent slopes	highly erodible
RbE	Riny loam, 20 to 30 percent slopes	highly erodible
RcD3	Riny sandy clay loam, 6 to 20 percent slopes, severely eroded	highly erodible
Rd	Robertsville silt loam	not highly erodible
RoE	Rock outcrop-Corydon complex, 12 to 30 percent slopes	highly erodible
SdA	Sadler silt loam, 0 to 2 percent slopes	not highly erodible
SdB	Sadler silt loam, 2 to 6 percent slopes	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued  
 Hardin and Larue Counties, Kentucky: Detailed Soil Map Legend  
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
SdC	Sadler silt loam, 6 to 12 percent slopes	highly erodible
Sg	Sensabaugh silt loam	not highly erodible
SnB	Sonora silt loam, 2 to 6 percent slopes	highly erodible
SnC	Sonora silt loam, 6 to 12 percent slopes	highly erodible
SnC3	Sonora silt loam, 6 to 12 percent slopes, severely eroded	highly erodible
VrC	Vertrees silt loam, 6 to 12 percent slopes	highly erodible
VrD	Vertrees silt loam, 12 to 20 percent slopes	highly erodible
VrE	Vertrees silt loam, 20 to 30 percent slopes	highly erodible
VtD3	Vertrees silty clay loam, 6 to 20 percent slopes, severely eroded	highly erodible
W	Water	
WbC	Waynesboro loam, 6 to 12 percent slopes	highly erodible
WbD	Waynesboro loam, 12 to 20 percent slopes	highly erodible
WbE	Waynesboro loam, 20 to 30 percent slopes	highly erodible
WcC3	Waynesboro clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
WcD3	Waynesboro clay loam, 12 to 20 percent slopes, severely eroded	highly erodible
WlB	Wellston silt loam, 2 to 6 percent slopes	highly erodible
WlC	Wellston silt loam, 6 to 12 percent slopes	highly erodible
WlC3	Wellston silt loam, 6 to 12 percent slopes, severely eroded	highly erodible