

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT
 Breckinridge and Meade Counties, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
BaB2	Baxter very gravelly silt loam, karst, 2 to 6 percent slopes, eroded	highly erodible
BaC2	Baxter very gravelly silt loam, karst, 6 to 12 percent slopes, eroded	highly erodible
BaD2	Baxter very gravelly silt loam, karst, 12 to 20 percent slopes, eroded	highly erodible
BaE2	Baxter very gravelly silt loam, karst, 20 to 30 percent slopes, eroded	highly erodible
BbC3	Baxter very gravelly silty clay loam, karst, 6 to 12 percent slopes, severely eroded	highly erodible
BbD3	Baxter very gravelly silty clay loam, karst, 12 to 20 percent slopes, severely eroded	highly erodible
BeC4	Baxter soils, karst, 6 to 12 percent slopes, very severely eroded	highly erodible
BeD4	Baxter soils, karst, 12 to 20 percent slopes, very severely eroded	highly erodible
CaC2	Caneyville silt loam, 6 to 12 percent slopes, eroded	highly erodible
CaD2	Caneyville silt loam, 12 to 20 percent slopes, eroded	highly erodible
CeC3	Caneyville silty clay, 6 to 12 percent slopes, severely eroded	highly erodible
CeD3	Caneyville silty clay, 12 to 20 percent slopes, severely eroded	highly erodible
CkD	Caneyville-Rock outcrop complex, 12 to 30 percent slopes	highly erodible
Cn	Chagrin fine sandy loam, occasionally flooded	not highly erodible
Co	Clifty gravelly silt loam, 0 to 2 percent slopes, occasionally flooded	not highly erodible
CrB2	Crider silt loam, 2 to 6 percent slopes, eroded	highly erodible
CrC2	Crider silt loam, 6 to 12 percent slopes, eroded	highly erodible
CrD2	Crider silt loam, 12 to 20 percent slopes, eroded	highly erodible
CtC3	Crider silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
CtD3	Crider silty clay loam, 12 to 20 percent slopes, severely eroded	highly erodible
Cu	Cuba silt loam, occasionally flooded	not highly erodible
DAM	Dam, large	
EkA	Elk silt loam, 0 to 2 percent slopes, rarely flooded	not highly erodible
EkB	Elk silt loam, 2 to 6 percent slopes, rarely flooded	highly erodible
EkC2	Elk silt loam, 6 to 12 percent slopes, eroded	highly erodible
EkD2	Elk silt loam, 12 to 20 percent slopes, eroded	highly erodible
EkD3	Elk silt loam, 12 to 20 percent slopes, severely eroded	highly erodible
EkE	Elk silt loam, 20 to 40 percent slopes, rarely flooded	highly erodible
FcC2	Fredonia-Crider complex, karst, rocky, 6 to 12 percent slopes, eroded	highly erodible
FcD2	Fredonia-Crider complex, karst, rocky, 12 to 20 percent slopes, eroded	highly erodible
FrD3	Fredonia-Crider complex, karst, very rocky, 6 to 20 percent slopes, severely eroded	highly erodible
GaB2	Gatton silt loam, 2 to 6 percent slopes, eroded	highly erodible
Glc2	Gilpin silt loam, 6 to 12 percent slopes, eroded	highly erodible
Glc3	Gilpin silt loam, 6 to 12 percent slopes, severely eroded	highly erodible
GwF	Gilpin-Dekalb-Rock outcrop complex, 30 to 60 percent slopes	highly erodible
HaB2	Hammack silt loam, 2 to 6 percent slopes, eroded	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued
 Breckinridge and Meade Counties, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
HbC2	Hammack-Baxter complex, karst, 6 to 12 percent slopes, eroded	highly erodible
HbC3	Hammack-Baxter complex, karst, 6 to 12 percent slopes, severely eroded	highly erodible
Hu	Huntington silt loam, occasionally flooded	not highly erodible
LaB	Lakin loamy fine sand, 2 to 6 percent slopes	not highly erodible
LaC	Lakin loamy fine sand, 6 to 15 percent slopes	highly erodible
Ld	Lindside silt loam, occasionally flooded	not highly erodible
Ln	Lindside silt loam, depressional, frequently flooded	not highly erodible
MaC3	Markland silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
MaD3	Markland silty clay loam, 12 to 35 percent slopes, severely eroded	highly erodible
Mc	McGary silt loam	not highly erodible
Me	Melvin silt loam, occasionally flooded	not highly erodible
Mf	Melvin silt loam, depressional, frequently flooded	not highly erodible
Na	Newark silt loam, occasionally flooded	not highly erodible
Ne	Newark silt loam, depressional, frequently flooded	not highly erodible
NhB2	Nicholson silt loam, 2 to 6 percent slopes, eroded	highly erodible
NhC2	Nicholson silt loam, 6 to 12 percent slopes, eroded	highly erodible
NkC4	Nicholson soils, 4 to 12 percent slopes, very severely eroded	highly erodible
No	Nolin silt loam, 0 to 2 percent slopes, occasionally flooded	not highly erodible
Nv	Nolin silt loam, depressional, frequently flooded	not highly erodible
PeA	Pekin silt loam, 0 to 2 percent slopes	not highly erodible
PeB	Pekin silt loam, 2 to 6 percent slopes	highly erodible
Pt	Pits, quarries	
RaC2	Riney loam, 6 to 12 percent slopes, eroded	highly erodible
ReD	Riney-Lily complex, 12 to 20 percent slopes	highly erodible
ReE	Riney-Lily complex, 20 to 30 percent slopes	highly erodible
RkF	Rock outcrop-Caneyville complex, 30 to 90 percent slopes	highly erodible
RmD	Rock outcrop-Corydon complex, 12 to 30 percent slopes	highly erodible
RnC2	Rosine silt loam, 6 to 12 percent slopes, eroded	highly erodible
RoC3	Rosine silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
RSD2	Rosine-Gilpin-Lenberg complex, 12 to 20 percent slopes, eroded	highly erodible
RSD3	Rosine-Gilpin-Lenberg complex, 12 to 20 percent slopes, severely eroded	highly erodible
RsE	Rosine-Gilpin-Lenberg complex, very rocky, 20 to 30 percent slopes	highly erodible
SaA	Sadler silt loam, 0 to 2 percent slopes	not highly erodible
SaB2	Sadler silt loam, 2 to 6 percent slopes, eroded	highly erodible
ScA	Sciotoville silt loam, 0 to 2 percent slopes	not highly erodible
ScB	Sciotoville silt loam, 2 to 6 percent slopes	highly erodible
Sf	Steff silt loam, occasionally flooded	not highly erodible
St	Stendal silt loam, occasionally flooded	not highly erodible
uAlfB2	Alford silt loam, 2 to 6 percent slopes, eroded	highly erodible
uAlfC2	Alford silt loam, 6 to 12 percent slopes, eroded	highly erodible
uAlfD2	Alford silt loam, 12 to 20 percent slopes, eroded	highly erodible
uHosB2	Hosmer silt loam, 2 to 6 percent slopes, eroded	highly erodible
uHosC2	Hosmer silt loam, 6 to 12 percent slopes, eroded	highly erodible
uRobA	Robbs silt loam, 0 to 2 percent slopes	not highly erodible
VrF	Varilla-Gilpin-Rock outcrop complex, very bouldery, 20 to 65 percent slopes	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued
 Breckinridge and Meade Counties, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
W	Water	
We	Weinbach silt loam	not highly erodible
WtF	Westmoreland-Caneyville-Rock outcrop complex, 30 to 80 percent slopes	highly erodible
WxB	Wheeling fine sandy loam, 2 to 6 percent slopes	highly erodible
WxC2	Wheeling fine sandy loam, 6 to 12 percent slopes, eroded	highly erodible
Ya	Yeager loamy sand, occasionally flooded	not highly erodible
ZaB2	Zanesville silt loam, 2 to 6 percent slopes, eroded	highly erodible
ZaC2	Zanesville silt loam, 6 to 12 percent slopes, eroded	highly erodible
ZnC3	Zanesville silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible