

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT
 Metcalfe County, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
BaB	Baxter cherty silt loam, 2 to 6 percent slopes	not highly erodible
BaB2	Baxter cherty silt loam, 2 to 6 percent slopes, eroded	not highly erodible
BaC	Baxter cherty silt loam, 6 to 12 percent slopes	highly erodible
BaC2	Baxter cherty silt loam, 6 to 12 percent slopes, eroded	highly erodible
BaD	Baxter cherty silt loam, 12 to 20 percent slopes	highly erodible
BaD2	Baxter cherty silt loam, 12 to 20 percent slopes, eroded	highly erodible
BaE	Baxter cherty silt loam, 20 to 30 percent slopes	highly erodible
BaE2	Baxter cherty silt loam, 20 to 30 percent slopes, eroded	highly erodible
BcC3	Baxter cherty silty clay loam, 6 to 12 percent slopes, severely eroded	highly erodible
BcD3	Baxter cherty silty clay loam, 12 to 20 percent slopes, severely eroded	highly erodible
BcE3	Baxter cherty silty clay loam, 20 to 30 percent slopes, severely eroded	highly erodible
BeC2	Baxter-Talbott rocky silt loams, 6 to 12 percent slopes, eroded (baxter, caneyville rocky)	highly erodible
BeD2	Baxter-Talbott rocky silt loams, 12 to 20 percent slopes, eroded (baxter, caneyville rocky)	highly erodible
BeE2	Baxter-Talbott rocky silt loams, 20 to 30 percent slopes, eroded (baxter, caneyville rocky)	highly erodible
BfD3	Baxter-Talbott rocky silty clay loams, 12 to 20 percent slopes, severely eroded (baxter, caneyville rocky)	highly erodible
BoD	Bodine cherty silt loam, 12 to 20 percent slopes	highly erodible
BoE	Bodine cherty silt loam, 20 to 35 percent slopes	highly erodible
CaE	Caneyville rocky complex, 20 to 30 percent slopes	highly erodible
CaE3	Caneyville rocky complex, 20 to 30 percent slopes, severely eroded	highly erodible
CaF	Caneyville rocky complex, 30 to 50 percent slopes	highly erodible
CbA	Captina silt loam, 0 to 2 percent slopes	not highly erodible
CbB	Captina silt loam, 2 to 6 percent slopes	highly erodible
CcD3	Christian clay loam, 6 to 20 percent slopes, severely eroded	highly erodible
CdB	Christian loam, 2 to 6 percent slopes	highly erodible
CdC	Christian loam, 6 to 12 percent slopes	highly erodible
CdC2	Christian loam, 6 to 12 percent slopes, eroded	highly erodible
CdD2	Christian loam, 12 to 20 percent slopes, eroded	highly erodible
CeD	Christian rocky soils, 12 to 20 percent slopes (caneyville rocky)	highly erodible
CkB	Clarksville cherty silt loam, 2 to 6 percent slopes	not highly erodible
CkC	Clarksville cherty silt loam, 6 to 12 percent slopes	highly erodible
CkC2	Clarksville cherty silt loam, 6 to 12 percent slopes, eroded	highly erodible
CkD2	Clarksville cherty silt loam, 12 to 20 percent slopes, eroded	highly erodible
CkE2	Clarksville cherty silt loam, 20 to 30 percent slopes, eroded	highly erodible
CrB	Crider silt loam, 2 to 6 percent slopes	not highly erodible
CrB2	Crider silt loam, 2 to 6 percent slopes, eroded	not highly erodible
CrC2	Crider silt loam, 6 to 12 percent slopes, eroded	highly erodible
CuB	Cumberland cherty silt loam, 2 to 6 percent slopes (frederick)	not highly erodible
CuB2	Cumberland cherty silt loam, 2 to 6 percent slopes, eroded (frederick)	not highly erodible
CuC2	Cumberland cherty silt loam, 6 to 12 percent slopes, eroded (frederick)	highly erodible

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT--Continued
 Metcalfe County, Kentucky: Detailed Soil Map Legend
 (FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
CuD2	Cumberland cherty silt loam 12 to 20 percent slopes, eroded (frederick)	highly erodible
CvD3	Cumberland cherty silty clay, 12 to 20 percent slopes, severely eroded (frederick)	highly erodible
DaD	Dandridge and Westmoreland shaly silt loams, 12 to 20 percent slopes (dandridge, garmon)	highly erodible
DaF	Dandridge and Westmoreland shaly silt loams, 20 to 50 percent slopes (dandridge, garmon)	highly erodible
DAM	Dam, large	
DbD3	Dandridge and Westmoreland shaly silty clay loams, 12 to 20 percent slopes, severely eroded (dandridge, garmon)	highly erodible
DbF3	Dandridge and Westmoreland shaly silty clay loams, 20 to 50 percent slopes, severely eroded (dandridge, garmon)	highly erodible
DcB	Dandridge and Westmoreland silt loams, 2 to 6 percent slopes (dandridge, garmon)	highly erodible
DcC	Dandridge and Westmoreland silt loams, 6 to 12 percent slopes (dandridge, garmon)	highly erodible
DeB	Dewey silt loam, 2 to 6 percent slopes	not highly erodible
DeC2	Dewey silt loam, 6 to 12 percent slopes, eroded	highly erodible
DeD2	Dewey silt loam, 12 to 20 percent slopes, eroded	highly erodible
DkA	Dickson silt loam, 0 to 2 percent slopes	not highly erodible
DkB	Dickson silt loam, 2 to 6 percent slopes	highly erodible
DkB2	Dickson silt loam, 2 to 6 percent slopes, eroded	highly erodible
DkC2	Dickson silt loam, 6 to 12 percent slopes, eroded	highly erodible
EkB	Elk silt loam, 2 to 6 percent slopes	not highly erodible
Gu	Gullied land	
HcB	Humphreys cherty silt loam, 2 to 6 percent slopes	not highly erodible
HcC	Humphreys cherty silt loam, 6 to 12 percent slopes	highly erodible
HcC2	Humphreys cherty silt loam, 6 to 12 percent slopes, eroded	highly erodible
Hg	Huntington gravelly silt loam (sensabaugh)	not highly erodible
Hu	Huntington silt loam	not highly erodible
LaC2	Landisburg cherty silt loam, 6 to 12 percent slopes, eroded (tarklin)	highly erodible
LdB	Landisburg silt loam, 2 to 6 percent slopes (captina)	highly erodible
LdC2	Landisburg silt loam, 6 to 12 percent slopes, eroded (captina)	highly erodible
Ls	Lindside silt loam	not highly erodible
Me	Melvin silt loam	not highly erodible
MoB	Mountview silt loam, 2 to 6 percent slopes	not highly erodible
MoC2	Mountview silt loam, 6 to 12 percent slopes, eroded	highly erodible
MuC	Muse silt loam, 6 to 12 percent slopes	highly erodible
Nk	Newark silt loam	not highly erodible
PmB	Pembroke silt loam, 2 to 6 percent slopes	not highly erodible
PmC2	Pembroke silt loam, 6 to 12 percent slopes, eroded	highly erodible
Pt	Pits, quarries	
RaC	Renox silt loam, 6 to 12 percent slopes	highly erodible
Rb	Robertsville silt loam	not highly erodible
Rf	Robinsonville fine sandy loam	not highly erodible
Rk	Rock land (rock outcrop)	
SaA	Sango silt loam, 0 to 2 percent slopes	not highly erodible
SaB	Sango silt loam, 2 to 6 percent slopes	highly erodible
Ta	Taft silt loam	not highly erodible
TbC	Talbott silt loam, 6 to 12 percent slopes (caneyville rocky)	highly erodible
TcC2	Talbott silty clay loam, 6 to 12 percent slopes, eroded (caneyville rocky)	highly erodible
W	Water	