

HIGHLY ERODIBLE LAND CLASSIFICATION REPORT
 Montgomery County, Kentucky: Detailed Soil Map Legend
 FOR OFFICE DETERMINATIONS ONLY)

Map Symbol	Soil Mapunit Name	HEL Classification
AaB	Aaron silt loam, 2 to 6 percent slopes	highly erodible
AaC	Aaron silt loam, 6 to 12 percent slopes	highly erodible
BaB	Beasley silt loam, 2 to 6 percent slopes	highly erodible
BaC	Beasley silt loam, 6 to 12 percent slopes	highly erodible
BcF	Bledsoe-Rock outcrop-Caneyville complex, 6 to 60 percent slopes	highly erodible
BlE	Bledsoe-Shelocta complex, 12 to 35 percent slopes	highly erodible
Bo	Boonesboro silt loam, frequently flooded	not highly erodible
BrC	Brassfield silty clay loam, 6 to 12 percent slopes, eroded	highly erodible
BsE	Brassfield-Shrouts-Beasley complex, 12 to 35 percent slopes, eroded	highly erodible
CoF	Colyer-Trappist complex, 25 to 60 percent slopes, eroded	highly erodible
CrB	Crider silt loam, 2 to 6 percent slopes	not highly erodible
CrC	Crider silt loam, 6 to 15 percent slopes	highly erodible
Cu	Cuba silt loam, occasionally flooded	not highly erodible
DAM	Dam	
EkB	Elk silt loam, 2 to 6 percent slopes	not highly erodible
EkC	Elk silt loam, 6 to 12 percent slopes	highly erodible
ErB	Elk silt loam, 2 to 6 percent slopes, rarely flooded	not highly erodible
ErC	Elk silt loam, 6 to 12 percent slopes, rarely flooded	highly erodible
FaC	Faywood silt loam, 6 to 12 percent slopes	highly erodible
FaD	Faywood silt loam, 12 to 20 percent slopes	highly erodible
FcE	Faywood-Cynthiana complex, 12 to 35 percent slopes, eroded	highly erodible
Fle	Faywood-Lowell complex, 12 to 35 percent slopes	highly erodible
Hu	Huntington silt loam, occasionally flooded	not highly erodible
LaF	Latham-Shelocta-Rock outcrop complex, 20 to 60 percent slopes	highly erodible
LeC	Lenberg silt loam, 6 to 12 percent slopes, eroded	highly erodible
LeD	Lenberg silt loam, 12 to 20 percent slopes, eroded	highly erodible
LnF	Lenberg-Shelocta complex, 20 to 50 percent slopes, eroded	highly erodible
LoB	Lowell silt loam, 2 to 6 percent slopes	highly erodible
LoC	Lowell silt loam, 6 to 12 percent slopes	highly erodible
LoD	Lowell silt loam, 12 to 20 percent slopes	highly erodible
Me	Melvin silt loam, frequently flooded	not highly erodible
Mo	Morehead silt loam, rarely flooded	not highly erodible
NiB	Nicholson silt loam, 2 to 8 percent slopes	highly erodible
OtB	Otwell silt loam, 2 to 8 percent slopes	highly erodible
Pe	Peoga silt loam, rarely flooded	highly erodible
Pt	Pits-Dumps complex	
Ro	Robertsville silt loam	not highly erodible
Sc	Sees silt loam, 0 to 4 percent slopes, rarely flooded	not highly erodible
SeB	Shelbyville silt loam, 2 to 6 percent slopes	highly erodible
SeC	Shelbyville silt loam, 6 to 12 percent slopes	highly erodible
ShC	Shelocta silt loam, 6 to 15 percent slopes	highly erodible
SLF	Shelocta-Berks complex, 20 to 60 percent slopes	highly erodible
St	Stendal silt loam, frequently flooded	not highly erodible
TiB	Tilsit silt loam, 0 to 6 percent slopes	highly erodible
TiC	Tilsit silt loam, 6 to 12 percent slopes	highly erodible
TrC	Trappist silt loam, 6 to 12 percent slopes, eroded	highly erodible
TsE	Trappist-Colyer-Shelocta complex, 12 to 30 percent slopes, eroded	highly erodible
Ty	Tyler silt loam	not highly erodible
W	Water	
WoB	Woolper silty clay loam, 2 to 8 percent slopes	highly erodible