

Ecological Site Description

Alfic Wet Footslope Forest

F134XY014MO


- (*Liquidambar styraciflua* - *Quercus pagoda*/*Ilex decidua*/*Carex* - *Chasmanthium latifolium*)
- (sweetgum – cherrybark oak/possum haw/sedge – river oats)

An Ecological Site Description (ESD) is a reference document of ecological knowledge regarding a particular land area (ecological site). An ESD describes ecological potential and ecosystem dynamics of land areas and their potential management. Ecological sites are linked to soil survey map unit components, which allows for mapping of ecological sites. (**NOTE:** *This is a “provisional” ESD, and is subject to change. It contains basic ecological information sufficient for conservation planning and land management in Missouri. After additional information is developed and reviewed, a “Correlated” ESD will be published and will be available via the Web Soil Survey <http://websoilsurvey.nrcs.usda.gov> .)*)

Major Land Resource Area: 134 – Southern Mississippi Valley Loess

Introduction

The Southern Mississippi Valley Loess (area outlined in red on the map; northern portion only) is a relatively narrow strip of the coastal plain bordering the Mississippi River valley, that is blanketed with loess. The northern part of this MLRA, discussed here, is locally referred to as Crowley’s Ridge. Elevation ranges from about 300 feet on the footslopes to nearly 600 feet on the highest ridges. Loess caps the summits and upper slopes, and Pliocene-aged sand and gravel deposits of the coastal plain influence soils on lower, steeper slopes.



Alfic Wet Footslope Forests are within the green areas on the map (Missouri portion only; distributions farther south are currently under review). These sites are extensive on footslopes and drainageways on Crowley’s Ridge and adjacent areas. Soils are very deep, with a seasonal high water table.

Physiographic Features

This site is on footslopes and drainageways with slopes of less than 3%. The site receives runoff from adjacent upland sites. Some areas are subject to flooding.

Soil Features

These soils have no rooting restriction. The soils were formed under forest vegetation, and have thin, light-colored surface horizons. Parent material is colluvium or local alluvium from loess. They have silt loam surface horizons, with loamy to clayey subsoils. They are affected by a seasonal high water table during the spring months. Soil series associated with this site include Calhoun, Dundee, Falaya, Overcup, and Zachary.

Ecological Dynamics

Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information presented is representative of very complex vegetational communities. Not all scenarios or plants are included. Key indicator plants, animals and ecological processes are described to help inform land management decisions.

Alfic Wet Footslope Forests in this region are on footslope and drainageway positions. The historic reference plant community is dominated by a wide variety of deciduous hardwood tree species, tolerant of seasonally wet conditions including bur oak, cherrybark oak, willow oak, sweetgum, pin oak, Nuttall oak, water oak, American elm, sugarberry, and green ash. Trees are generally large and tall forming a dense, closed canopy. Both historically and today, these forests are structurally and compositionally diverse, with occasional tree-fall gaps and natural mortality providing opportunities for regeneration of overstory species. The understory is also complex, with multiple layers of shade tolerant species such as blue beech, spicebush, and Ohio buckeye. Grape vines, greenbriar, and trumpet creeper are also present along with a diverse array of ground flora species that carpets the forest floor.

While the upland woodlands had an estimated fire frequency of 5-10 years, these sites burned much less frequently (estimated 10-25 years) and with lower intensity. Alfic Wet Footslope Forests are also subject to occasional disturbances from wind and ice, which periodically open the canopy up by knocking over trees or breaking substantial branches of canopy trees. Such canopy disturbances allow more light to reach the ground and favor reproduction of the dominant oak species.

Today, these communities have been cleared and converted to pasture, or have undergone repeated timber harvest and domestic grazing. Most existing occurrences have a younger (50-80 years) canopy layer whose composition may have been altered by timber harvesting practices. These blocks of forest play an important role as a source of food and shelter for migrating birds.

Alfic Wet Footslope Forests are productive timber sites. Timber harvest in this region typically is done using single-tree selection, and often results in removal of the most productive trees, or high-grading of the stand. This can result in poorer quality timber and a shift in species composition away from more valuable oak species.

Carefully planned single tree selection or the creation of group openings can help regenerate more desirable oak species and increase vigor on the residual trees. Clear-cutting does occur and results in dense, even-aged stands of primarily oak. This may be most beneficial for existing stands whose composition has been highly altered by past management practices. However, without some thinning of the dense stands, the ground flora diversity can be shaded out and productivity of the stand may suffer.

Prescribed fire can play a beneficial but limited role in the management of this ecological site. The higher productivity of these sites makes it more challenging than on other forest and woodland sites in the region. Control of woody species will be more difficult. Footslope forests did evolve with some fire, but their composition often reflects more closed, forested conditions, with fewer woodland ground flora species that can respond to fire. Consequently, while having these sites in a burn unit is acceptable, targeting them solely for woodland restoration is not advisable.

Reference State Plant Community

Canopy Trees

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
BUR OAK	<i>Quercus macrocarpa</i>	5-20	80
GREEN ASH	<i>Fraxinus pennsylvanica</i>	5-20	80
WILLOW OAK	<i>Quercus lyrata</i>	5-20	70
WATER OAK	<i>Quercus nigra</i>	5-20	70
NUTTALL OAK	<i>Quercus nuttallii</i>	5-20	70
SUGARBERRY	<i>Celtis laevigata</i>	5-20	70
SWEETGUM	<i>Liquidambar styraciflua</i>	5-20	90
SWAMP CHESTNUT OAK	<i>Quercus michauxii</i>	5-20	80
SHELLBARK HICKORY	<i>Carya lacinosa</i>	5-20	70
AMERICAN ELM	<i>Ulmus americana</i>	5-20	60
SYCAMORE	<i>Platanus occidentalis</i>	5-20	90
CHERRYBARK OAK	<i>Quercus pagoda</i>	5-20	80
PECAN	<i>Carya illinoensis</i>	5-20	70

Understory Trees

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
OHIO BUCKEYE	<i>Aesculus glabra</i>	10-20	40
BLUE BEECH	<i>Carpinus caroliniana</i>	10-20	20

Shrubs

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
POSSUM HAW	<i>Ilex decidua</i>	10-20	8
SPICE BUSH	<i>Lindera benzoin</i>	10-20	6

Vines

Common Name	Botanical Name	Cover % (low-high)
TRUMPET CREEPER	<i>Campis radicans</i>	10-20
RACCOON GRAPE	<i>Ampelopsis cordata</i>	10-20
FOX GRAPE	<i>Vitis vulpina</i>	10-20
GREENBRIAR	<i>Smilax glauca</i>	10-20

Forbs

Common Name	Botanical Name	Cover % (low-high)
PALE TOUCH-ME-NOT	<i>Impatiens pallida</i>	5-20
FALSE NETTLE	<i>Boehmeria cylindrica</i>	5-20
CLEARWEED	<i>Pilea pumila</i>	5-20
WHITE WOODLAND ASTER	<i>Aster lateriflorus</i>	5-20
WOOD NETTLE	<i>Laportea canadensis</i>	5-20
WATERLEAF	<i>Hydrophyllum virginianum</i>	5-20
HISPID BUTTERCUP	<i>Ranunculus hispidus</i>	5-20
SKULLCAP	<i>Scutellaria nervosa</i>	5-20

Grasses and sedges

Common Name	Botanical Name	Cover % (low-high)
RIVER OATS	<i>Chasmanthium latifolium</i>	5-10
HOP SEDGE	<i>Carex lupulina</i>	5-10
WOOD REED GRASS	<i>Cinna arundinacea</i>	5-10
SMOOTH CONE SEDGE	<i>Carex laeviconica</i>	5-10

Site Interpretations

Wildlife Species

- Moist conditions with abundant coarse woody debris make this type of ecological site important for many herptiles.
- Ephemeral pools provide important amphibian breeding habitat. Periodic inundation and acorns provide important habitat and food for migrating ducks (especially mallards) and breeding ducks including wood ducks and hooded mergansers.
- Tall emergent trees along with an uneven canopy structure and canopy gaps are important for heron colonies, eagle nesting, Mississippi kites, cerulean warblers and other bird species.
- Birds associated with late-successional to mature forests are Wood Duck, Hooded Merganser, Barred Owl, Cerulean Warbler, Yellow-throated Warbler, Prothonotary Warbler, Pileated Woodpecker, Yellow-throated Vireo, Brown Creeper, and Yellow-crowned Night Heron.
- Reptiles and amphibians associated with ecological site include: small-mouthed salamander, central newt, midland brown snake, gray treefrog, northern spring peeper, Blanchard's cricket frog, southern leopard frog, western painted turtle, and red-eared slider.

Glossary

Alfic – soil that has a clay-dominated subsoil (argillic horizon) with moderate to high amounts of bases such as calcium, and were typically formed under woody vegetation.

Backslope – a hillslope profile position that forms the steepest and generally linear, middle portion of the slope.

Backswamp – marshy or swampy, depressed areas of flood plains between natural levees and valley sides or terraces

Calcareous – the presence of calcium carbonate in the soil parent material within the rooting zone; relatively alkaline

Claypan – a dense, compact, slowly permeable layer in the subsoil having much higher clay content than the overlying material

Chert – hard, extremely dense or compact crystalline sedimentary rock, consisting dominantly of interlocking crystals of quartz

Cliff – a significant vertical, or near vertical, rock exposure

Dolomite – a type of sedimentary rock that is a carbonate mineral composed of calcium magnesium carbonate

Drainageway – the upper most reach of a stream channel system characterized by little meandering

Dry – a site where soil moisture is limiting during the growing season; low available water capacity

Dune – a low mound, ridge, bank or hill of loose, wind-blown sand

Exposed – steep, south and west-facing slopes, which are warmer and drier than other slope aspects

Flatwoods – a type of woodland that occurs on soils with a root restricting subsoil layer within 20 to 30 inches, resulting in very slow runoff and ponding that remains saturated for most of the winter and early spring months but dries out and becomes very dry in the summer months; plants that grow there must be adapted to both conditions

Floodplain – the nearly level plain that borders a stream and is subject to inundation under flood-stage conditions

Footslope – a hillslope position at the base of a slope where hillslope sediment (colluvium) accumulates

Forest – a vegetative community dominated by trees forming a closed canopy and interspersed with shade-tolerant understory species

Fragipan – a dense, brittle subsoil horizon that is extremely hard and compact when dry

Glade – open, rocky, barren vegetative community dominated by drought-adapted forbs and grasses, typically with scattered, stunted woody plants

Igneous –bedrock formed by cooling and solidification of magma. Granite and rhyolite are typical igneous bedrocks in Missouri

Limestone – a type of sedimentary rock composed largely of calcium carbonate

Loess – material transported and deposited by wind and consisting predominantly of silt-size particles

Loamy – soil material containing a relatively equal mixture of sand and silt and a somewhat smaller proportion of clay

Marsh – a type of wetland that is dominated by herbaceous rather than woody plant species

Moist – a site that is moderately well to well drained and has high available water capacity, resulting in a well-balanced supply of moisture (neither too dry nor too wet).

Mollic – soil that has a thick, dark surface horizon and was typically formed under prairie vegetation

Mudstone – blocky or massive, fine-grained sedimentary rock in which the proportions of clay and silt are approximately equal

Natric – a soil horizon that displays a blocky, columnar, or prismatic structure and has a subhorizon with an exchangeable-sodium saturation of over 15%

Outwash – stratified sediments of sand and gravel removed or “washed out” from a glacier by melt-water streams

Pinery – a vegetative community within the historic pine range in Missouri that has shortleaf pine as a significant tree species

Prairie – a vegetative community dominated by perennial grasses and forbs with scattered shrubs and very few trees

Protected – steep, north- and east-facing slopes, which are cooler and moister than other slope aspects

Residuum - unconsolidated, weathered, or partly weathered mineral material that accumulates by disintegration of bedrock in place

Riser – a component of terraces and flood-plain steps consisting of the steep side slope; the escarpment

Riverfront – a vegetative community in the floodplain immediately adjacent and generally parallel to a river or stream channel

River hills – a geographic area characterized by thick, dissected loess deposits, formed immediately adjacent to the edges of the Missouri and Mississippi River floodplains

Sandy – a coarse-sized soil containing a large mixture of sand and gravels and a somewhat smaller proportion of silts and clays with excessive drainage

Sandstone – a sedimentary rock containing dominantly sand-size particles

Savanna – grasslands interspersed with open-grown scattered trees, groupings of trees, and shrubs

Shale – a sedimentary rock formed from clay, silty clay, or silty clay loam deposits and having the tendency to split into thin layers

Shallow – a site with bedrock within 20 inches of the surface

Shoulder – the slope profile position that forms the convex surface near the top of a hill slope; it comprises the transition zone from summit to backslope

Sinkhole – a closed, circular or elliptical depression, commonly funnel-shaped, characterized by subsurface drainage and formed either by dissolution of the surface of underlying bedrock or by collapse of underlying caves within bedrock

Summit – the top or highest area of a hillslope

Swale – shallow, closed depressions irregularly spaced across a floodplain or terrace with an irregularly undulating surface.

Swamp – an area of low, saturated ground, intermittently or permanently covered with water, and predominantly vegetated by shrubs and trees.

Talus – rock fragments of any size or shape (usually coarse and angular) derived from and lying at the base of a cliff or very steep rock slope.

Terrace – a step-like surface, bordering a valley floor that represents the former position of a flood plain

Till – dominantly unsorted and unstratified soil material deposited directly by a glacier

Ultic – soil that has a clay-dominated subsoil (argillic horizon) with low amounts of bases such as calcium, and were typically formed under woody vegetation

Upland – a general term for the higher ground of a region, in contrast with a low-lying, adjacent land such as a valley or floodplain

Wet – a somewhat poorly, poorly or very poorly drained site that has an oversupply of moisture during the growing season

Woodland – a highly variable vegetative community with a canopy of trees ranging from 30 to 100 percent closure with a sparse midstory and a dense ground flora of grasses, sedges and forbs

References

Nelson, Paul W. 2010. *The Terrestrial Natural Communities of Missouri*. Missouri Department of Conservation, Jefferson City, Missouri.

Nigh, Timothy A., and Walter A. Schroeder. 2002. *Atlas of Missouri Ecoregions*. Missouri Department of Conservation, Jefferson City, Missouri.

MDC, 2006. *Missouri Forest and Woodland Community Profiles*. Missouri Department of Conservation, Jefferson City, Missouri.