

Mapunit Name	HEL Class	HEL Water	HEL Wind
Badlands	null	null	null
Chistochina silt loam, 0 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Copper River peat, 0 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Copper River peat, 7 to 12 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Copper River peat, 12 to 20 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Copper River peat, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Kenny Lake silt loam, 2 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Kenny Lake silt loam, 7 to 12 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Kenny Lake silt loam, 12 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Chitina silt loam, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Chitina silt loam, 2 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Chitina silt loam, 7 to 12 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Chitina silt loam, 12 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina silt loam, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina silt loam, 2 to 7 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina silt loam, 7 to 12 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina silt loam, 12 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina-Hanagita complex, 2 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Copper River-Hanagita complex, 2 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Tonsina-Hanagita complex, 20 to 70 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Cryochrepts-Rock outcrop complex, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Cryochrepts-Rock outcrop complex, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Cryochrepts-Rock outcrop complex, 30 to 70 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Cryochrepts and Cryochrepts, 30 to 70 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Dagina peat, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Dagina-Kanalechechena complex, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Dagina-Tolsona complex, 0 to 5 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Pits, gravel	Not highly erodible land	Not highly erodible land	Not highly erodible land
Gulkana silt loam, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Gulkana silt loam, 2 to 7 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Gulkana silt loam, 7 to 12 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Gulkana silt loam, 12 to 20 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Klawasi peat, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Klawasi peat, 2 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Klawasi peat, 7 to 12 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Klawasi peat, cool, 0 to 7 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Klawasi peat, cool, 7 to 20 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Klawasi peat, depression, 0 to 2 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Gakona silt loam, cool, 0 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Gakona silt loam, 7 to 20 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Gakona silt loam, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Gakona silt loam, 2 to 7 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Gakona silt loam, 7 to 12 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Gakona silt loam, 12 to 20 percent slopes	Potentially highly erodible land	Potentially highly erodible land	Not highly erodible land
Klawasi-Tolsona complex, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Gakona-Stuck complex, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Gakona-Chetasilina complex, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Gakona-Chetasilina complex, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Klawasi-Wrangell complex, 0 to 2 percent slopes	Not highly erodible land	Not highly erodible land	Not highly erodible land
Klutina-Klutina, rarely flooded, complex, 0 to 2 percent slopes	Highly erodible land	Highly erodible land	Not highly erodible land
Klutina silt loam, rarely flooded, 2 to 7 percent slopes	Highly erodible land	Potentially highly erodible land	Highly erodible land

