

UNITED STATES DEPARTMENT OF AGRICULTURE  
NATURAL RESOURCES CONSERVATION SERVICE

CONSERVATION PRACTICE STANDARD

**FOREST STAND IMPROVEMENT**

(Acre)

CODE 666

**DEFINITION**

The manipulation of species composition, stand structure, and stocking by cutting or killing selected trees and understory vegetation.

**PURPOSE**

- Increase the quantity and quality of forest products by manipulating stand density and structure.
- Timely harvest of forest products
- Development of renewable energy systems.
- Initiate forest stand regeneration.
- Reduce wildfire hazard.
- Improve forest health reducing the potential of damage from pests and moisture stress.
- Restore natural plant communities.
- Achieve or maintain a desired native understory plant community for special forest products, grazing, and browsing.
- Improve aesthetic and recreation values.
- Improve wildlife habitat.
- Alter water yield.
- Increase carbon storage in selected trees.

**CONDITIONS WHERE PRACTICE APPLIES**

All forest land.

This standard is not applicable for Alley Cropping (311), Multi-story Cropping (379), Windbreak/Shelterbelt Establishment (operation and maintenance) (380) and Windbreak/Shelterbelt Renovation (650).

**CRITERIA**

**General Criteria Applicable to All Purposes**

The harvest-regeneration strategy will be identified for all planned forest improvement harvesting:

Uneven-aged management systems (e.g., single-tree selection, group selection, coppice selection)

Even-aged management (e.g., clear-cut, seed-tree, shelterwood, coppice)

The extent or size and orientation of treatment area(s) shall be identified as part of practice design.

Preferred tree and understory species are identified and retained to achieve all planned purposes.

Spacing, density, size class, number and amounts of trees and understory species to be retained will follow established guidelines for the intended purposes.

Stocking guidelines shall contain stocking in terms of basal area, spacing or trees per acre by species and size class distribution.

The method, felling direction and timing of tree cutting for harvesting shall protect site resources, e.g., residual trees, wetlands, cultural resources, improvements and utilities. Time tree cutting to avoid buildup of insect or disease populations. Felling direction must be compatible with trail layout as specified by Alabama NRCS conservation practice standard (AL NRCS CPS), [Forest Trails and Landings - Code 655](#). Forest stand improvement activities shall be performed to minimize soil erosion, compaction, rutting, and damage to remaining vegetation and maintain hydrologic conditions.

Refer to the AL NRCS CPS, [Access Road – Code 560](#), standard for roads associated with forest stand improvement activities.

Slash and debris will be treated such that they do not present an unacceptable fire, safety, environmental or pest hazard. Such remaining material will not interfere with the intended purpose or other management activities. Refer to AL NRCS CPS, Woody Residue Treatment - Code 384. Burning of slash and other debris on-site shall follow the AL NRCS CPS, [Prescribed Burning – Code 338](#).

#### **Additional Criteria to Develop Renewable Energy Systems**

Bioenergy intensity and frequency of energy biomass removals will be managed to prevent long-term negative impacts on the stand.

The harvesting of energy biomass shall be accomplished in a manner that will not compromise the other intended purpose(s) and functions. If applicable refer to State woody biomass Best Management Practices (BMPs).

#### **Additional Criteria to Reduce Wildfire Hazard**

Reduce stocking rates of trees to minimize crown-to-crown spread of fire.

Remove “ladder” fuels to minimize the occurrence of crown fires.

Further treat or eliminate slash accumulations next to roads and trails.

Reduce or eliminate species with high volatility but not to a level that would compromise other intended purposes.

For additional wildfire risk and damage reduction, refer to the AL NRCS CPS, [Fuel Break – Code 383](#), Firebreak - Code 394, and Alabama Job Sheet, [Reduce Wildfire Hazard, AL666](#).

#### **Additional Criteria to Improve Wildlife Habitat**

Manage for a variety of native tree species and stocking rates that meet desired wildlife and pollinator species food and cover requirements.

Create, recruit and maintain sufficient snags and down woody material to meet requirements of desired species and secondary cavity nesting species in balance with conditions needed to achieve other intended purposes.

Minimize improvement actions that disturb seasonal wildlife activities.

Refer to Early Successional Habitat Development/Management (647), Rare and Declining Habitats (643), Upland Wildlife Habitat Management (645), and Wetland Wildlife Habitat Management (644) to further develop and manage wildlife-related activities.

**NRCS, AL  
August 2011**

#### **Additional Criteria to Increase Carbon Storage in Selected Trees**

Manage for tree species and stocking rates that have higher rates of growth and potential for carbon sequestration.

#### **Additional Criteria for Thinning**

A. Trees to Leave: The selection of trees to leave in intermediate cuttings should be based on management objectives, adaptability of species to certain soils, the condition of individual trees as related to insects and diseases, and the form and vigor of individual trees. Species to favor on various soils are listed in the Woodland Reference #13-2, entitled "Considerations for Forest Management on Alabama Soils", in Section I of the Field Office Technical Guide. Merchantable dead trees and trees which are likely to die should be harvested during periodic thinnings.

B. Stocking: Stand density is a measure of the stocking of a stand of trees. Basal area is the common method used by foresters to measure stand density.

Stands should be thinned to the desired basal area or spacing according to the species and DBH.

<b>DESIRED STOCKING**</b>			
<b>HARDWOOD</b>			
<b>Avg. DBH Main Stand</b>	<b>Spacing</b>		<b># Trees per ac.</b>
	<b>ft.</b>	<b>BA</b>	
4	7-9	50-80	575-920
6	10-12	60-90	307-460
8	12-16	60-110	172-315
10	14-18	70-120	129-221
12	17-22	70-120	90-153
14	19-25	75-125	71-117
16	21-27	80-135	58-97
<b>PINE</b>			
4	8-10	40-60	460-690
6	10-12	60-80	307-409
8	12-14	80-105	230-301
10	14-17	85-115	156-212
12	17-20	90-120	115-153
14	19-22	95-130	89-122
16	22-25	95-130	69-94
** The smallest number is the "thin to" criteria under basal area (BA) and # of trees per ac. The largest number indicates when the stand needs to be thinned.			

C. Cutting Cycles: The time between commercial thinnings will vary according to site quality, the degree of thinning based on a diameter increase of 2 in. at breast height for pine and upland hardwoods and a diameter increase of 4 in. at breast height for the other hardwood species.

Site Index	CUTTING CYCLES (years)		
	Pine & Upland Oak	Bottomland Hardwood	Sweetgum & Yellow-Poplar
60	8	---	---
70	7	---	---
80	6	11	12
90	5	10	10
100	5	9	9
110	5	---	8

D. Multiple Use:

1. Grazing - Where grazing is a management objective the residual basal area should be approximately 60 sq. feet.

2. Wildlife - Mast bearing hardwoods should be retained where possible when wildlife is a management objective.

3. Aesthetics - Flowering trees and shrubs should be favored when aesthetics is a management objective.

4. Recreation - Intermediate thinnings should be timed in order not to conflict with recreational activities.

### Criteria for Harvest Cutting

A. Definition: Harvesting merchantable trees that are either financially or biologically mature.

B. Purpose: To harvest forest products and to ensure that the forest is regenerated for both soil protection and the production of wood products and other multiple uses. To create openings for regeneration, harvest trees should be removed in groups, strips, or blocks. Openings should be at least 100 ft. wide to permit sufficient sunlight. Sawtimber rotations are generally from 35 to 60 years, and pulpwood/chip-n-saw rotations are usually 25 to 30 years. To maintain a sustained yield of forest products, a percentage of the total forest acreage can be harvest cut and regenerated each cutting cycle. This percentage is calculated by dividing the cutting cycle by the number of years in the rotation. To maintain a sustained yield of forest products on large acreages (1000 acres or more), a percentage of the total forest acreage can be

clearcut and regenerated each year. This percentage is calculated by dividing the total acreage by the number of years in the rotation.

C. Pine:

1. Seed Tree Cut - Seed trees should be dominant trees of good quality that are at least 9 to 10 inches in diameter. Seed trees should be removed within 3 to 5 years to reduce damage to seedlings.

MINIMUM RECOMMENDED NUMBER OF SEED TREES (Per Acre)					
DBH =	9	10	12	14	16+
Shortleaf	---	20	14	12	12
Loblolly	---	12	9	6	4
Slash	---	12	9	6	4
Longleaf	---	55	38	28	21
Virginia	6	5	4	4	4
** Shelterwood cut of 30 sq. ft. of basal area.					

2. Shelterwood Cut - A shelterwood cut involves leaving a large number of seed trees per acre. The trees are thinned to approximately 30 to 60 sq. ft. of basal area depending on species. The residual stand of trees should be removed within 3 to 5 years to reduce damage to seedlings.

3. Clearcutting - All merchantable trees are removed. Site preparation is usually needed to enhance regeneration. See Alabama NRCS Conservation Practice Standard, Code 490, Forest Site Preparation. Natural regeneration methods such as seed-in-place, seedlings in place and seeding from the side may be used with clearcuts.

Tree planting and direct seeding are artificial regeneration options. Improved seedlings and improved seed should be used when possible. See Alabama NRCS Conservation Practice Standard, Code 612, Tree/Shrub Planting.

4. Individual Tree Selection - The removal of trees individually or in small clumps. This type of harvesting creates an uneven-aged stand of timber. The stand is regulated by periodic volume removal. Regeneration occurs either continuously or periodically.

D. Hardwood:

1. Clearcutting - Clearcutting is one method of regenerating bottomland hardwoods. All merchantable trees should be harvested and small

trees should be removed either mechanically or by the use of herbicides.

Yellow-poplar is a species which is easily managed by clearcutting if a seed source has been present for 3 to 5 years. Yellow-poplar seeds remain viable in the forest litter for up to 8 years. Normal harvesting operations will expose mineral soil and allow adequate germination of yellow-poplar seeds.

2. Shelterwood Cut - The shelterwood system can be used to obtain oak regeneration. The shelterwood system is preferable from a wildlife perspective. Stand density is reduced to 60% stocking to allow seedling establishment. When a stand of 500 or more seedlings per acre is 4.5 ft. or higher, the overstory stand should be removed. It may take as long as 15 to 20 years to obtain the desired reproduction once stand density is reduced.

E. Multiple Uses:

1. Grazing - Areas to be regenerated should have a tree spacing of 8'x10' or 8'x12' where forest grazing is a management objective.

2. Wildlife - Mast bearing hardwoods should be retained where possible when wildlife is a management objective. Openings should be retained or developed as needed. See AL NRCS CPS, [Upland Wildlife Habitat Management – Code 645](#).

3. Aesthetics - Clearcuts and harvested areas should be irregular in shape where practical and flowering trees and shrubs should be favored after regeneration.

4. Recreation - Harvesting should be timed in areas used for recreation in order to avoid conflicts with recreational activities.

F. Release of Established Seedlings: Competing vegetation may be controlled by one or more of the methods listed in **Criteria for Timber Stand Improvement**, C. Methods of Removal or Control. If necessary, seedlings should be released from overtopping trees and shrubs no later than the end of the second growing season.

G. Protection of Young Trees: Livestock should be excluded from stands of desirable hardwoods and overgrazing should be prevented on areas planted to pine. Control of wildfire is necessary throughout the life of all stands of trees.

## **Criteria for Timber Stand Improvement**

A. Trees to Release: Trees to release are determined by the owner's objectives, by species adaptation and desirability, and by the form and vigor of individual trees. Favor the most vigorous and best formed trees of desired species. See Woodland Reference #13-2 entitled "[Considerations for Forest Management on Alabama Soils](#)" in Section I of the Field Office Technical Guide, for species to favor on various soils.

B. Spacing and Number of Release Trees: For pine trees less than 4 inches in diameter, released trees should average no farther apart than 12 ft. or about 300 trees per acre. Release pine trees 4 to 8 inches in diameter at intervals of 12 to 16 ft. or about 200 to 300 trees per acre. Release hardwood trees 4 to 8 in. in diameter at intervals of 14 to 19 ft. or about 120 to 220 trees per acre. If the minimum number of desired trees is not present, either replant or interplant according to specifications in Alabama NRCS Conservation Practice Standard, Code 612, Tree/Shrub Planting.

C. Methods of Removal or Control: Either remove or control unwanted hardwood trees, shrubs, vines, and weeds by either chemical treatment, mechanical treatment, or prescribed burning.

Factors to consider when choosing a method are:

1) species and size of trees and other vegetation to be killed, 2) presence of desired trees, 3) density of unwanted trees and other plants, 4) site quality, 5) size and location of the area to be treated, and 6) costs of equipment and materials.

1. Mechanical treatment - Trees can be girdled with either an axe or a mechanical tree girdler. Trees smaller than 12 inches in diameter may sprout, and such trees should be treated with a herbicide.

2. Cut and fell - Larger trees may be removed by the cut and fell method. Trees should be utilized for firewood where feasible. Tree stumps 14 inches in diameter or less should be treated with herbicides to prevent sprouting.

3. Tree injection - Herbicides are applied to individual stems at the rate of 1 to 2 ml per in. of stem diameter.

4. Spraying - Both high volume and low volume sprays may be used to kill undesirable woody plants. Low volume sprays are usually applied by aircraft.

5. Soil treatment - Certain herbicides may be applied to the soil to control undesirable plants. Application rates will vary according to soil types and plant species.

6. Prescribed burning - Prescribed burning can be used to control undesirable hardwoods usually 2 in. or less in diameter. For specifications, see AL NRCS CPS, [Prescribed Burning- Code 338](#).

Caution: If not properly handled and applied, herbicides can be injurious to people, domestic animals, beneficial insects, desirable plants, and to fish and other wildlife. Use herbicides selectively and carefully. Follow recommendations when disposing of surplus herbicides and herbicide containers.

### **Criteria for Precommercial Thinning**

Precommercial thinning is needed: 1) where stands of desirable trees are of unmerchantable size and are overstocked, thereby preventing satisfactory growth; and 2) where thinning will either increase the growth of remaining trees or enhance the beauty of the stand. Precommercial thinnings are usually applied to seedling and sapling-sized stands (stands with trees less than 5 in. in diameter).

A. Species to Favor: Favor trees according to desirable species, form, vigor, and crown development. See Woodland Reference #13-2, entitled "[Consideration for Forestry Management on Alabama Soils](#)", in Section I of the Field Office Technical Guide for species to favor on various soils.

#### 1. Selecting Trees to Leave -

a. Seedling-Sized Stands - In seedling-sized stands where all trees are about the same size, it is usually best to leave clumps of seedlings 6 to 10 ft. apart.

b. Sapling-Sized Stands - In sapling-sized stands, release the best trees at 10 to 12 ft. intervals.

2. Removing Trees - Remove trees by one or more of the following methods:

a. Mowing Machine: Where trees are generally less than 2 in. dbh, mow strips 6 to 10 ft. wide and either cross-check or cross-mow strips of the same width. Remaining clumps should be 2 ft. wide.

b. Bush and Bog Harrow, Chopper, or Bulldozer Strips - Disk trees in the seedling to small sapling size with a bush and bog harrow or knock

them down with either a bulldozer or chopper. Remove in strips of the same width and cross-check in the same manner as for mowing. To prevent root damage, strips that remain should be 2 ft. wide.

c. Hand Tools - Seedlings and saplings may be thinned with machetes, axes, bush hooks, hoes, or with power equipment such as chain saws. Seedlings up to 20 in. tall should be cut near the ground level. Seedlings and saplings taller than 20 in. may be cut at waist height if they are cut below living branches and stem needles.

### 3. Optimum Seasons for Thinning

a. Pine Types - In pine stands, precommercial thinnings should be done from October through March. Thinnings at that time reduce the possibility of insect infestation.

b. Hardwood Types - Precommercial thinnings should be done in hardwood stands during midsummer because there is less sprouting then than during other seasons.

c. Pine-Hardwood Types - Precommercial thinnings should be done in pine-hardwood stands in late fall and winter. Sprouting of hardwoods may be controlled by herbicides.

4. Slash Disposal - To reduce the possibility of loss from insects and fire, leave no trees cut in precommercial thinnings either leaning against remaining trees or touching them. In high fire-risk areas, cut trees should be removed.

## **CONSIDERATIONS**

Silvicultural objectives and harvest-regeneration strategies may change over time and may be limited by prior management.

Successful regeneration of desirable species is usually dependent upon timely application of forest stand improvement and other practices, e.g., prescribed burning, site preparation, tree and shrub establishment, prescribed grazing and access control.

The extent, timing, size of treatment area, or the intensity of the practice should be adjusted to minimize cumulative effects (onsite and offsite), e.g., hydrologic and stream alteration, habitat fragmentation, nutrient cycling, biodiversity and visual resources.

For purposes other than improving wildlife habitat, the practice should be timed to minimize disturbance of seasonal pollinator and wildlife activities.

Landowners should secure a written contract with any service provider that specifically describes the extent of activity, duration of activity, liability and responsibilities of each party and amount and timing of payments for services provided

Slash, debris and other vegetation (biomass) removed during stand improvement may be used to produce energy. Management alternatives should consider the amount of energy required to produce and convert the biomass into energy with the amount produced by the biomass. Wildlife and sustainability requirements should also be considered.

Invasive or noxious woody vegetation should be controlled.

Clients should be advised of responsibilities of wildfire control and consider the development of a wildfire control plan including "defensible" space, access routes, fire-season water source, and location of wildfire control facilities.

Timing of treatment and retaining dead or dying trees will minimize impacts on wildlife.

Wildlife food and cover can be retained by minimizing modifications to composition and spacing regardless of the purpose for treatment. Forested wildlife corridors can minimize fragmentation effects.

Select appropriate tree removal techniques that would reduce the potential for erosion and sedimentation.

Erosion rates and sediment yields may increase as a result of harvesting activities.

Proper selection and application of pesticides should be considered to minimize surface water transport and potential leaching to ground water.

## PLANS AND SPECIFICATIONS

Specifications for applying this practice shall be prepared for each site and recorded using approved specification sheets, job sheets, narrative statements in the conservation plan, or other acceptable documentation.

## OPERATION AND MAINTENANCE

Periodic inspections during and after treatment activities are necessary to ensure that purposes are achieved and resource damage is minimized, e.g., assessment of insects, disease and other pests, storm damage, and damage by trespass. The results of inspections shall determine the need for additional treatment under this practice.

## REFERENCES

Alabama Cooperative Extension System. Auburn University and Alabama A&M University. October 1988. Herbaceous Weed Control in Young Pine Plantations. Circular ANR-518.

Alabama Cooperative Extension System, Auburn University and Alabama A&M University. Loblolly Pine Management. ANR-490 Prepared by: Frank A. Roth II.

Alabama Cooperative Extension System, Auburn University and Alabama A&M University. Management of Hardwood Forests For Timber in Alabama. ANR-581. Prepared by: Robert Jones.

American Pulpwood Association. 1988. Regenerating and Managing Natural Stands of Bottomland Hardwoods. APA 88-A-6. Prepared by: R.C. Kellison, J.P. Martin, G.D. Hansen, and R. Lea.

Cantrell, Rick. November 1985. A Guide to Silvicultural Herbicide Use in the Southern United States. Auburn University.

Holt, Harvey and B. Fischer, ed. 1979. Regenerating Oaks in Upland Hardwood Forests. Purdue University Research Foundation.

U.S. Department of Agriculture, Forest Service. 1985. Thinning Practices in Southern Pines - With Pest Management Recommendations. Technical Bulletin 1703. Prepared by: T.E. Nebeker, J.D. Hodges, B.K. Karr, and D.M. Moehrin

U.S. Department of Agriculture, Forest Service, Southern Region. 1985. Proceedings - Third Symposium on Southeastern Hardwoods.

USDA Forest Service Miscellaneous Report, July 1986. Service Foresters Handbook. SA-MR-10.