United States Department of Agriculture
Natural Resources Conservation Service
Ecological Site Description

Section I: Ecological Site Characteristics

Ecological Site Identification and Concept

Site stage: **Provisional**

Provisional: an ESD at the provisional status represents the lowest tier of documentation that is releasable to the public. It contains a grouping of soil units that respond similarly to ecological processes. The ESD contains 1) enough information to distinguish it from similar and associated ecological sites and 2) a draft state and transition model capturing the ecological processes and vegetative states and community phases as they are currently conceptualized. The provisional ESD has undergone both quality control and quality assurance protocols. It is expected that the provisional ESD will continue refinement towards an approved status.

Site name: **Semidesert Juniper Loam**
Juniperus osteosperma - *Pinus edulis* / *Pleuraphis jamesii*
(Utah juniper - twoneedle pinyon / James' galleta)
Site type: Rangeland
Site ID: R036XY113CO
Major land resource area (MLRA): 036-Southwestern Plateaus, Mesas, and Foothills



Location Map

Semidesert Juniper Loam ecological site is found on dipslopes, dipslopes on cuestas, cuestas, and eroded dipslopes on cuestas in MLRA 36 (Southwestern Plateaus Mesas and Foothills). The MLRA 36 is illustrated orange color on the map. The ecological site locations as assigned in soil survey map units are shown in pink color.

The site concept was established within the MLRA 36 semi-desert. This zone is 9 to 12 inches of precipitation and has a mesic temperature regime. This site has bimodal precipitation that is dominated by Utah Juniper.

Ecological Site Concept

The 36X Semidesert Juniper Loam ecological site was drafted from the existing Semidesert Juniper Loam Range Site 34B and 35X (NRCS, March, 1996). This site was written prior to MLRA 36 being mapped in Colorado and this area was in MLRA 34X when it was written. This site occurs on dipslopes, dipslopes on cuestas, cuestas, and eroded dipslopes on cuestas on loamy textured soils (sandy loam, fine sandy loam and loam) derived from slope alluvium over residuum weathered from sandstone, slope alluvium derived from sandstone, and slope alluvium derived from sandstone and shale over residuum weathered from sandstone. It is a Utah Juniper-Pinyon. It has an ustic aridic moisture regime and mesic temperature regime. The effective precipitation ranges from 9 to 12 inches.

Physiographic Features

This site occurs on dipslopes, dipslopes on cuestas, cuestas, and eroded dipslopes on cuestas. Slopes typically range from 3-30%, and elevations are generally 5400-6800 ft., but it can go down to 5200 on north and easterly slopes and up to 7000 on south and westerly slopes.

Landform: (1) Dipslope
(2) Cuesta
(3) Structural bench

	Minimum	Maximum
Elevation (feet):	5400	6800
Slope (percent):	3	30
Flooding		
Frequency:	None	None
Ponding		
Frequency:	None	None
Runoff class:	High	Very high
Aspect:	No Influence on this site	

Climatic Features

Average annual precipitation is about 9 to 12 inches. This area is located where there is winter precipitation and summer monsoonal rains meet. Of this, 45-50% falls as snow, and 50-55% falls as rain. Snow usually falls from November to March. Rains are falls April 1 thru October 31. The driest period is usually May to June. Plant growth begins late March and early April. Cool-season plants start a dormancy period during June. Summer thundershowers are common in July to September. The summer moisture will favor growth from the warm season plants. When late summer and fall rains occur, warm-season plants accelerate growth, and some regrowth occurs on cool-season species. Shrub species continue growth through the entire growing season. The average annual total snowfall is 17.8 inches. The highest winter snowfall record in this area is 44.8 inches which occurred in 1972-1973. The lowest snowfall record is zero inches during the 1999-2000 winter. The highest yearly precipitation recorded was 19.02 in 2015 and the lowest was 5.17 in 1989. Mean daily annual air temperature is about 50ºF to 54ºF, averaging about 33ºF for the winter and 61ºF through the growing season, March through October. Summer temperatures of 100ºF or more are not unusual. The frost-free period typically ranges from 125 to 165 days at Hovenweep NM (national monument). The last spring frost is the first part of May to the end of May. The first fall frost is the end of September to the middle of October. Mean annual temperature ranges from 55 to 49ºF. Average annual temperature is 51.9ºF. The coldest winter temperature recorded was -24ºF on December 24, 1990 and the coldest summer temperature recorded was 26ºF on June 12, 1970. The hottest day on record is 106 ºF on July 15, 1998. Wide yearly and seasonal fluctuations are common for this climatic zone. Data taken from Western Regional Climate Center (2017) for Hovenweep NM, Utah Climate Station. Hovenweep NM is on the Western edge of the MLRA. Hovenweep NM is the only station occurring in the MLRA in this zone. It is on the upper end of precipitation. There is a need for climate date in the zone. Typical warm dry weather in late spring and early summer puts warm season plants at a distinct disadvantage compared to plants that grow earlier on stored winter moisture and spring rains. Deep storage of winter moisture allows sagebrush and small trees to compete strongly with shallower rooted grasses during this dry period where other factors encourage their spread.

	Averaged
Frost-free period (days):	131
Freeze-free period (days):	146
Mean annual precipitation (inches):	11.24

Monthly Precipitation (Inches):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
High	1.31	1.36	1.70	1.43	0.93	0.41	1.41	1.98	1.77	1.63	1.95	1.26
Medium	0.65	0.67	0.59	0.55	0.51	0.22	0.83	1.06	0.98	1.14	0.88	0.76
Low	0.40	0.43	0.30	0.25	0.14	0.03	0.32	0.37	0.68	0.52	0.35	0.34



Monthly Temperature (°F):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
High	41.6	47.6	57.4	65.9	76.8	88.0	93.4	90.3	81.7	68.3	53.4	41.8
Low	18.0	23.4	29.2	34.8	44.1	52.6	60.9	59.6	49.9	37.5	26.8	18.5

30 Year Annual Rainfall (inches):

1981 N	1982 H	1983 H	1984 N	1985 H	1986 H	1987 H	1988 N	1989 D	1990 N	1991 N	1992 H	1993 N	1994 N	1995 N
10.46	14.59	15.13	10.84	14.41	15.04	15.99	10.24	5.19	10.44	9.68	14.12	13.8	9.83	11.17
1996 N	1997 H	1998 N	1999 D	2000 N	2001 N	2002 D	2003 N	2004 N	2005 N	2006 N	2007 N	2008 N	2009 D	2010 N
12.59	14.67	9	8.05	8.82	10.18	6.48	10.65	10.55	12	10.35	11.33	11	7.1	13.46

D-Drought N-Normal H-Heavy



Climate stations: (1) HOVENWEEP NM [USC00424100], San Juan County UT 84534. Period of record 1981-2010

Influencing Water Features

None.

Wetland Description (Cowardin System)		
System	Subsystem	Class
None	N/A	N/A

Representative Soil Features

Soils are very shallow to shallow in depth (5-20 inches). The surface soils of this site are sandy loams, fine sandy loams, and loams. The surface layer texture is usually a sandy loam with 15-20% clay. The subsoils are loamy. The subsurface can be sandy loam, sandy clay loam, and loam with approximately 19-30% clay. The most common parent materials are slope alluvium over residuum weathered from sandstone, slope alluvium derived from sandstone, and slope alluvium derived from sandstone and shale over residuum weathered from sandstone.

Soils assigned to this site and these soil map units needing to be evaluated for which ESD (ecological site description) they belong to are: Rizno, Dolcan, Romberg and Wauquie.

This ecological site has been used in the following Soil Surveys: CO670 (Ute Mountain Area), and CO677 (Ridgeway Area), CO680 (Mesa County Area), CO679 (Paonia Area), CO676 (Uncompahgre Area).

Typical soils assigned to this ecological site are:

Loamy - Barboncito, Lazear, Rizno, Dolcan

Loamy-Skeletal – Walknolls, Reef

Parent materials

Kind: Slope alluvium, Residuum

Origin: Sandstone, Sandstone and shale

Surface texture: (1) Sandy loam
(2) Fine sandy loam
(3) Loam

Subsurface texture group: Loamy

	Minimum	Maximum
Surface fragments <=3" (% cover):	0	10
Surface fragments >3" (% cover):	0	5
Subsurface fragments <=3" (% volume):	0	10
Subsurface fragments >3" (% volume):	0	5

Drainage class: Well drained

	Minimum	Maximum
Depth (inches):	5	20
Available water capacity (inches):	0.70	2.30
Electrical conductivity (mmhos/cm):	0	2
Sodium adsorption ratio:	0	0
Calcium carbonate equivalent (percent):	0	14
Soil reaction (1:1 water):	7.4	8.2

Plant Communities

Ecological Dynamics of the Site

This area has a long history of past prehistoric human use for thousands of years. They used pinyon-juniper woodlands for hunting, fuelwood, for food, such as pinon nuts. MLRA 36 have archaeological evidence indicating pinyon-juniper woodlands where modified by prehistoric humans and not pristine and thus where altered at the time of European settlement (Cartledge & Propper, 1993). This area is characterize by broken topography, and lack of perennial water sources. Most pinyon-juniper Northern half of MLRA 36 (Colorado and Utah) can be describe as a persistent woodland type. There is a winter-summer bimodal precipitation pattern on the Colorado Plateau. Meaning that this site developed under climatic conditions that include wet, cold winters, and hot, dry summers with summer rains. This area so included natural influences of herbivory, fire, and climate. This area rarely served as habitat for large herds of native herbivores or large frequent historic fires due to the broken topography. The precipitation and climate of MLRA 36 are conducive to producing Pinyon/juniper, and sagebrush complexes.

Pinyon-Juniper expansion began during the late 1800s into deeper well drained soils. (Tausch et al. 1981, Miller and Tausch, 2001). The causes of woodland expansion are often attributed to an reduction in fires, introduction of livestock grazing, shifts in climate, and increases in

atmospheric CO2 (Miller and Rose 1999). Prior to European settlement, PJ woodland species were primarily found on shallow soils and rocky ridges. Few fire history studies and pinyon-juniper chronologies have been done in the southwest. It appears that woodland on the Colorado Plateau are more susceptible to die off from severe drought (Miller and Tausch, 2001). Historically, fires before European settlement in the southwest occurred late spring to mid-summer (Miller and Tausch, 2001).

Historic fire return intervals (300-1000 years) are long, possibly indicating that fire did not play a frequent role in community dynamics. Pinyon and Juniper communities near Mesa Verde were established before European settlement with a fire return interval approximately 400 years (Floyd et al., 2000). Shinneman and Baker (2009) estimated the FRI on the Uncompahgre Plateau to be 400 to 600 years. Mesa Verde (Floyd et al., 2000) and Uncompahgre (Shinneman and Baker, 2009) are in the foothills/upland zone (12 to 16 inches annual precipitation) in MLRA 36. One other known study in the Colorado National Monument on the north eastern part of the Uncompahgre Plateau suggest that lower ecological site zone (semi-desert) (9 to 12 inches of annual precipitation) have a fire return interval of 300 to 1,000 years (Kennard and Moore, 2013). One other difference is that in the semi-desert zone smaller fire of only a few trees maybe more common than the infrequent larger fires found in other studies.

In lower elevations and lower precipitation areas, Utah Juniper maybe dominant over Pinyon. As the precipitation increase and effect moisture increase so will pinyon. The lower end of the pinyon-juniper woodland would be almost entirely Utah Juniper with the reverse happening and pinyon being dominant in the upper end of the pinyon-juniper belt.

The driving factors in Pinyon Juniper woodlands seem to be weather patterns. Drought and insects outbreaks appear to be the main driving factors for mortality in many of the Pinyon/Juniper communities. (Shinneman and Baker, 2009, Floyd et al., 2004) Wet periods seem to enhance and promote pinyon and juniper establishment. Betancourt (1993), noted that Pinyon and Juniper woodlands in the southwest appear to be more susceptible to large die offs during droughts, than in other locations. As severe droughts persist, the Pinyon trees, being more susceptible to drought and insects, seem to die out, while the Utah juniper trees survive. This action could open the canopy for a few years and with sufficient moisture, grasses and forbs would be expected to respond favorably. Two studies illustrated this on the Uncompahgre Plateau found that pinyon began increasing in the 1700s, during a wet period that followed a long dry period. So, tree infill and expansion began before European settlement. Associated fire reduction and livestock grazing effect of European settlers can after the trees started the current expansion. Since the 1900s there has been 2 very wet period in the southwest, during 1900s to 1920s and 1970s to 1990s. These periods saw an increase in Pinon establishment. During the drought of the 1950s and the drought mid-1990s to early 2000s, Pinyon mortality was extensive. (Romme, et al. 2009)

Disturbances such as improper grazing (continuous season long grazing, heavy stocking rates, etc.), recreation activities, etc., can remove herbaceous vegetation and compact the soils. The unpredictability of the annual growing conditions make these communities susceptible to the loss of understory and the resulting accelerated erosion. This ecological site has been grazed by domestic livestock since they were introduced into the area, though grazing has been light due to the lack of water and difficult terrain. The introduction of domestic livestock and the use of fencing and reliable water sources have influenced the disturbance regime of this site. As of this date, invasive annual grasslands that are so common in the Great Basin after a severe disturbance are not as prevalent in MLRA 36, potentially due to the remote location, the climate, and/or the soils.

PJ fire intervals can be influences by the landscape it occurs on. PJ that is complexed with sagebrush site would burn more frequently do to the fine fuels in the sagebrush sites to start the fires. So, the more rough broken terrain would burn less frequently than the gentler and broader landscapes. PJ sites on the Colorado Plateau generally don't have enough fine fuels to start large scale fires. The exception would be several wet years in a row that would create the fine fuels necessary for a fire to start.

As vegetation communities respond to changes in management or natural occurrences, thresholds can be crossed, which usually means that a return to the previous state may not be possible without major energy inputs. The amount of energy input needed to affect vegetative shifts depends on the present biotic and abiotic features and the desired results.

Pinyon-juniper sites were treated as one vegetation dynamic type when developing the provision ecological site initiative for MLRA 36. These sites will need to be altered as more data and knowledge in the future becomes available. Variability in climate, soils, aspect and complex biological processes will cause the plant communities to differ. These factors contributing to annual production variability include wildlife use, drought, and insects. Factors contributing to special variability include soil texture, depth, rock fragments, slope, aspect, and micro-topography. The species lists are representative and not a complete list of all occurring or potentially occurring species on this site. The species lists are not intended to cover the full range of conditions, species and responses of the site. The State & Transition model depicted for this site is based on available research, field observations and interpretations by experts and could change as knowledge increases. As more data is collected, some of these plant communities may be revised or removed, and new ones may be added. The following diagram does not necessarily depict all the transitions and states that this site may exhibit, but it does show some of the most common plant communities.

State-and-Transition Diagram



STM

Legend



Legend

State 1: Reference State

This state represents the natural variability and dynamics of this site that occurred naturally. This state includes the dominant biotic communities that would have occurred on this ecological site prior to European Settlement. The dominant aspect of this site is Pinyon and Utah Juniper with an understory of shrubs and associated grasses. Fluctuations in species compositions and relative production may change from year to year dependent upon abnormal precipitation or other climatic factors. The primary disturbance mechanisms for this site in reference condition include drought, insects, and infrequent fire. Because catastrophic disturbances like a crown fire or drought happen with long intervals, these communities have long periods of succession, (i.e. long periods of dense Pinyon and Juniper)—300-600 years in upland/foothills ecological site zone and 300 to 1,000 in semi-desert ecological site zone. According to Shinneman (2006), the pinon-juniper zone on the Uncompahgre Plateau typically burns in high-intensity, stand-replacing fires with a 400–600 years rotation (Shinneman, 2006). In the semi-arid environment of this ecological site, fine fuels are typically not continuous, reducing the likelihood of short fire return intervals. Typically, fires occurred in late spring through mid-summer following several wet years that allowed the fine fuels to become more contiguous (Baisan and Swetnam, 1990, and Swetnam and Baisan, 1996). The higher in elevation and higher precipitation area would burn more frequently as they would have more fine fuels in the understory. The timing of drought, and fire, coupled with surface disturbance can dictate whether the community can stay within the reference state or if the community transitions into another state.

Community Phase 1.1: Pinyon-Juniper Woodland

A well-developed understory with a canopy of younger Pinyon and Utah juniper. At this stage Utah juniper may be dominant over Pinyon. Pinyon trees are more susceptible to drought, insects, and disease than Utah Juniper trees. In fact, it is difficult to identify methods beside fire that naturally reduce Utah juniper. After long periods of drought weaken the Pinyon trees, beetle kills can become quite extensive, especially after the droughts. Drought periods can also weaken and reduce the understory. Plant establishment is mainly limited by the available moisture. Biological crusts can be highly developed and diversified in the large interspaces between trees.

Community Phase Pathway 1.1A
This pathway occurs when events create a wetter climate cycle, favor pinyon and perennial bunch grass establishment. Following several favorable precipitation years and lack of surface disturbances, native perennial plants will reestablish.

Community Phase Pathway 1.1B
This pathway is very unlikely, but can occur when a fire is able to move through the community. Two situations can make this occur: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 1.2: Mature Pinyon-Juniper Woodland

Mature pinyon and Utah juniper woodland characterized this community phase. When weather patterns favor an increase of pinyon and Utah juniper canopy with the associated understory of shrubs, grasses and forbs. Depending on the timing of precipitation, cool season grasses, like Indian ricegrass or warm season grasses like galleta could be dominant. Interspaces supporting highly developed biological crusts are common.

Community Phase Pathway 1.2A
This pathway occurs during and after events such as drought or insect/pathogen outbreaks. Droughts and insects can kill the trees, increasing nutrient availability in the system. Due to the natural conditions of drought, grasses typically do not take up the extra nutrients in the long term. In the short term, grasses and forbs may increase for a few years until juniper and pinyon recover.

Community Phase Pathway 1.2B
This pathway is very unlikely but can occur when a fire is able to move through the community phase. Two situations can make this occur: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 1.3: Perennial Shrubland with scattered PJ

The overall aspect of this community phase is grasses and shrubs with scattered Utah juniper and maybe a few pinyon. The herbaceous understory has a mix of grasses and forbs. This community phase is a result of a crown fire or sufficiently large and hot ground fire that will kill many of the trees, combined with sufficient seed-banks and moisture for reestablishment of grasses and forbs. It is common that after a crown fire many patches of trees will remain unburned, because of fire's unpredictability and broken topography. This leaves a seed bank for the burned areas. This community phase is very short lived in comparison to the other community phases in this state.

The following is from the 1996 Range Site:

Tree canopy 15 to 30% - Ground cover and structure:

% Canopy cover vertical view

Grasses 20
Forbs 2

Shrubs 10
Trees 5

Average height (FT)

Grasses 1.0
Forbs 0.5
Shrubs 1.5
Trees 8.0

%Basal Cover

Grasses 8
Forbs 1
Shrubs 4
Trees 2

Total annual production:
In an average year, the approximate total annual production (air-dry) is as follows:
Tree canopy cover 0 to 15% 200 to 300 lbs/Ac.

Community Phase Pathway 1.3A
This pathway occurs when the climate favors the establishment and growth of trees. More energy is taken-up and stored in the trees as the length between fires and droughts increase. In addition, when shrubs establish on the site they can provide safe-sites for tree establishment furthering the presence of trees.

Perennial Shrubland with scattered PJ Plant Species Composition

Grass/Grasslike

Group	Group name	Common name	Symbol	Scientific name	Annual Production (pounds per acre) Low	Annual Production (pounds per acre) High	Foliar cover (percent) Low	Foliar cover (percent) High
1					75	200		
		Indian ricegrass	ACHY	*Achnatherum hymenoides*	25	45		
		Fendler threeawn	ARPUL	*Aristida purpurea var. longiseta*	0	15		
		blue grama	BOGR2	*Bouteloua gracilis*	0	15		
		squirreltail	ELEL5	*Elymus elymoides*	10	30		
		needle and thread	HECO26	*Hesperostipa comata*	25	45		
		prairie Junegrass	KOMA	*Koeleria macrantha*	0	15		
		James' galleta	PLJA	*Pleuraphis jamesii*	25	60		

Forb

Group	Group name	Common name	Symbol	Scientific name	Annual Production (pounds per acre) Low	Annual Production (pounds per acre) High	Foliar cover (percent) Low	Foliar cover (percent) High
2					25	65		
		rosy pussytoes	ANRO2	*Antennaria rosea*	0	15		
		Douglas' dustymaiden	CHDO	*Chaenactis douglasii*	0	5		
		hairy false goldenaster	HEVI4	*Heterotheca villosa*	0	10		
		hoary tansyaster	MACAC3	*Machaeranthera canescens var. canescens*	0	5		
		Colorado four o'clock	MIMU	*Mirabilis multiflora*	0	5		
		spiny phlox	PHHO	*Phlox hoodii*	0	10		
		threadleaf ragwort	SEFLF	*Senecio flaccidus var. flaccidus*	0	5		
		scarlet globemallow	SPCO	*Sphaeralcea coccinea*	1	10		
		stemless mock goldenweed	STAC	*Stenotus acaulis*	0	5		
		desert princesplume	STPI	*Stanleya pinnata*	0	3		

Shrub/Vine

Group	Group name	Common name	Symbol	Scientific name	Annual Production (pounds per acre) Low	Annual Production (pounds per acre) High	Foliar cover (percent) Low	Foliar cover (percent) High
3					50	150		
		black sagebrush	ARNO4	*Artemisia nova*	0	15		
		Wyoming big sagebrush	ARTRW8	*Artemisia tridentata subsp. wyomingensis*	10	30		
		shadscale saltbush	ATCO	*Atriplex confertifolia*	10	30		
		littleleaf mountain mahogany	CEIN7	*Cercocarpus intricatus*	0	5		
		longflower rabbitbrush	CHDE2	*Chrysothamnus depressus*	0	10		
		yellow rabbitbrush	CHVI8	*Chrysothamnus viscidiflorus*	0	15		
		mormon tea	EPVI	*Ephedra viridis*	0	10		
		spreading buckwheat	EREF	*Eriogonum effusum*	0	10		
		broom snakeweed	GUSA2	*Gutierrezia sarothrae*	0	15		
		winterfat	KRLA2	*Krascheninnikovia lanata*	2	15		
		plains pricklypear	OPPO	*Opuntia polyacantha*	0	15		
		soapweed yucca	YUGL	*Yucca glauca*	0	5		

Tree

Group	Group name	Common name	Symbol	Scientific name	Annual Production (pounds per acre) Low	Annual Production (pounds per acre) High	Foliar cover (percent) Low	Foliar cover (percent) High

4				10	70
	Utah juniper	JUOS	*Juniperus osteosperma*	10	60
	twoneedle pinyon	PIED	*Pinus edulis*	0	15

Annual Production by Plant Type

	Annual Production (lbs/ac)		
Plant type	Low	Representative value	High
Grass/Grasslike	105	120	135
Forb	25	40	55
Shrub/Vine	40	50	60
Tree	30	40	50
Total	200	250	300

Plant Growth Curve

Growth curve number: CO0102

Growth curve name: MLRA 34B & 36 - Semidesert Sites

Growth curve description: MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Transition T1A

This transition from the native perennial bunchgrass and shrub understory in the reference state to a state that has been invaded by naturalized species such as crested wheatgrass (blown in or seeded), cheatgrass, annual wheatgrass and other introduced or exotic plants. This transition occurs as natural and/or management actions favor an increase in non-native grasses and forbs, especially annuals. Possible events include the presence of invasive species, improper livestock grazing, extended droughts, and fire combined with an available seed source of non-native species.

State 2: Current Potential State

This state is very similar to the reference state, except that non-native grasses and/or forbs are now present in all community phases. The current potential state may include introduced (seeded) or invasive nonnative species. The invasive plants are present in sparse amounts in this state. Natural disturbance are still drought, insects, and infrequent fires still influence the community shifts. The human caused disturbance drivers (i.e. domestic livestock grazing, vegetation manipulation, and recreational activities (i.e. OHV use)) are now present. This shift in species composition could affect nutrient cycling, hydrology and soil stability. At this time there is no known way to effectively remove the non-native plants from the site once they have become established. State 2 is in jeopardy of moving to State 3 (Pinyon-Juniper Invasive State) when remaining native understory plants are stressed and invasive species have increased till they are dominant.

Community Phase 2.1: Pinyon-Juniper Woodland

A well-developed understory with a canopy of younger Pinyon and Utah juniper. At this stage Utah juniper may be dominant over Pinyon. Pinyon trees are more susceptible to drought, insects, and disease than Utah Juniper trees. In fact, it is difficult to identify methods beside fire that naturally reduce Utah juniper. After long periods of drought weaken the Pinyon trees, beetle kills can become quite extensive, especially after the droughts. Drought periods can also weaken and reduce the understory. Plant establishment is mainly limited by the available moisture. Biological crusts can be highly developed and diversified in the large interspaces between trees. Sparse invasive introduced plants species would be present in this phase.

Community Phase Pathway 2.1A
This pathway occurs when events create a wetter climate cycle, favor Pinyon and perennial bunch grass establishment. Following several favorable precipitation years and lack of surface disturbances, native perennial bunch grasses and forbs will reestablish.

Community Phase Pathway 2.1B
This pathway is very unlikely, but can occur when a fire or vegetation manipulation happens to the trees. Two situations can make this occur:

1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires. Seeding after the tree removal may be necessary to help facilitate the return of understory species. Seeding depending on the species may take this community phase into state 4 (Seeded State).

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 2.2: Mature Pinyon-Juniper Woodland

Mature pinyon and Utah juniper woodland with a well-developed understory would characterized this community phase. This phase supports a diverse understory of grasses, forbs and shrubs. Depending on the timing of precipitation, cool season grasses, like Indian ricegrass or warm season grasses like galleta could be dominant. Interspaces supporting highly developed biological crusts are common. Sparse invasive introduced plants species would be present in this phase.

Community Phase Pathway 2.2A
This pathway occurs during and after events such as drought or beetle infestations. Droughts and insects can kill pinyon trees, increasing nutrient availability in the system. Due to the natural conditions of drought, grasses typically do not take up the extra nutrients in the long term. In the short term, grasses and forbs may increase for a few years until Juniper recover. Utah Juniper are more able to compete for these nutrients and became the dominant overstory tree over time.

Community Phase Pathway 2.2B
This pathway is very unlikely to occur naturally with fire. But, vegetation manipulation can be used to remove trees. Two situations occur naturally: 1) a fire can carry in the understory after several wet years allow fine fuels to accumulate, or 2) as the woodland approaches the later stages of development where canopies become dense and crown sizes have increased, and thus community phase becomes susceptible to crown fires. Seeding after the tree removal may be necessary to help facilitate the return of understory species. Seeding depending on the species may take this community phase into state 4 (Seeded State).

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 2.3: Perennial Shrubland with scattered PJ

The overall aspect of this community phase is grassland with scattered pinyon and Utah juniper. The herbaceous understory has a mix of grasses and forbs. This community phase is a result of a crown fire or sufficiently large and hot ground fire that will kill many of the trees, combined with sufficient seed-banks and moisture for reestablishment of grasses and forbs. It is common that after a crown fire many patches of trees will remain unburned, because of fire's unpredictability and broken topography. This leaves a seed bank for the burned areas. This community phase is very short lived in comparison to the other community phases in this state. Sparse invasive introduced plants species would be present in this phase.

Community Phase Pathway 2.3A
This pathway occurs when the climate favors the establishment and growth of trees. More energy is taken-up and stored in the trees as the length between fires and droughts increase. In addition, when shrubs establish on the site they can provide safe-sites for tree establishment furthering the presence of trees.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Transition T2B

This transition is from tree canopy reduction and re-establishment of grasses and forbs. If the community is approaching state 3 (pinyon-juniper invasive state), due to a loss of understory and increase invasive plants this pathway of seeding could be preferable to doing nothing. This pathway may facilitate the recovery of the soils. The infrequent naturally occurring fires could also cause this transition. Reseeding after a fire may be the only way to successfully restore the ecological dynamics to a site. Either way this pathway involves large energy and monetary inputs by man.

Transition T2A

When this transition to state 3 occurs the site has lost much of its expected resistance and resilience. At this point natural and/or management actions have decreased the understory to a point where erosion increases. Reduced influence from fire, insects, and drought could cause the tree canopy to close, effectively reducing the herbaceous understory thus facilitating the transition. Improper grazing and or increase surface disturbance combined with periods of drought can facilitate this transition because soil stability is lost and susceptibility to soil loss increases.

State 3: Pinyon-Juniper Invasive State

This state occurs when there is an absence of natural disturbance (i.e. Insects and drought and/or fire) over long time frames (Zlatnik, 1999). Also, management actions could have allowed trees to become very mature and have effectively closed out the understory. Invasive plants have increased in abundance. This state has the lowest resiliency and resistance of any state in this model. There may be no practicable way back to the Current Potential State (State 2), due to the large amounts of energy and monetary inputs that are needed. Seeding, with either natural disturbance and/or vegetation management to transition it to State 3 (Seeded State) may be the best long term option for this site.

Community Phase 3.1: PJ Woodland with Invasives

A lack of understory with a canopy of older Pinyon and Juniper, where plant interspaces very large and connected. This community phase occurs when natural or management actions allow for the increase in Pinyon and Utah juniper and a decrease in the grass and forb understory. Invasive introduced plants species would be present in this phase and are increasing.

Community Phase Pathway 3.1A
This pathway occurs when events such as frequent fire or drought remove the trees and shrubs, and facilitate the continued establishment of cheatgrass or other invasive annuals. Cheatgrass will typically invade/increase in tree/shrub interspaces when PJ communities are degraded. Once the cheatgrass establishes the amount and continuity of fine fuels increases. This can reduce the fire return interval and shorten the time between fires. When fire eliminates the tree/shrub/native grass component, it completes the conversion to annual dominant community

phase. Cheatgrass and other invasive annuals can persist for long periods of time. Once a fire or a drought removes the trees/shrubs, it is difficult to reestablish because, not only has the fire return interval been shortened to a time that will not allow seedling establish, the soil and other abiotic factors have been altered.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 3.2: Invasive Annuals

This state is characterized by annual grasses like cheatgrass, annual wheatgrass dominating the understory. Also, invasive forbs like storkbill, halogeton and others may be present. This community phase has active erosion under the pinyon and Utah juniper canopy. Utah Juniper has allelopathic effects on some plant (i.e. Sandberg bluegrass, blue grama), which cheatgrass does not appear to suffer this effect when growing under juniper canopies (Zlatnik, 1999).

Community Phase Pathway 3.2A
This pathway is when there is a lack of fire and/or disturbance. The fire return interval lengthens. This could be done by having firebreaks and/or fire suppression which will allow the perennial species a chance to establish with natural processes or with vegetation manipulation.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Transition T3A

Vegetation treatment can transition it to a seeded state. Because of the soils (shallow and/or rocky) and the unpredictable precipitation, this pathway should be used cautiously. This pathway involves large energy and monetary inputs by man.

State 4: Seeded State

This state is a result seeding plants species. Vegetation manipulation may or may not have been done depending on disturbance history of the location. The trees were removed and adapted grasses, forbs and shrubs are established. Plants can be native or introduced depending on the desired management goals. If grazing tolerant species were established these communities can better withstand grazing and other

disturbances. Due to the shallow or rocky soils and unpredictable precipitations patterns, it is difficult to establish grasses from seed, so this state may be hard to achieve and require large energy inputs.

Community Phase 4.1: Seeded Grassland/Shrubland

This community phase appears as a grassland with scattered shrubs and trees. The vegetative production is typically higher than in the current potential state, depending on grass species seeded; however the grass is still sparse due to the low water holding capacity of soils associated with pinyon and juniper.

Community Phase Pathway 4.1A
This pathway occurs when events favor the establishment of shrubs and trees, including long periods without disturbances.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Community Phase 4.2: Seeded with PJ

This community phase has a dense under story of introduced grasses and forbs, but a canopy of pinyon and Utah juniper are establishing. Native perennial grasses, forbs, and shrubs may also be starting to establish. Interspaces are filled with biological crusts and herbaceous plants.

Community Phase Pathway 4.2A
This pathway occurs as trees and shrubs are removed from the community, either naturally through insect herbivory or through vegetation manipulation by man.

Plant Growth Curve

Growth curve number:	CO0102
Growth curve name:	MLRA 34B & 36 - Semidesert Sites
Growth curve description:	MLRA 34B and 36

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	20	30	30	10	0	0	5	5	0	0



Transition T4A

This transition occurs when events favor the establishment and dominance of invasive annuals. Events may include an extended drought, surface disturbance such as off road vehicle use, and/or a shortened fire return interval, all of which can stress the native perennial bunchgrasses.

Section II: Ecological Site Interpretations

Animal Community

The following is from 1996 Range Site:

Grazing:
This site is only of low value for grazing due to low production. It is best used during the late fall, winter, and early spring months when snow still covers the high elevation grazing lands. Care needs to be exercised to prevent site deterioration when grazing. Many areas are short of livestock water so these places must be used during times when snow can provide the animals with a source of water.

The following is from 1996 Range Site:

Wildlife values:

Due to the low productivity of this site, very little manipulation of the vegetation is recommended. If areas of juniper or sagebrush are dense enough to create monoculture situations then some form of chemical or mechanical brush management should be considered. If adequate fuel is available prescribed burning may be considered. In areas where sagebrush and other shrubs become dominant, manipulation that restores a mix of grasses and forbs is beneficial for many wildlife species. Areas of sagebrush along drainages and south and west facing slopes should be left undisturbed to provide critical winter forage for mule deer. Water developments for livestock and wildlife can be a useful management tool on this site.

Wildlife species list:
mule deer, coyote, cottontail, bushy tailed rat, golden eagle, bald eagle, pinyon jay, rock wren, Rocky Mountain elk, mountain lion, white-tailed prairie dog, white-tailed jackrabbit, side blotched lizard, red-tailed hawk, ash-throated flycatcher, western blue bird, bobcat, rock squirrel, gopher snake, sagebrush lizard, chuckar, mourning dove, and plain titmouse.

Plant Preference by Animal Kind

Animal kind: Mature antelope

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Leaves	N	N	N	P	P	P	N	N	N	D	D	D
rosy pussytoes	*Antennaria rosea*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
black sagebrush	*Artemisia nova*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
Fendler threeawn	*Aristida purpurea var. longiseta*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Leaves	P	P	P	D	D	D	D	D	D	P	P	P
shadscale saltbush	*Atriplex confertifolia*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
blue grama	*Bouteloua gracilis*	Entire plant	D	D	D	P	P	P	P	P	P	D	D	D
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
longflower rabbitbrush	*Chrysothamnus depressus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
squirreltail	*Elymus elymoides*	Entire plant	U	U	U	D	D	D	U	U	U	U	U	U
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
needle and thread	*Hesperostipa comata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hairy false goldenaster	*Heterotheca villosa*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Utah juniper	*Juniperus osteosperma*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
prairie Junegrass	*Koeleria macrantha*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
winterfat	*Krascheninnikovia lanata*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Colorado four o'clock	*Mirabilis multiflora*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

plains pricklypear	*Opuntia polyacantha*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
James' galleta	*Pleuraphis jamesii*	Leaves	U	U	U	D	D	P	P	D	D	U	U	U
threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Animal kind: Mature cattle

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Leaves	D	D	P	P	P	D	U	U	U	D	D	D
rosy pussytoes	*Antennaria rosea*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
black sagebrush	*Artemisia nova*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Fendler threeawn	*Aristida purpurea var. longiseta*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Leaves	D	D	D	U	U	U	U	U	U	D	D	D
shadscale saltbush	*Atriplex confertifolia*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
blue grama	*Bouteloua gracilis*	Entire plant	U	U	U	U	D	D	P	P	P	U	U	U
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
longflower rabbitbrush	*Chrysothamnus depressus*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
squirreltail	*Elymus elymoides*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
needle and thread	*Hesperostipa comata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hairy false goldenaster	*Heterotheca villosa*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Utah juniper	*Juniperus osteosperma*	Entire plant	E	E	E	E	E	E	E	E	E	E	E	E
prairie Junegrass	*Koeleria macrantha*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
winterfat	*Krascheninnikovia lanata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Colorado four o'clock	*Mirabilis multiflora*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
plains pricklypear	*Opuntia polyacantha*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	N	N	N	N	N	N	N	N	N	N	N	N
James' galleta	*Pleuraphis jamesii*	Leaves	U	U	U	U	D	P	P	D	D	U	U	U
threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Animal kind: Mule deer

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Leaves	N	N	N	P	P	P	N	N	N	D	D	D
rosy pussytoes	*Antennaria rosea*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
black sagebrush	*Artemisia nova*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
Fendler threeawn	*Aristida purpurea var. longiseta*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Leaves	P	P	P	U	U	U	D	D	P	P	P	P
shadscale saltbush	*Atriplex confertifolia*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
blue grama	*Bouteloua gracilis*	Entire plant	D	D	D	P	P	P	P	P	P	D	D	D
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
longflower rabbitbrush	*Chrysothamnus depressus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P

squirreltail	*Elymus elymoides*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
needle and thread	*Hesperostipa comata*	Entire plant	U	U	U	D	D	D	D	D	U	U	U	U
hairy false goldenaster	*Heterotheca villosa*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Utah juniper	*Juniperus osteosperma*	Entire plant	D	D	X	X	X	X	X	X	D	D	D	D
prairie Junegrass	*Koeleria macrantha*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
winterfat	*Krascheninnikovia lanata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Colorado four o'clock	*Mirabilis multiflora*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
plains pricklypear	*Opuntia polyacantha*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	X	X	X	X	X	X	X	X	X	X	X	X
James' galleta	*Pleuraphis jamesii*	Leaves	U	U	U	D	D	P	P	D	D	U	U	U
threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Animal kind: Mature elk

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Leaves	D	D	P	P	P	D	U	U	U	D	D	D
rosy pussytoes	*Antennaria rosea*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
black sagebrush	*Artemisia nova*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
Fendler threeawn	*Aristida purpurea var. longiseta*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Leaves	D	D	D	U	U	U	U	U	U	D	D	D
shadscale saltbush	*Atriplex confertifolia*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
blue grama	*Bouteloua gracilis*	Entire plant	U	U	U	U	D	D	P	P	D	U	U	U
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
longflower rabbitbrush	*Chrysothamnus depressus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	D	D	D	U	U	U	U	U	U	D	D	D
squirreltail	*Elymus elymoides*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
needle and thread	*Hesperostipa comata*	Entire plant	D	D	D	P	P	P	D	D	D	D	D	D
hairy false goldenaster	*Heterotheca villosa*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Utah juniper	*Juniperus osteosperma*	Entire plant	X	X	X	X	X	X	X	X	X	X	X	X
prairie Junegrass	*Koeleria macrantha*	Entire plant	U	U	D	D	D	U	U	U	U	U	U	U
winterfat	*Krascheninnikovia lanata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
plains pricklypear	*Opuntia polyacantha*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	N	N	N	N	N	N	N	N	N	N	N	N
James' galleta	*Pleuraphis jamesii*	Leaves	U	U	U	U	D	P	P	D	D	U	U	U
threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Animal kind: Adult horses

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
rosy pussytoes	*Antennaria rosea*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U

black sagebrush	*Artemisia nova*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
purple threeawn	*Aristida purpurea*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
shadscale saltbush	*Atriplex confertifolia*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
blue grama	*Bouteloua gracilis*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
longflower rabbitbrush	*Chrysothamnus depressus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
squirreltail	*Elymus elymoides subsp. elymoides*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
needle and thread	*Hesperostipa comata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hairy false goldenaster	*Heterotheca villosa*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Utah juniper	*Juniperus osteosperma*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
prairie Junegrass	*Koeleria macrantha*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
winterfat	*Krascheninnikovia lanata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Colorado four o'clock	*Mirabilis multiflora*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
plains pricklypear	*Opuntia polyacantha*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
James' galleta	*Pleuraphis jamesii*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Animal kind: Mature sheep

Common name	Scientific name	Plant part	J	F	M	A	M	J	J	A	S	O	N	D
Indian ricegrass	*Achnatherum hymenoides*	Leaves	N	N	N	P	P	P	N	N		D	D	D
rosy pussytoes	*Antennaria rosea*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
black sagebrush	*Artemisia nova*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
Fendler threeawn	*Aristida purpurea var. longiseta*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
Wyoming big sagebrush	*Artemisia tridentata subsp. wyomingensis*	Leaves	D	D	D	U	U	U	U	U	U	D	D	D
shadscale saltbush	*Atriplex confertifolia*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
blue grama	*Bouteloua gracilis*	Entire plant	D	D	D	P	P	P	P	P	P	D	D	D
littleleaf mountain mahogany	*Cercocarpus intricatus*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
Douglas' dustymaiden	*Chaenactis douglasii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
yellow rabbitbrush	*Chrysothamnus viscidiflorus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
squirreltail	*Elymus elymoides*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
mormon tea	*Ephedra viridis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spreading buckwheat	*Eriogonum effusum*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
broom snakeweed	*Gutierrezia sarothrae*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
needle and thread	*Hesperostipa comata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hairy false goldenaster	*Heterotheca villosa*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Utah juniper	*Juniperus osteosperma*	Entire plant	E	E	E	E	E	E	E	E	E	E	E	E
prairie Junegrass	*Koeleria macrantha*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
winterfat	*Krascheninnikovia lanata*	Entire plant	P	P	P	P	P	P	P	P	P	P	P	P
hoary tansyaster	*Machaeranthera canescens var. canescens*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
Colorado four o'clock	*Mirabilis multiflora*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
plains pricklypear	*Opuntia polyacantha*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
spiny phlox	*Phlox hoodii*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
twoneedle pinyon	*Pinus edulis*	Entire plant	N	N	N	N	N	N	N	N	N	N	N	N
James' galleta	*Pleuraphis jamesii*	Leaves	U	U	U	D	D	P	P	D	D	U	U	U

threadleaf ragwort	*Senecio flaccidus var. flaccidus*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
scarlet globemallow	*Sphaeralcea coccinea*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D
stemless mock goldenweed	*Stenotus acaulis*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
desert princesplume	*Stanleya pinnata*	Entire plant	U	U	U	U	U	U	U	U	U	U	U	U
soapweed yucca	*Yucca glauca*	Entire plant	D	D	D	D	D	D	D	D	D	D	D	D

Legend: P=Preferred; D=Desirable; U=Undesirable; N=Not consumed; E=Emergency; T=Toxic; X=Used, but degree of utilization unknown

Hydrology Functions

Soils were originally assigned to hydrologic soil groups based on measured rainfall, runoff, and infiltrometer data (Musgrave 1955). Since the initial work was done to establish these groupings, assignment of soils to hydrologic soil groups has been based on the judgment of soil scientists. Assignments are made based on comparison of the characteristics of unclassified soil profiles with profiles of soils already placed into hydrologic soil groups. Most of the groupings are based on the premise that soils found within a climatic region that are similar in depth to a restrictive layer or water table, transmission rate of water, texture, structure, and degree of swelling when saturated, will have similar runoff responses. Four (4) Hydrologic Soil Groups are recognized (A-D). For specific definitions of each hydrologic soil group see the National Engineering Handbook, Chapter 7, Part 630 Hydrology, or visit:http://policy.nrcs.usda.gov/OpenNonWebContent.aspx?content=22526.wba The hydrologic soil groups are based on the following factors:

-intake and transmission of water under the conditions of maximum yearly wetness (thoroughly wet)
- soil not frozen
- bare soil surface
- maximum swelling of expansive clays

The slope of the soil surface is not considered when assigning hydrologic soil groups. In its simplest form, the hydrologic soil group is determined by the water transmitting soil layer with the lowest saturated hydraulic conductivity and depth to any layer that is more or less water impermeable (such as a fragipan or duripan) or depth to a water table (if present) (Caudle, et. al, 2013). The runoff curve numbers are determined by field investigations using hydrologic cover conditions and hydrologic soil groups.

Soils Hydrologic Group

Loamy Soils

Lazear - D
Barboncito - D

Loamy-Skeletal Soils

Walknolls - D

Hydrologic soil groups are based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms (Soil Survey Staff, 2015).

The soils in the United States are assigned to four groups (A, B, C, and D) and three dual classes (A/D, B/D, and C/D). The groups are defined as follows:

Group A. Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.

Group B. Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C. Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D. Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission (Soil Survey Staff, 2015).

Recreational Uses

The following is from 1996 Range Site:

This site occurs at lower elevations so is accessible for recreation nearly year around. It offers beauty in the spring with blooming forbs and hunting opportunities in the fall. Summers are hot but many higher elevation recreation areas are only accessible by travelling through this site.

Wood Products

The following is from 1996 Range Site:

This site does produce scattered Utah juniper trees. The trees grow very slowly and are scrubby in appearance. They are of low value for fence posts and firewood.

Other Information

The following is from 1996 Range Site:

Endangered Plants and Animals:

Bald eagles can be found on this site during the winter. The spineless hedgehog cactus occurs on this site.

Supporting Information

Associated Sites

Site name	Site ID	Site narrative
Saltdesert Overflow	*R034BY407CO*	Saltdesert Overflow is located run-in positions on the landscape. Soils are deep and formed from alluvium and are sandy (i.e. fine sandy loam, sandy loams) in texture. This site is dominated by alkali sacaton, basin wildrye, galleta and, four-wing saltbush, basin big sagebrush and greasewood. This site is in the 8 to 12 inch precipitation zone of semidesert.
Shallow Desert	*R035XY409CO*	Shallow Desert is located edges of mesa tops and high benches. Soils are shallow and are sandy loam and loamy sands surfaces textures. This site is dominated by Saline wildrye, New Mexico Feathergrass, Indian ricegrass, galleta, shadscale, green Mormontea, and basin big sagebrush. This site is in the 8 to 12 inch precipitation zone of semidesert.
Semidesert Loam	*R036XY325CO*	Semidesert Loam are loamy texture soils. Particle control section is fine-loamy. Clay content is higher in these soils than those found in Semidesert Sandy Loam and lower than those found in Semidesert Clay Loam. It is a Wyoming big sagebrush dominated site. Indian Ricegrass and galleta are the dominant grass on this site. The soils on this site are moderately deep to very deep.

Similar Sites

Site name	Site ID	Site narrative
Shallow Clay Loam - (Pinyon-Juniper)	*R036XY110CO*	Shallow Clay Loam Pinyon-Juniper is a gentle sloped (<25% slopes) site with shallow soils that are clayey in texture. This site is dominated by Utah Juniper and scattered pinyon. This site may have Wyoming big sagebrush in the understory. This site is in the 8 to 12 inch precipitation zone of semidesert.
Steep Shallow Clay Loam - (Pinyon-Juniper)	*R036XY111CO*	Steep Shallow Clay Loam Pinyon-Juniper is a very steep sloped (> 25% slopes) site with shallow soils that are clayey in texture. This site is dominated by Utah Juniper and scattered pinyon. This site may have Wyoming big sagebrush in the understory. This site is in the 8 to 12 inch precipitation zone of semidesert.
Mountain Pinyon	*R036XY114CO*	Mountain Pinyon is a gentle sloped (<25% slope) site with very shallow and shallow soils that are loamy in texture. This site is dominated by Pinyon, Utah Juniper. This site may have oakbrush in the understory. This site is in the 12 to 16 inch precipitation zone of foothills/upland.
Shallow Loamy Mesa Top - (Pinyon-Juniper)	*R036XY141CO*	Shallow Loamy Mesa Top is a gentle sloped (<25% slope) site with very shallow and shallow soils that are loamy in texture. This site is dominated by Pinyon, Utah Juniper, muttongrass and Indian ricegrass. This site is in the 15 to 18 inch precipitation zone of foothills/upland.
Loamy Mesa Top - (Pinyon-Juniper)	*R036XY142CO*	Loamy Mesa Top is a gentle sloped (<15% slope) site with moderately deep to deep soils that are coarse loamy in texture. This site is shallow to calcic horizon. The typical profile is border-line skeletal which reduces the water holding capacity of this site. It is dominated by Pinyon, Utah Juniper, muttongrass and Indian ricegrass. This site is in the 15 to 18 inch precipitation zone of foothills/upland.
Stony Foothills	*R036XY287CO*	Stony Foothill is a gentle sloped (<25% slope) site with moderately deep to deep that are loamy-skeletal in texture. This site is dominated by Pinyon, Utah Juniper. This site may have oakbrush in the understory. This site is in the 12 to 16 inch precipitation zone of foothills/upland.
Cobbly Foothills	*R036XY346CO*	Cobbly Foothill is a gentle sloped (<20% slope) site with moderately deep to deep soils that are loamy-skeletal in texture. Common surface textures are cobbly or gravelly loam. This site is dominated by Big sagebrush, western wheatgrass, Pinyon, and Utah Juniper. This site is in the 12 to 16 inch precipitation zone of foothills/upland.
Steep Colluvial Slopes	*R036XY445CO*	Steep Colluvial Slopes is a very steep (>25% slope) sloped site with very shallow to shallow soils that are clayey in texture. This site is dominated by Utah Juniper and pinyon. This site may have Wyoming big sagebrush in the understory. This site has higher precipitation (12 to 16") than Semidesert Loam (8 to 12"). The temperature is slightly cooler than the semidesert site.

		Foothill site will be found at elevations above the semidesert site. The soils are similar in nature.
Southwestern Mountain - (Pinyon-Juniper)	*R036XY446CO*	Southwestern Mountain (Pinyon-Juniper) is a gentle sloped (<25% slope) site with very shallow and shallow soils that are loamy or loamy-skeletal in texture. This site is dominated by Pinyon, Utah Juniper, Wyoming big sagebrush, muttongrass and Indian ricegrass. This site may have oakbrush in the understory. This site is in the 12 to 16 inch precipitation zone of foothills/upland.

State Correlation
This site has been correlated with the following states: CO

Inventory Data References

Baisan, C. H. and T. W. Swetnam. 1990. Fire history on a desert mountain range: Rincon Mountain Wilderness, Arizona, USA. Canadian Journal of Forest Research. 20:1559-1569.

Betancourt, J. L., E. A. Pierson, K. A. Rylander, J. A. Fairchild-Parks, and J. S. Dean. 1993. Influence of history and climate on New Mexico pinyon-juniper woodlands. In: Gen. Tech. RM-236 - Managing Pinon-Juniper Ecosystems for Sustainability and Social Needs.

Cartledge, T. R., and J. G. Propper. 1993. Pinon-Juniper Ecosystems through Time: Information and Insights from the Past. In Gen. Tech. RM-236 - Managing Pinon-Juniper Ecosystems for Sustainability and Social Needs.

Floyd, M.L., W.H. Romme, and D.D. Hanna. 2000. Fire History and vegetation pattern in Mesa Verde National Park, Colorado, USA. Ecological Applications. 10:1666-1680.

Floyd, M. L., D. D. Hanna, W. H. Romme. 2004. Historical and recent fire regimes in pinyon-juniper woodlands on Mesa Verde, Colorado, USA. Forest Ecology and Management. 198:269-289.

Kennard, D.K. and A.J. Moore. 2013. Fire history, woodland structure, and mortality in pinon-juniper woodland in the Colorado National Monument. Natural Areas Journal. 33:296-306.

Miller, R. F. and R. J. Tausch. 2001. The role of fire in juniper and pinyon woodlands: a descriptive analysis. In: Galley, K.E.M.; Wilson. T.P., [EDs]. Proceedings of the invasive species workshop: the role of fire in the control and spread on invasive species. Fire conference 2000. Tallahassee, Fl: Tall Timbers Research Station: Miscellaneous publication 11:15-30.

Miller, R. F. and J. A. Rose. 1999. Fire history and western juniper encroachment in sagebrush steppe. Journal of Range Management. 52:550-559.

Musgrave, G.W. 1955. How much of the rain enters the soil? In Water: U.S. Department of Agriculture Yearbook. Washington, D.C. P. 151-159.

National Engineering Handbook. US Department of Agriculture, Natural Resources Conservation Service. Available: http://www.info.usda.gov/CED/Default.cfm#National%20Engineering%20Handbook Accessed February 25, 2008.

Natural Resources Conservation Service (NRCS). March 1996. Range Site Description for Semidesert Juniper Loams #113: USDA, Denver Colorado

Passey, H. B., W. K. Hugie, E. W. Williams, and D. E. Ball. 1982. Relationships between soil, plant community, and climate on rangelands of the Intermountain west. USDA, Soil Conservation Service, Tech. Bull. No. 1669.

Romme, W. H., C.D. Allen, J.D. Bailey, W.L. Baker, B.T. Bestelmeyer, P.M. Brown, K.S. Eisenhart, M.L. Floyd, D.W. Huffman, B.F. Jacobs, R.F. Miller, E.H. Muldavin, T.W. Swetnam, R.J. Tausch, and P.J. Weisberg. 2009. Historical and Modern Disturbance Regimes, Stand Structures, and Landscape Dynamics in Pinon-Juniper Vegetation of the Western United States. Rangeland Ecology and Management 62:203-222.

Shinneman, D. J. and W. L, Baker. 2009. Historical fir and multidecadal drought as context for pinon-juniper woodland restoration in western Colorado. Ecological Applications 19: 1231-1245.

Soil Survey Staff, Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey. Available online at http://websoilsurvey.nrcs.usda.gov/. Accessed [5/3/2017].

Swetnam, T. and Baisan, C. 1996. Historical fire regime patterns in the southwestern United States since AD 1700. In: CD Allen (ed.) Fire Effects in Southwestern Forest: Proceedings of the 2nd La Mesa Fire Symposium, pp. 11-32. USDA Forest Service, Rocky Mountain Research Station, General Technical Report RM-GTR-286.

Tausch, R. J., N. E. West, and A. A. Nabi. 1981. Tree age and dominance patterns in Great Basin pinyon-juniper woodlands. Journal of Rangeland Management 34:259-264.

Western Regional Climate Center. Retrieved from http://www.wrcc.dri.edu/summary/Climsmco.html on February 9, 2017.

Zlatnik, E. 1999. Juniperus osteosperma. In: Fire Effects Information System. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). http://www.fs.fed.us/database/feis/. Accessed March 27, 2017.

Type Locality

State:	CO
County:	Montezuma
Township:	13S
Range:	99W
Section:	8
General legal description:	1,200' w & 1300' S of the NE corner, Sec.8 T13S, R99W, Mesa County. E and N side of Sleeping Ute mountain, non-sectionalized T33 1/2N, R18W, Montezuma County.

Hierarchical Classification Relationships

NRCS & BLM:
Major Land Resource Area 36, Southwestern Plateaus Mesas and Foothills (United States Department of Agriculture, Natural Resources Conservation Service, 2006).

USFS:
313Aa-San Juan Basin-Mesa Verde, and 313Ab-Canyon of Ancients-Blanding Basin subsections <313A Grand Canyons Section < 313 Colorado Plateau Semi-Desert (Cleland, et al., 2007).

313Bc Chuska Valley Cold Desert Shrubland Subsection <313B Navaho Canyonlands Section < 313 Colorado Plateau Semi-Desert (Cleland, et al., 2007).

341Ba-Mancos Shale Lowlands-Grand Valley, 341Bp-Uncompahgre Plateau and 341Bg-Northeast Flank Subsections <341B Northern Canyonlands Section < 341 Intermountain Semi-desert and Desert (Cleland, et al., 2007).

EPA:
20b Shale Deserts and Sedimentary Basins and 20c Semiarid Benchlands and Canyonlands, < 20 Colorado Plateau < 10.I Cold Deserts < 10 North American Deserts (Griffith, 2006).

USGS:
Colorado Plateau Province (Canyonlands Section)

Acknowledgments:

Project Staff:
Suzanne Mayne-Kinney, Ecological Site Specialist, NRCS MLRA, Grand Junction SSO
Chuck Peacock, MLRA Soil Survey Leader, NRCS MLRA Grand Junction SSO

Program Support:
Rachel Murph, NRCS CO State Rangeland Management Specialist, Denver
Scott Woodhall, NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ
Eva Muller, Regional Director, Rocky Mountain Regional Soil Survey Office, Bozeman, MT
B.J. Shoup, CO State Soil Scientist, Denver
Eugene Backhaus, CO State Resource Conservationist, Denver

Partners/Contributors:
Those involved in developing earlier versions of this site description include: Herman Garcia, retired CO State RMS and NRCS MLRA Ecological Site Specialist-QA Phoenix, AZ.

--Site Development and Testing Plan--:

Future work to validate and further refine the information in this Provisional Ecological Site Description is necessary. This will include field activities to collect low-, medium-, and high-intensity sampling, soil correlations, and analysis of that data.

Additional information and data is required to refine the Plant Production and Annual Production tables for this ecological site. The extent of MLRA 36 must be further investigated.

Field testing of the information contained in this Provisional ESD is required. As this ESD is moved to the Approved ESD level, reviews from the technical team, quality control, quality assurance, and peers will be conducted.

Site Authors

Suzanne Mayne Kinney
Jim Kellogg John Murray LJJ

Certifications

NRCS Science and Technology Certified by Rachel Murph (9/18/2017)

Quality Assurance

Provisional Status Verfied by Scott Woodall (12/21/2017)

Reference Sheet

Certification: Reference Sheet Certified by Rachel Murph (9/18/2017)

Author(s)/participant(s): Original written by Steve Meyer (5/12/2006) Used the 2017 verison of R036XY110CO worksheet to updated Semidesert Juniper Loams worksheet 9/18/2017 - Suzanne Mayne-Kinney

Contact for lead author:

Date: 9/18/2017 **MLRA:** 036X **Ecological Site:** Semidesert Juniper Loam R036XY113CO This *must* be verified based on soils and climate (see Ecological Site Description). Current plant community cannot be used to identify the ecological site.

Composition (indicators 10 and 12) based on: X Annual Production, Foliar Cover, Biomass

---

Indicators. For each indicator, describe the potential for the site. Where possible, (1) use numbers, (2) include expected range of values for above- and below-average years for **each** community and natural distrurbance regimes within the reference state, when appropriate and (3) cite data. Continue descriptions on separate sheet.

---

1. **Number and extent of rills:** On more gentle slopes (< 15 %): Common and occur throughout site. Rills may be 10 or more feet in length. Sides of rills may be up to 4 inches high. Rills are most likely to form below adjacent exposed bedrock or water flow patterns where sufficient water accumulates to cause erosion. B. On steep slopes (> 15 %): Frequent. Occur throughout the site. Rills may extend down entire slope.

---

2. **Presence of water flow patterns:** Flow paths are expected and frequent. They are disconnected with debris dams, As slope increases, flow paths become more prominent and connected.

---

3. **Number and height of erosional pedestals or terracettes:** Pedestals are rare and may form at the base of plants that occur on the edge of flow paths. Terracettes are rare to few, forming behind debris dams of small to medium sized litter (up to 2 inches in diameter) may form in water flow patterns. These debris dams may accumulate smaller litter (leaves, grass and forb stems) and sediment. Terracettes or debris dams are more obvious following intense rainfall events.

---

4. **Bare ground from Ecological Site Description or other studies (rock, litter, standing dead, lichen, moss, plant canopy are not bare ground):** 20-30% bare ground. (Soil surface is typically covered 0 to 15 percent surface fragments). Most bare ground is associated with water flow patterns, rills, and gullies. Areas with well-developed biological soil crusts should not be counted as bare ground. Poorly developed biological soil crusts that are interpreted as functioning as bare ground (therefore they would be susceptible to raindrop splash erosion) should be recorded as bare ground. Ground cover is based on first raindrop impact, and bare ground is the opposite of ground cover. Ground cover + bare ground = 100%.

---

5. **Number of gullies and erosion associated with gullies:** None to few on gentle slopes (< 15 %). On steep slopes and areas below adjacent exposed bedrock, gullies are more prone to occur. Length often extends from exposed bedrock until gully reaches a stream or an area where water and sediment accumulate. Gullies may remove soil from base of plants exposing roots. Gullies may show slightly more erosion as slope increases, or as the site occurs adjacent to steep sites with concentrated flow patterns.

---

6. **Extent of wind scoured, blowouts and/or depositional areas:** None to very few. Trees break the wind and reduce the potential for wind erosion. The surface fragments armor the soil surface and help to reduce the potential for wind erosion.

---

7. **Amount of litter movement (describe size and distance expected to travel):** Most litter accumulates at base of plants. Woody stems from trees are not moved unless present in water flow patterns, rills, or gullies.

---

8. **Soil surface (top few mm) resistance to erosion (stability values are averages - most sites will show a range of values):** This site should have an erosion rating of 5 to 6 under the plant canopies, and a rating of 3 to 4 in the interspaces. The average should be a 4. Vegetation cover, litter, biological soil crusts and surface rock reduce erosion. This site can have cryptogamic crusts which help to stabilize the soil surface and places with cryptograms should rate a 5-6.

9. **Soil surface structure and SOM content (include type and strength of structure, and A-horizon color and thickness):** Soil organic matter content ranges from 0.25-2%. Soils are very shallow and shallow in depth. The surface soils of this site range from fine sandy loam, sandy loam, and loam. Structure is usually weak fine granular structure. The soil surface (A horizon) ranges from 3 to 6 inches in depth. Refer to soil survey for more detailed information about your specific site.

10. **Effect on plant community composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff:** The presence of trees, perennial grasses and forbs, and shrubs will breakup raindrop impact and splash erosion. The spatial distribution of the plants, biological crusts and interspaces will provide small pockets for water storage and surface roughness that slows down runoff, allowing time for infiltration. The tree and shrub canopy is effective in intercepting rain drops and preventing splash erosion on the reference state. But, with increased tree canopy, understory canopy is reduced, increased bare soil and litter accumulates under trees, it can forms micro-topography that can help water accumulate which can cause more rapid runoff.

11. **Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site):** None, although bedrock is found within 20 inches of soil surface. In addition, there may be layers of calcium carbonate or other naturally occurring hard layers found in the soil subsurface. These should not be considered to be compaction layers.

12. **Functional/Structural Groups (list in order of descending dominance by above-ground weight using symbols: >>, >, = to indicate much greater than, greater than, and equal to) with dominants and sub-dominants and "others" on separate lines:**
    Dominant: Trees (Utah Juniper, pinyon) >
    Sub-dominant: cool season bunchgrasses (Indian ricegrass, squirreltail, needle-and-thread) > shrubs (Wyoming big sagebrush, shadscale, winterfat, yellow rabbitbrush, broom snakeweed) > warm season short bunchgrass (Galleta, Blue grams) >
    Other: forbs (scarlet globemallow, Hood's phlox, rosy pussytoes, hairy goldenaster, stemless goldenweed) > cryptograms
    Additional:

13. **Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence):** A mix of young, middle aged and old pinyon and Utah juniper are expected to be found on this site. In years with average or above average precipitation, shrubs, grasses and forbs should have little mortality or decadence. Tree mortality, especially pinyon, can be expected under severe and/or extended drought and subsequent insect infestations. Under a dense tree canopy, understory has increased decadence and mortality.

14. **Average percent litter cover (20-35%) and depth (0.5-3.0 inches):** Litter cover ranges from 20-35% at a depth of 0.5 to 3.0 inches. Most litter is at the base and under the canopy of the plants.

15. **Expected annual production (this is TOTAL above-ground production, not just forage production):** Tree canopy cover 0-15%: 200-300 lbs./ac. Production figures are for total annual vegetation.

16. **Potential invasive (including noxious) species (native and non-native). List Species which BOTH characterize degraded states and have the potential to become a dominant or co-dominant species on the ecological site if their future establishment and growth is not actively controlled by management interventions. Species that become dominant for only one to several years (e.g., short-term response to drought or wildfire) are not invasive plants. Note that unlike other indicator, we are describing what is NOT expected in the reference state for the ecological site:** Few invasives capable of dominating this site. Cheatgrass, Broom snakeweed, and Mustard may invade the community.

17. **Perennial plant reproductive capability:** All plants have the ability to reproduce in most years. Limitations are weather related, wildfire, natural disease, inter-species competition, and insects may temporarily reduce reproductive capability. Increased tree canopy will result in decreased understory reproductive capability.

Reference Sheet Approval

Approval	Date
Rachel Murph, State Rangeland Management Spec., USDA NRCS Colorado	9/18/2017