

FORAGE SUITABILITY GROUP

Loamy and Clayey Soils on Stream Terraces, Flood Plains or in Depressions

FSG No.: G138XA345FL

Major Land Resource Area (MLRA 138): North-Central Florida Ridge

Map Unit List*

- Meggett fine sand, frequently flooded
- Oleno clay (FL023)
- Pantego fine sandy loam (FL121)

*NOTE: Some members do not identify flooding or depressional in the map unit name but are subject to flooding or ponding. In these cases, please refer to the water features data on the Web Soil Survey or Soil Data Mart. Information in parenthesis refers to soil survey code where map unit occurs.

Adapted Species List

The native forage species listed are considered adapted to grow on the soils in this group at their natural pH levels. All introduced grasses will need native pH raised to min. 5.5 (unless noted) for best production. Consult with state extension service for current cultivar or germplasm recommendations (<http://agronomy.ifas.ufl.edu/foragesofflorida/>).

Perennial Species:

Grasses

Warm season (Introduced)

- Limpograss (*Hemarthria altissima*)

Warm season (Native)

- Maidencane (*Panicum hemitomom*)
- Blue Maidencane (*Amphicarpum muhlenbergianum*)

Annual Species:

Grasses

Warm season

- Japanese Millet (*Echinochloa esculenta*)

Legumes

Warm season

- Aeschynomene (*Aeschynomene americana*)

Seasonal and Total Production Estimates

Unless previously drained, soils in this FSG have very few forage species adapted to their seasonal high water table (1 to 2 feet above the soil surface). If previously drained see forage list and discussion with FSG G138XA341FL.


Expected Range in Dry Matter Production and Animal Unit Months (AUM) for Different Forages				
Forage	Range in Dry Matter, lbs/acre		Range in AUM/acre†	
Limpograss (≈400 lb N/acre) ^{3,5}	8,000	13,000	5.1	8.2
Maidencane ^{1#}	5,000	6,700	3.2	4.3
Blue Maidencane ^{1#}	2,100	2,500	1.3	1.6
Japanese Millet ²	4,000	6,000	2.5	3.8
Aeschynomene ⁴	2,000	3,000	1.3	1.9

†Animal Unit Month based on 50% grazing efficiency and 2.6% intake per day.

#Superscript numbers refer to references.

#Dry matter estimated based on the assumption air dried yield in reference had ≈16% moisture.

Production Curves:



Physiographic Features

Dominantly very deep, nearly level, poorly drained or very poorly drained soils formed in loamy and clayey marine deposits or alluvial deposits. These soils are on flood plains, or in depressions of marine terraces. Diagnostic subsurface horizon is an argillic horizon above 20 inches. A few members have either a mollic or umbric horizon. The organic matter content of the surface layer is dominantly medium or high. Unless limed, the reaction in the surface layer ranges from extremely acid to neutral.

Climatic Features

Freeze-free period (>28° F 9 years in 10 at least):
 averages 270 d (range 266-278 d)

Length of growing season (>32° F 9 years in 10 at least): averages 234 d (range 225-247 d)

Annual minimum temperature (° F in month of January):
 averages 41.1 (range 38.6-43.5)

USDA Plant Hardiness Zone:
 8b (15-20° F, Jacksonville)
 8a (10-15° F, Glen St. Mary)

Mean annual precipitation (inches):
 averages 54.40 (range 52.24-59.65)

Soil Properties

Percent Slope: 0 to 2 percent

Surface Texture: Dominantly sandy loam, fine sandy loam, loam, clay, fine sand, loamy sand, loamy fine sand, and their mucky analogs. A few members are sand, very fine sandy loam, silt loam, sandy clay loam, clay loam, silty clay loam, and sandy clay.

Sand Content of Surface Layer: 26 to 95 percent

Clay Content of Surface Layer: 5 to 74 percent

Organic Matter Content of Surface Layer: 1 to 20 percent

Cation Exchange Capacity of Surface Layer (meq/100g):
 1.3 to 54.4

Effective Cation Exchange Capacity of Surface Layer (meq/100g): 2.4 to 72.1

Bulk Density of Surface Layer (g/cc): 1.3 to 1.68

Saturated Hydraulic Conductivity of Surface Layer:
 Moderate to very rapid

Soil Reaction of Surface Layer: 3.5 to 7.3

Available Water Capacity (0 to 30 inches): 0.2 to 6.0 inch per inch

Depth to Finer Textured Material: Less than 20 inches

Depth to Bedrock: Greater than 80 inches. Some members have bedrock at less than 80 inches.

Drainage Class (Agronomic): Very poorly, poorly

Depth to Seasonal High Water Table (during wet periods): 1.0 to 2.0 feet above the surface

Flooding: If flooded, frequent or occasional with brief to very long duration

Ponding: If ponded, Long or very long duration

Monthly precipitation (inches) and temperature (F):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip avg	4.89	3.82	5.09	3.28	3.27	6.23	6.84	7.12	4.86	2.98	2.69	3.34
Avg Min	41.1	43.6	49.4	54.3	62.0	68.5	71.1	70.8	67.9	57.7	50.0	43.2
Avg Temp	55.1	56.3	62.3	67.5	74.4	79.6	81.6	81.2	78.4	69.8	62.4	55.4
Avg Max	65.7	68.9	75.2	80.5	86.8	90.7	92.0	91.5	88.9	82.1	74.7	67.6

Climate Station Locations (averages from 1971 to 2000; see Appendix 1)

FSG Documentation

Inventory Data References:

1. ----- . 1987. Range Management for Important Native Grasses of Florida. USDA, NRCS. Gainesville, FL. 168 p.
2. Barnhart, S. 2008. Forage and Cover Crop Considerations for Delayed Planting and Flooded Sites. Iowa State Univ. Ext.
<http://www.extension.iastate.edu/CropNews/2008/0611SteveBarnhart.htm> (accessed Oct. 12, 2011).
3. Kalmbacher, R.S., P.H. Everett, F.G. Martin, K.H. Quesenberry, E.M. Hodges, O.C. Ruelke, and S.C. Schank. 1987. Yield and Persistence of Perennial Grasses at Immokalee, Florida: 1981 to 1984. Univ. Florida, IFAS, Agric. Exp. Stn., Bull. 865.
<http://ufdc.ufl.edu/UF00027614/00001?search=kalmbacher>, accessed August 22, 2011).
4. Mislevy, P., R.S. Kalmbacher, and F.G. Martin. 1981. Cutting Management of the Tropical Legume American Jointvetch. Agron. J. 73:771-775
<https://www.agronomy.org/publications/aj/abstracts/73/5/AJ0730050771>, accessed September 29, 2011).
5. Newman Y.C., A. Agyin-Birikorang, M.B. Adjei, J.M. Silveira, J.M.B. Vendramini, J.E. Rechcigl, and L.E. Soltenberger. 2009. Nitrogen Fertilization Effect on Phosphorus Recommendation Potential of Three Perennial Warm-Season Forages. Agron. J. 101:1243-1248.
<https://www.soils.org/publications/aj/pdfs/101/5/1243>, accessed August 22, 2011).

State Correlation: (NA)

Forage Suitability Group Approval:



Greg Hendricks, State Resource Conservationist



Tom Weber, State Soil Scientist

Appendix 1: Climate Station Locations		
COOP ID (FL=08)	Location	County
4731	Lake City	Columbia
4394	Jasper	Hamilton
5539	Mayo	LaFayette
9120	Usher Tower	Levy
5275	Madison	Madison
5099	Live Oak	Suwannee