

## FORAGE SUITABILITY GROUP

### Sandy Over Loam, Loamy and Clayey Soils on Ridges and Side Slopes of Hydric Uplands

FSG No.: G138XA443FL

**Major Land Resource Area (MLRA 138):** North-Central Florida Ridge

#### Map Unit List

Bivans loamy sand, 8 to 12 percent slopes

#### Adapted Species List

The native forage species listed are considered adapted to grow on the soils in this group at their natural pH levels. All introduced grass and legume species will need the pH level raised to min. 5.5 (unless noted) for best production. All forages listed are adapted to dryland conditions. Consult with state extension service for current cultivar or germplasm recommendations (<http://agronomy.ifas.ufl.edu/foragesofflorida/>).

#### Perennial Species:

##### Grasses

Warm season (Introduced)

- Bahiagrass (*Paspalum notatum*, pH 5.0 – 6.5)
- Bermudagrass (*Cynodon dactylon*)

Warm season (Native)

- Big Bluestem (*Andropogon gerardii*)
- Purple Bluestem (*Andropogon glomeratus* var. *glaucopsis*)
- Yellow Indiangrass (*Sorghastrum nutans*)
- Lopsided Indiangrass (*Sorghastrum secundum*)
- Switchgrass (*Panicum virgatum*)

##### Legumes

Warm season (Introduced)

- Rhizoma Perennial Peanut (*Arachis glabrata*, pH 5.8-7.0)

#### Annual Species:

##### Grasses

Warm season

- Browntop Millet (*Urochloa ramosa*; =*Panicum ramosum*)
- Pearl Millet (*Pennisetum glaucum*)
- Sorghum (*Sorghum bicolor*; includes forage sorghums, sudangrass, and their hybrids)

Cool season

- Ryegrass, annual (*Lolium perenne* ssp. *multiflorum*; =*L. multiflorum*)

- Oat (*Avena sativa*)
- Rye (*Secale cereale*)
- Wheat (*Triticum aestivum*)
- Triticale (x *Triticosecale*)

##### Legumes

Warm season

- Aeschynomene (*Aeschynomene americana*)
- Hairy Indigo (*Indigofera hirsuta*)

Cool season

- White Clover (*Trifolium repens*, pH 6.0-7.5)
- Berseem Clover (*Trifolium alexandrinum*, pH 6.5-8.0)
- Ball Clover (*Trifolium nigrescens*, pH >6.5)

#### Seasonal and Total Production Estimates

Production on in this FSG would be expected to be similar to FSG G138XA341FL most times of the year due to loam and/or clay in the subsoil. The water table will perch above the loamy layer and seepage slopes will be present on sites with slopes >2%. Due to erosion concerns, annual forages should only be used when no-till is an option on those sites with 8% slope.

For this FSG, use of cool season forages such as annual ryegrass, oats, and wheat planted for use as a winter feed supply for the whole cow herd may be practical most years. In years of above average winter rainfall (El Niño winters), cool season annual grass forage growth may be limited on this FSG due to saturated soil conditions. Overseeding annual ryegrass also is an option.

Several clovers and other legumes are recommended for this FSG if planted no-till or by overseeding. Grazing management and fertilization need to favor the legume component for persistence, productivity, and seed production when natural reseeding of annuals is desired. Grazing management for seed production also is important for white clover, which is a weak a perennial in Florida and are heavily dependent upon reseeding to persist. Due to bloat issue, clovers should be used only in mixtures with cool season grasses, overseeded on bahiagrass pastures when grazed, or when bloat preventative supplements are fed.

Initial growth of perennial warm season grasses and legumes or establishment of warm season annual grasses may be delayed in the spring due to low rainfall. Often production of perennial species also dips during the

April/May dry period. Once normal summer rainfall begins, plant production should resume. Warm season legumes such as aescynomene can also be oversown onto warm season grasses in this forage suitability group, although fertilization (no N fertilizer) and grazing man-

agement needs to favor legume establishment and persistence. Additional lime may be needed to maintain a pH of 5.5 to 6.0.


<b>Expected Range in Dry Matter Production and Animal Unit Months (AUM) for Different Forages†</b>				
<b>Forage</b>	<b>Range in Dry Matter,</b>		<b>Range in AUM/acre‡</b>	
Bahiagrass (0 lb N/acre) <sup>13,14#</sup>	3,750	6,000	2.4	3.8
Bahiagrass (60 lb N/acre) <sup>14</sup>	8,750	10,000	5.6	6.4
White clover/bahiagrass <sup>13</sup>	7,500	8,000	4.8	5.1
Bermudagrass, (200 lb N/acre) <sup>11</sup>	12,500	14,000	8.0	9.0
Eastern Gamagrass, Pete (100-300 lb N/A) <sup>5,6,7</sup>	4,000	6,750	2.5	4.3
Big Bluestem (100-300 lb N/acre) <sup>5,6,7</sup>	900	1,800	0.6	1.2
Pearl Millet (225 to 300 lb N/acre) <sup>1,8</sup>	7,500	12,000	4.8	7.7
Sorghum X Sudangrass (225 to 300 lb N/acre) <sup>1,8</sup>	12,500	24,000	8.0	15.4
Ryegrass (120 lb N/A) <sup>3,4</sup>	3,500	7,200	2.2	4.6
Small Grain Forage (oat, wheat, etc.; 120 lb N/acre) <sup>2</sup>	6,000	7,200	3.8	4.6
Rhizoma Perennial Peanut <sup>12</sup>	10,000	14,000	6.4	9.0
Aeschynomene <sup>10</sup>	2,500	3,000	1.6	1.9
Hairy Indigo <sup>9</sup>	2,500	3,000	1.6	1.9
Cool-Season Clovers, overseeded on bahiagrass <sup>4,6,7</sup>	300	1,080	0.2	0.7
Cool-Season Clovers, prepared seedbed <sup>4,6</sup>	1,300	3,600	0.8	2.3

†Production data based on 25% increase in lower range values for FSG G138XA141FL for introduced warm season species and production similar to FSG G138XA231 for warm season natives and cool season species.

‡Animal Unit Month based on 50% grazing efficiency and 2.6% intake per day.

#Superscript numbers refer to references.

**Production Curves:**



Dry Matter Production Distribution by Month												
Forage	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<b>Introduced Warm Season Perennial Grasses</b>												
Bahiagrass (Pensacola)		0.01	0.03	0.14	0.14	0.17	0.19	0.17	0.14	0.03		
Bahiagrass (Argentine)		0.01	0.04	0.08	0.09	0.18	0.25	0.22	0.11	0.02		
Bermudagrass (Tifton 85)		0.02	0.02	0.13	0.15	0.13	0.21	0.22	0.08	0.03	0.01	0.00
<b>Native Warm Season Perennial Grasses</b>												
Native Warm Season Grasses (Generic)	0.01	0.02	0.03	0.05	0.1	0.2	0.27	0.19	0.07	0.03	0.02	0.01
Eastern Gamagrass	0.01	0.02	0.04	0.16	0.18	0.2	0.16	0.13	0.06	0.02	0.01	0.01
Switchgrass	0.01	0.02	0.03	0.07	0.15	0.19	0.2	0.19	0.09	0.03	0.01	0.01
<b>Legumes or Legume/Grass Combinations</b>												
Rhizoma Perennial Peanut			0.05	0.15	0.14	0.15	0.20	0.18	0.14			
White clover/Argentine Bahiagrass	0.01	0.02	0.07	0.14	0.17	0.21	0.18	0.12	0.09	0.02		
<b>Cool Season Annual Grasses</b>												
Annual Ryegrass	0.18	0.18	0.2	0.18	0.1						0.02	0.14
Small Grains (Wheat, Rye, etc.)	0.18	0.2	0.2	0.16	0.08						0.02	0.16
<b>Warm Season Annual Grasses</b>												
Sorghum-Sudangrass					0.07	0.2	0.3	0.25	0.15	0.03		
Millet (Pearl and Browntop)				0.07	0.15	0.26	0.39	0.07	0.04	0.02		

## Physiographic Features

Dominantly very deep, gently sloping to sloping, poorly drained soils formed in sandy over loamy, loamy, or clayey marine deposits. These soils are on summits, shoulders and back slopes of marine terraces. Diagnostic sub-surface horizon is an argillic horizon above 40 inches. The organic matter content of the surface layer is dominantly low or medium. Unless limed, the reaction in the surface layer ranges from extremely acid to moderately acid.

## Climatic Features

**Freeze-free period (>28° F 9 years in 10 at least):**  
 averages 270 d (range 266-278 d)

**Length of growing season (>32° F 9 years in 10 at least):** averages 234 d (range 225-247 d)

**Annual minimum temperature (° F in month of January):**  
 averages 41.1 (range 38.6-43.5)

**USDA Plant Hardiness Zone:**  
 8b (15-20° F, Jacksonville)  
 8a (10-15° F, Glen St. Mary)

**Mean annual precipitation (inches):**  
 averages 54.40 (range 52.24-59.65)

## Soil Properties

**Percent Slope:** Dominantly 2 to 12 percent

**Surface Texture:** Dominantly loamy sand, sand, or their gravelly analogs. A few members are fine sand, coarse sand, fine sandy loam, or loamy fine sand.

**Sand Content of Surface Layer:** 70 to 97 percent

**Clay Content of Surface Layer:** 1 to 15 percent

**Organic Matter Content of Surface Layer:** 0.5 to 10 percent

**Cation Exchange Capacity of Surface Layer (meq/100g):**  
 Not available

**Effective Cation Exchange Capacity of Surface Layer (meq/100g):** 1.1 to 5.5

**Bulk Density of Surface Layer (g/cc):** 1.3 to 1.53

**Saturated Hydraulic Conductivity of Surface Layer:**  
 Rapid

**Soil Reaction of Surface Layer:** 3.5 to 6.0 (unless limed)

**Available Water Capacity (0 to 30 inches):** 0.3 to 2.4 inch per inch

**Depth to Finer Textured Material:** Less than 40 inches to more than 80 inches

**Depth to Bedrock:** Greater than 80 inches

**Drainage Class (Agronomic):** Poorly

**Depth to Seasonal High Water Table (during wet periods):** 0 to 1 feet

**Flooding:** None

**Ponding:** None

## Monthly precipitation (inches) and temperature (F):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<b>Precip avg</b>	4.89	3.82	5.09	3.28	3.27	6.23	6.84	7.12	4.86	2.98	2.69	3.34
<b>Avg Min</b>	41.1	43.6	49.4	54.3	62.0	68.5	71.1	70.8	67.9	57.7	50.0	43.2
<b>Avg Temp</b>	55.1	56.3	62.3	67.5	74.4	79.6	81.6	81.2	78.4	69.8	62.4	55.4
<b>Avg Max</b>	65.7	68.9	75.2	80.5	86.8	90.7	92.0	91.5	88.9	82.1	74.7	67.6

## Climate Station Locations (averages from 1971 to 2000; see Appendix 1)

## FSG Documentation

### Inventory Data References:


1. Adjei, M.B., and P. Mislevy. 1988. Forage Yield and Quality of Summer Annual Grasses and Legumes in Central Florida. Univ. Florida, Range Cattle AREC. Research Rep. RC-1988-3. 6 p.
2. Barnett, R.D., D.L. Wright, A.R. Soffes Blount, and R.L. Stanley. 1997. Small Grain Production Recommendations for the 1997-98 Growing Season. Univ. Florida, IFAS, Florida Coop. Ext. Ser. SS-AGR-46.
3. Dunavin, L.S. 1997. Cool-Season Forage Trials, 1996-1997. Univ. Florida, IFAS, Florida Agric. Exp. Stn., WFREC Res. Rep. WF97-5.
4. -----, and C.G. Chambliss. 2000. Cool-Season Forage Variety Trials, WFREC, Jay, FL 1999-2000. Univ. Florida, IFAS, Florida Exp. Stn. SS-AGR-85.
5. -----, and D.W. Gorbet. 2000. Variety and Other Trials of Several Forage Grasses and Legumes, Temperate Corn and Grain Sorghum. Univ. Florida, IFAS, Florida Agric. Exp. Stn., WFREC Res. Rep. WF00-03.
6. -----, 2001. Variety and Other Trials of Several Forage Grasses and Legumes, Temperate Corn and Grain Sorghum. Univ. Florida, IFAS, Agric. Exp. Stn., WFREC Res. Rep. WF01-03.
7. -----, 2002. Variety and Other Trials of Several Forage Grasses and Legumes and Grain Sorghum. Univ. Florida, IFAS, Florida Agric. Exp. Stn., WFREC Res. Rep. WF02-03.
8. Green, V.E., Jr., D.W. Gorbet, L.S. Dunavin, Jr., H.A. Peacock, J.T. Johnson, R.S. Kalmbacher, C.G. Chambliss, R.J. Allen, Jr., G.M. Prine, A.M. Akhanda, B.R. Tyree, and P.H. Everett. 1978. Statewide uniform tests with grain sorghums, silage sorghums, annual summer grasses (sorghum x sudangrasses and pearl millets) and sweet sorghums (sorgos). Dep. Agron., Univ. Florida. Agron. Res. Rep. AG 78-7. 95 p.
9. Kalmbacher, R.S., E.M. Hodges, F.G. Martin. 1980. Effect of Plant and Cutting Height on Yield and Quality of *Indigofera hirsuta*. Tropical Grassld. 14:14-18.
10. Mislevy, P., R.S. Kalmbacher, and F.G. Martin. 1981. Cutting Management of the Tropical Legume American Jointvetch. Agron. J. 73:771-775 (<https://www.agronomy.org/publications/aj/abstracts/73/5/AJ0730050771>, accessed September 29, 2011).
11. Mislevy P., O.P. Miller, F.G. Martin. 2008. Influences of Grazing on *Cynodon* Grasses in Peninsular Florida., Online. Forage and Grazinglands doi; 10.1094/FG-2008-0429-01-RS. (<http://www.plantmanagementnetwork.org/pub/fg/research/2008/cynodon>; accessed August 19, 2011).
12. Mislevy, P., M.J. Williams, A.S. Blount, and K.H. Quesenberry. 2007. Influence of Harvest Management on Rhizoma Perennial Peanut Production, Nutritive Value, and Persistence on Flatwoods Soils. Online. Forage and Grazinglands. doi:10.1094/FG-2007-1108-01-RS. (<http://www.plantmanagementnetwork.org/pub/fg/research/2007/harvest/>; accessed August 19, 2011).
13. Prine, G.M., C. Ruelke, C.E. Dean, and D.D. Baltensperger. 1984. Performance of Winter Forage Legumes at Gainesville, Florida Over Eight Growing Seasons 1975-76 to 1982-83. Univ. Florida, IFAS. Agron. Res. Rep. AY 84-10. 32 p.
14. Sumner S., W. Wayne, J. Selph, J. Southwell, V. Hoge, P. Hogue, E. Jennings, P. Miller, and T. Seawright. 1991. Fertilization of Established Bahiagrass Pasture in Florida. Univ. Florida, IFAS, Agric. Exp. Stn., Cir. 916. (<http://rcrec.ona.ifas.ufl.edu/pdf/publications/fertilization-established-bahia-pasture-FL.pdf>; accessed August 20, 2011).

### State Correlation: (NA)

### Forage Suitability Group Approval:



Greg Hendricks, State Resource Conservationist



Tom Weber, State Soil Scientist

<b>Appendix 1: Climate Station Locations</b>		
<b>COOP ID (FL=08)</b>	<b>Location</b>	<b>County</b>
4731	Lake City	Columbia
4394	Jasper	Hamilton
5539	Mayo	LaFayette
9120	Usher Tower	Levy
5275	Madison	Madison
5099	Live Oak	Suwannee