

FORAGE SUITABILITY GROUP

Sandy Soils on Ridges and Dunes of Xeric Uplands

FSG No.: G152AA111FL

Major Land Resource Area (MLRA 152A):

Eastern Gulf Coast Flatwoods

Soil Series List

Due to the large list of map units in this group, please refer to Appendix 1.

Alpin	Lakeland
Astatula	Orlando
Candler	Paola
Apopka	Penney
Kershaw	Troup
Kureb	

Adapted Species List

The native forage species listed are considered adapted to grow on the soils in this group at their natural pH levels. All introduced grass and legume species will need the pH level raised to min. 5.5 (unless noted) for best production. All forages listed are adapted to dryland conditions. Consult with state extension service for current cultivar or germplasm recommendations (<http://agronomy.ifas.ufl.edu/foragesofflorida/>).

Perennial Species:

Grasses

Warm season (Introduced)

- Bahiagrass (*Paspalum notatum*; pH 5.0-6.5)
- Bermudagrass, (*Cynodon dactylon*)

Warm season (Native)

- Chalky Bluestem (*Andropogon virginicus* var. *glaucus*)
- Splitbeard Bluestem (*Andropogon ternarius*)
- Yellow Indiangrass (*Sorghastrum nutans*)
- Switchgrass (*Panicum virgatum*)

Legumes

Warm season (Introduced)

- Rhizoma Perennial Peanut (*Arachis glabrata*; pH 5.8-7.0)

Annual Species:

Grasses

Warm season (Introduced)

- Browntop Millet (*Urochloa ramosa*; =*Panicum ramosum*)
- Pearl Millet (*Pennisetum glaucum*)
- Sorghum (*Sorghum bicolor*; includes forage sorghum, sudangrass, and their hybrids)

Cool season (Introduced)

- Rye (*Secale cereale*)

Legumes

Warm season (Introduced)

- Alyceclover (*Alysicarpus vaginalis*)
- Cowpea (*Vigna unguiculata*)
- Hairy Indigo (*Indigofera hirsuta*)

Seasonal and Total Production Estimates

Seasonal and total forage production is limited in this group due to soils being well drained to excessively drained. Surface and subsurface texture is predominantly sandy with low or very low available water capacity, and a seasonal high water table greater than 6 feet. These factors increase drought effects. Total annual production is driven largely by rainfall; yields can increase by > 1,000 lbs/acre in years with above average rainfall. However greatly reduced production and even stand loss associated with over grazing can occur in years with below average rainfall. Irrigation is not recommended for these soils due to poor water holding capacity. Establishment of both annual and perennial warm season forages maybe delayed due to limited rainfall in the spring and short term drought periods in the summer months. Total production of all forage species is expected to be considerably less than other FSG, with a general growth curve weighted more towards the later part of the growing season.

Rye is the only cool season forage recommended for this FSG. Productivity of other cool season annuals will be very low without irrigation due to poor water holding capacity of the soils in this FSG.



Expected Range in Dry Matter Production and Animal Unit Months (AUM) for Different Forages				
Forage	Range in Dry Matter Yield, lb/acre		Range in AUM/acre†	
Bahiagrass (0 lb N/acre) ^{6‡}	2,500	3,500	1.6	2.2
Bahiagrass (60 lb N/acre) ^{6,10}	4,000	6,000	2.6	3.8
Bermudagrass (400 lb N/acre) ⁵	14,000	20,000	9.0	12.8
Switchgrass, Alamo ¹	6,000	8,000	3.8	5.1
Rhizoma Perennial Peanut, Florigraze ^{4,8}	7,000	10,000	4.5	6.4
Pearl Millet (limited irrigation, ≈400 lb N/acre) ⁷	8,000	16,000	5.1	10.3
Rye (120 lb N/acre) ^{3#}	3,600	4,800	2.3	3.1
Alyceclover ⁹	3,000	5,000	1.9	3.2
Hairy Indigo ²	6,000	12,000	3.8	7.7

†Animal Unit Month based on 50% grazing efficiency and 2.6% intake per day.

‡Superscript numbers refer to references.

#Production data based on 40% reduction yield range of FSG G152AA321FL.

Production Curves:



Physiographic Features

Dominantly very deep, nearly level to sloping, well to excessively drained soils formed in eolian or sandy marine deposits. These soils occur on summits, shoulders, and back slopes of marine terraces. They have 40 inches to greater than 80 inches of fine sand or sand. Diagnostic subsurface horizon is an argillic horizon below 40 inches or is absent. The organic matter content of the surface layer is dominantly very low or low. Unless limed, the reaction in the surface layer ranges from extremely acid to slightly acid.

Climatic Features

Freeze-free period (>28° F 9 years in 10 at least):
 averages 268 d (range 243-295 d)

Length of growing season (>32° F 9 years in 10 at least): averages 235 d (range 206-267 d)

Annual minimum temperature (° F in month of January):
 averages 53.7 (range 49.0-64.5)

USDA Plant Hardiness Zone:
 8b (15-20° F, Tallahassee)
 9a (20-25° F, Gainesville)

Mean annual precipitation (inches):
 averages 60.77 (range 52.83-69.20)

Soil Properties

Percent Slope: 0 to 8 percent

Surface Texture: Fine sand, sand, loamy fine sand, loamy sand, gravelly sand

Sand Content of Surface Layer: 81 to 99 percent

Clay Content of Surface Layer: 0.1 to 10 percent

Organic Matter Content of Surface Layer: 0.5 to 5 percent

Cation Exchange Capacity of Surface Layer (meq/100g):
 0.3 to 7.1

Effective Cation Exchange Capacity of Surface Layer (meq/100g): 0.1 to 6.7

Bulk Density of Surface Layer (g/cc): 1.3 to 1.7

Saturated Hydraulic Conductivity of Surface Layer: Rapid or very rapid

Soil Reaction of Surface Layer: 3.5 to 6.5 (unless limed)

Available Water Capacity (0 to 30 inches): 0.1 to 1.0 inch per inch

Depth to Finer Textured Material: 40 to more than 80 inches

Depth to Bedrock: Greater than 80 inches

Drainage Class (Agronomic): Well, Somewhat excessive, Excessive

Depth to Seasonal High Water Table (during wet periods): More than 6 feet

Flooding: None

Ponding: None

Monthly precipitation (inches) and temperature (F):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip avg	5.25	4.28	5.67	3.52	3.78	6.20	7.82	7.44	5.82	3.46	3.54	3.75
Avg Min	40.5	43.0	48.9	83.7	62.0	66.9	71.6	71.4	68.1	57.1	49.0	43.2
Avg Temp	53.7	56.7	62.7	68.1	75.3	80.8	82.6	82.3	79.4	70.6	62.6	55.8
Avg Max	64.1	67.3	73.3	79.1	85.6	90.1	91.3	90.9	88.3	81.2	73.3	66.2

Climate Station Locations (averages from 1971 to 2000; see Appendix 2)

FSG Documentation

Inventory Data References:

1. -----, 2000. Plant materials adaption study final report, Belleview-Perry sprayfield. Brooksville, FL, USA: US Department of Agriculture, Natural Resources Conservation Service, Brooksville Plant Material Center Brooksville, Florida. 25 p.
2. Baltensperger, D.D., E.C. French, G.M. Prine, O.C. Ruelke, and K.H. Quesenberry. Hairy indigo a summer legume for Florida. Gainesville, FL, USA: University of Florida, Institute of Food and Agricultural Sciences, Agriculture Experiment Station, Circular S-318. 12 p. Available at: <http://ufdc.ufl.edu/UF00055246/00001>. Accessed 17 December 2012.
3. Barnett, R.D., D.L. Wright, A.R. Soffes Blount, and R.L. Stanley. 1997. Small grain production recommendations for the 1997-98 growing season. Gainesville, FL, USA: University of Florida, Institute of Food and Agriculture S, Florida Cooperative Extension Service SS-AGR-46.
4. Beltranena, R., J. Breman, and G. Prine. 1981. Yield and quality of Florigraze rhizoma peanut (*Arachis glabrata* Benth.) as affected by cutting height and frequency. Proceedings Soil Crop Science Society Florida 40:153-156.
5. Coleman, S., and M. Williams. 2007. Bermudagrass yield and quality through the grazing season. In: Proceedings USDA-ARS STARS Field Day; 25 May 2007; Brooksville, FL, USA. Brooksville, FL, USA: US Department of Agriculture, Agriculture Research Service.
6. Engibous, J.C., W.J. Friedmann, and M.B. Gillis. 1958. Yield and quality of pangolagrass and bahiagrass as affected by rate and frequency of fertilization. Soil Science Society of America Journal. 22:423-425. Available at: <https://www.agronomy.org/publications/sssaj/abstracts/22/5/SS0220050423>. Accessed 17 December 2012.
7. Green, V.E., Jr., D.W. Gorbet, L.S. Dunavin, Jr., H.A. Peacock, J.T. Johnson, R.S. Kalmbacher, C.G. Chambliss, R.J. Allen, Jr., G.M. Prine, A.M. Akhanda, B.R. Tyree, and P.H. Everett. 1978. Statewide uniform tests with grain sorghums, silage sorghums, annual summer grasses (sorghum x sudangrasses and pearl millets) and sweet sorghums (sorgos). Gainesville, FL, USA: Department of Agronomy, University of Florida, Agronomy Research Report AG 78-7. 95 p.
8. Williams, M.J. 1994. Growth characteristics of rhizoma peanut and nitrogen-fertilized bahiagrass swards. Agronomy Journal 86:819-823. Available at: <https://www.agronomy.org/publications/aj/abstracts/86/5/AJ0860050819>. Accessed 17 December 2012.
9. Williams, M.J., C.G. Chambliss, and J.D. Brolmann. 1993. Potential of 'Savanna' stylo as a stockpiled forage for the subtropical USA. Journal of Production Agric. 6:553-556.
10. Williams, M.J., and R.S. Kalmbacher. 1996. Renovation effects on bahiagrass productivity. Agronomy Journal 88:191-198. Available at: <https://www.agronomy.org/publications/aj/abstracts/88/2/AJ0880020191>. Accessed 17 December 2012.

State Correlation: (NA)

Forage Suitability Group Approval:



Greg Hendricks, State Resource Conservationist



Tom Weber, State Soil Scientist

Appendix 1. Map Unit List

Alpin sand, 0 to 5 percent slopes	Lakeland sand, 5 to 8 percent slopes
Astatula fine sand, 1 to 8 percent slopes	Orlando fine sand, 1 to 5 percent slopes
Candler fine sand, 1 to 5 percent slopes	Orlando fine sand, 1 to 5 percent slopes
Candler fine sand, 5 to 8 percent slopes	Orlando fine sand, 5 to 8 percent slopes
Candler-Apopka complex, 1 to 5 percent slopes	Paola fine sand, gently rolling
Kershaw fine sand, 0 to 8 percent slopes	Penney fine sand, 0 to 5 percent slopes
Kershaw sand, 0 to 5 percent slopes	Penney sand, 0 to 5 percent slopes
Kershaw sand, 2 to 5 percent slopes	Penney sand, 5 to 8 percent slopes
Kureb fine sand, 3 to 8 percent slopes	Penney-Wadley complex, 0 to 5 percent slopes
Kureb sand, 0 to 5 percent slopes	Troup sand, 0 to 5 percent slopes
Kureb sand, 2 to 5 percent slopes	Troup sand, 5 to 8 percent slopes
Lakeland sand, 0 to 5 percent slopes	

Appendix 2: Climate Station Locations		
COOP ID (FL=08)	Location	County
3230	Fountain	Bay
6842	Panama City	Bay
2008	Cross City	Dixie
3855	Pensacola Sherman NAS	Escambia
6997	Pensacola Regional Air.	Escambia
211	Apalachicola	Franklin
9566	Wewahitchka	Gulf
5539	Mayo	LaFayette
8758	Tallahassee Mun. Air.	Leon
9120	Usher Tower	Levy
5275	Madison	Madison
6240	Niceville	Okaloosa
3841	Whiting Field NAS	Santa Rosa
5099	Live Oak	Suwannee
7025	Perry	Taylor
8565	Steinhatchee	Taylor