

## FORAGE SUITABILITY GROUP

### Sandy Soils on Rises, Knolls, and Ridges of Mesic Uplands

FSG No.: G155XB121FL

**Major Land Resource Area (MLRA 155):** Southern Florida Flatwoods

#### Soil Series List

Due to the large list of map units in this group, please refer to Appendix 1.

Archbold	Hobe
Cassia moderately well	Jonathan
Centenary	Millhopper
Daytona	Orlando
Duette	Orsino
Florahome	Tavares

#### Adapted Species List

The native forage species listed are considered adapted to grow on the soils in this group at their natural pH levels. All introduced grass and legume species will need the pH level raised to min. 5.5 (unless noted) for best production. Irrigation is not recommended in these soils, and all forages listed are adapted to dryland conditions. Consult with state extension service for current cultivar or germplasm recommendations (<http://agronomy.ifas.ufl.edu/foragesofflorida/>).

#### Perennial Species:

##### Grasses

###### Warm season (Introduced)

- Bahiagrass (*Paspalum notatum*; pH 5.0-6.5)
- Bermudagrass (*Cynodon dactylon*)

###### Warm season (Native)

- Chalky Bluestem (*Andropogon virginicus* var. *glaucus*)
- Big Bluestem (*Andropogon gerardii*, northern half of the MLRA)
- Splitbeard Bluestem (*Andropogon ternarius*)
- Yellow Indiangrass (*Sorghastrum nutans*, northern half of MLRA)
- Switchgrass (*Panicum virgatum*)

##### Legumes

###### Warm season (Introduced)

- Rhizoma Perennial Peanut (*Arachis glabrata*; pH 5.8-7.0)

#### Annual Species:

##### Grasses

###### Warm season (Introduced)

- Browntop Millet (*Urochloa ramosa*; =*Panicum ramosum*)
- Pearl Millet (*Pennisetum glaucum*)
- Sorghum (*Sorghum bicolor*; includes forage sorghum, sudangrass, and their hybrids)

##### Legumes and Forbs

###### Warm season (Introduced)

- Alyceclover (*Alysicarpus vaginalis*)
- Cowpea (*Vigna unguiculata*)
- Hairy Indigo (*Indigofera hirsuta*)

#### Seasonal and Total Production Estimates

Seasonal and total forage production is somewhat higher than FSG G155XB111FL because soils in this FSG have slightly better water holding capacity and seasonal water table is higher (3 to 5 feet). These factors will decrease drought effects, but total annual production still is driven largely by rainfall. Yields can increase by > 1,000 lbs/acre in years with above average rainfall. However greatly reduced production and even stand loss associated with over grazing can occur in years with below average rainfall. Irrigation is not recommended for these soils due to poor water holding capacity. Establishment of both annual and perennial warm season forages maybe delayed due to limited rainfall in the spring and short term drought periods in the summer months. Total production of all forage species is expected to be higher than FSG G155XB111FL but less than other groups, with a general growth curve still weighted more towards the later part of the growing season.

Cool season forage production is very limited due to decreased and sporadic rainfall during winter months (November-March), therefore no cool season forages are recommended and no production data is given.


Expected Range in Dry Matter Production and Animal Unit Months (AUM) for Different Forages†				
Forage	Range in Dry Matter Yield, lb/acre		Range in AUM/acre‡	
	Bahiagrass (0 lb N/acre) <sup>5</sup> #	3,100	4,400	2.0
Bahiagrass (60 lb N/acre) <sup>5,9</sup>	5,000	7,500	3.2	4.8
Bermudagrass (400 lb N/acre) <sup>4</sup>	17,500	25,000	11.2	16.0
Switchgrass, Alamo <sup>1</sup>	7,500	10,000	4.8	6.4
Rhizoma Perennial Peanut, Florigraze <sup>3,7</sup>	8,750	12,500	5.6	8.0
Pearl Millet (limited irrigation, ~400 lb N/acre) <sup>6</sup>	10,000	20,000	6.4	12.8
Alyceclover <sup>8</sup>	3,750	6,250	2.4	4.0
Hairy Indigo <sup>2</sup>	7,500	15,000	4.8	9.6

† Production data based on a 25% increase from FSG G155XB111FL.

‡ Animal Unit Month based on 50% grazing efficiency and 2.6% intake per day.

#Superscript numbers refer to references.

**Production Curves:**



## Physiographic Features

Dominantly very deep, nearly level to gently sloping, well drained or moderately well drained soils formed in sandy marine deposits. These soils are on summits, shoulders, and back slopes of marine terraces. They have 40 inches to greater than 80 inches of fine sand or sand. Diagnostic subsurface horizon is an argillic horizon below 40 inches or is absent. The organic matter content of the surface layer is dominantly very low or low. Unless limed, the reaction in the surface layer ranges from extremely acid to slightly acid.

## Climatic Features

**Freeze-free period (>28° F 9 years in 10 at least):**  
 averages 337 d (range 290-365 d)

**Length of growing season (>32° F 9 years in 10 at least):** averages 309 d (range 253-365 d)

**Annual minimum temperature (° F in month of January):**  
 average 50.2 (range 45.2-59.2)

**USDA Plant Hardiness Zone:**  
 9a (20-25° F, Ocala)  
 9b (25-30° F, Orlando)  
 10a (30-35° F, Ft. Myers)

**Mean annual precipitation (inches):**  
 averages 51.89 (range 45.66-69.53)

## Soil Properties

**Percent Slope:** Dominantly 0 to 8 percent, but ranges to 10 percent

**Surface Texture:** Fine sand, sand, coarse sand, loamy sand, very fine sand

**Sand Content of Surface Layer:** 84 to 99 percent

**Clay Content of Surface Layer:** 0.1 to 8 percent

**Organic Matter Content of Surface Layer:** 0.5 to 3 percent

**Cation Exchange Capacity of Surface Layer (meq/100g):**  
 0.1 to 5.3

**Effective Cation Capacity of Surface Layer (meq/100g):**  
 0.1 to 5.8

**Bulk Density of Surface Layer (g/cc):** 1.3 to 1.6

**Saturated Hydraulic Conductivity of Surface Layer:** Rapid or very rapid

**Soil Reaction of Surface Layer:** 3.5 to 6.5 (unless limed)

**Available Water Capacity (0 to 30 inches):** 0.3 to 1.9 inch per inch

**Depth to Finer Textured Material:** 40 to more than 80 inches

**Depth to Bedrock:** Greater than 80 inches. A few members have bedrock at 60 to 80 inches.

**Drainage Class (Agronomic):** Moderately well, Well

**Depth to Seasonal High Water Table (during wet periods):** 3.0 to 5.0 feet below the surface

**Flooding:** None

**Ponding:** None

## Monthly precipitation (inches) and temperature (F):

	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
<b>Precip avg</b>	2.70	2.59	3.37	2.39	3.90	7.26	6.98	7.14	6.75	3.50	2.66	2.24
<b>Avg Min</b>	50.2	51.4	55.7	59.6	65.5	70.8	72.3	72.7	71.6	63.9	58.9	53.0
<b>Avg Temp</b>	62.3	63.5	67.8	70.5	77.1	81.1	82.0	82.3	81.1	75.8	69.6	63.9
<b>Avg Max</b>	72.7	74.4	78.6	82.7	87.5	90.2	91.5	91.3	89.5	84.8	79.2	74.0

## Climate Station Locations (averages from 1971 to 2000; see Appendix 2)

## FSG Documentation

### Inventory Data References:

1. ----- . 2000. Plant materials adaption study final report, Belleview-Perry sprayfield. Brooksville, FL, USA: US Department of Agriculture, Natural Resources Conservation Service, Brooksville Plant Material Center Brooksville, Florida. 25 p.
2. Baltensperger, D.D., E.C. French, G.M. Prine, O.C. Ruelke, and K.H. Quesenberry. Hairy indigo a summer legume for Florida. Gainesville, FL, USA: University of Florida, Institute of Food and Agricultural Sciences, Agriculture Experiment Station, Circular S-318. 12 p. Available at: <http://ufdc.ufl.edu/UF00055246/00001>. Accessed 17 December 2012.
3. Beltranena, R., J. Breman, and G. Prine. 1981. Yield and quality of Florigrass rhizoma peanut (*Arachis glabrata* Benth.) as affected by cutting height and frequency. Proceedings Soil Crop Science Society Florida 40:153-156.
4. Coleman, S., and M. Williams. 2007. Bermudagrass yield and quality through the grazing season. In: Proceedings USDA-ARS STARS Field Day; 25 May 2007; Brooksville, FL, USA. Brooksville, FL, USA: US Department of Agriculture, Agriculture Research Service.
5. Engibous, J.C., W.J. Friedmann, and M.B. Gillis. 1958. Yield and quality of pangolagrass and bahiagrass as affected by rate and frequency of fertilization. Soil Science Society of America Journal. 22:423-425. Available at: <https://www.agronomy.org/publications/ssaj/abstracts/22/5/SS0220050423>. Accessed 17 December 2012.
6. Green, V.E., Jr., D.W. Gorbet, L.S. Dunavin, Jr., H.A. Peacock, J.T. Johnson, R.S. Kalmbacher, C.G. Chambliss, R.J. Allen, Jr., G.M. Prine, A.M. Akhanda, B.R. Tyree, and P.H. Everett. 1978. Statewide uniform tests with grain sorghums, silage sorghums, annual summer grasses (sorghum x sudangrasses and pearl millets) and sweet sorghums (sorgos). Gainesville, FL, USA: Department of Agronomy, University of Florida, Agronomy Research Report AG 78-7. 95 p.
7. Williams, M.J. 1994. Growth characteristics of rhizoma peanut and nitrogen-fertilized bahiagrass swards. Agronomy Journal 86:819-823. Available at: <https://www.agronomy.org/publications/aj/abstracts/86/5/AJ0860050819>. Accessed 17 December 2012.
8. Williams, M.J., C.G. Chambliss, and J.D. Brolmann. 1993. Potential of 'Savanna' stylo as a stockpiled forage for the subtropical USA. Journal of Production Agric. 6:553-556.
9. Williams, M.J., and R.S. Kalmbacher. 1996. Renovation effects on bahiagrass productivity. Agronomy Journal 88:191-198. Available at: <https://www.agronomy.org/publications/aj/abstracts/88/2/AJ0880020191>. Accessed 17 December 2012.

State Correlation: (NA)

### Forage Suitability Group Approval:



Greg Hendricks, State Resource Conservationist



Tom Weber, State Soil Scientist

<b>Appendix 1: Map Unit List</b>	
Archbold fine sand	Millhopper sand, 0 to 5 percent slopes
Archbold fine sand, 0 to 5 percent slopes	Millhopper sand, 5 to 8 percent slopes
Archbold sand, 0 to 5 percent slopes	Orlando fine sand, moderately wet, 0 to 2 percent slopes
Cassia fine sand, moderately well drained	Orsino fine sand
Centenary fine sand	Orsino fine sand, 0 to 5 percent slopes
Daytona sand	Orsino fine sand, 0 to 8 percent slopes
Daytona sand, 0 to 5 percent slopes	Orsino sand, 0 to 5 percent slopes
Duette fine sand, 0 to 5 percent slopes	Tavares fine sand
Duette sand, 0 to 5 percent slopes	Tavares fine sand, 0 to 5 percent slopes
Florahome fine sand, 0 to 5 percent slopes	Tavares fine sand, cemented substratum, 2 to 5 percent slopes
Florahome sand	Tavares sand, 0 to 5 percent slopes
Hobe fine sand, 0 to 5 percent slopes	Tavares-Milhopper fine sands, 0 to 5 percent slopes
Jonathan fine sand	Tavares-Milhopper fine sands, 5 to 8 percent slopes
Jonathan sand	

<b>Appendix 2: Climate Station Locations</b>		
<b>COOP ID (FL=08)</b>	<b>Location</b>	<b>County</b>
8942	Titusville	Brevard
3163	Fort Lauderdale	Broward
7397	Punta Gorda	Charlotte
2850	Everglades	Collier
4210	Immokalee	Collier
228	Arcadia	DeSoto
5895	Moore Haven Lock	Glades
9401	Wauchula	Hardee
1654	Clewiston US Engin.	Hendry
2298	Devils Garden	Hendry
4662	La Belle	Hendry
236	Archbold Biol. Station	Highlands
369	Avon Park	Highlands
7205	Plant City	Hillsborough
8788	Tampa Intl. Air.	Hillsborough
9214	Vero Beach Muni. Air.	Indian River
9219	Vero Beach	Indian River
1641	Clermont	Lake
5076	Lisbon	Lake
3186	Fort Myers	Lee
6880	Parrish	Manatee
8620	Stuart	Martin
2137	Fort Drum	Okeechobee
6485	Okeechobee	Okeechobee
6628	Orlando Intl. Air.	Orange
4625	Kissimmee	Osceola
611	Belle Glade Exp. Stn.	Palm Beach
1276	Canal Point USDA	Palm Beach
5182	Loxahatchee	Palm Beach
9525	West Palm Beach Intl. Air.	Palm Beach
7851	St. Leo	Pasco
478	Bartow	Polk
4707	Lake Alfred Exp Stn	Polk
4797	Lakeland	Polk
5973	Mountain Lake	Polk
9707	Winter Haven	Polk
1978	Crescent City	Putnam
2915	Federal Point	Putnam
6753	Palatka	Putnam
6065	Myakka River State Park	Sarasota
9176	Venice	Sarasota
7982	Sanford Orlando	Seminole
3874	Hastings ARC	St. Johns
7826	St. Augustine WFOY	St. Johns
3207	Fort Pierce	St. Lucie
1163	Bushnell	Sumter
2158	Daytona Beach Inter. Air.	Volusia
2229	Deland	Volusia