## Technical Guide – Section II Correlation Date: April 2012 HEL List Amended: May 1, 2017

# HIGHLY ERODIBLE LAND Brantley County, Georgia

Map Symbol	Map Unit Name	HEL Class
AbA	Albany-Leefield complex, 0 to 3 percent slopes	NHEL
AqA	Aquents, ponded, 0 to 1 percent slopes	NHEL
BdA	Bladen loamy fine sand, 0 to 2 percent slopes, occasionally flooded	NHEL
ВоВ	Bonifay loamy sand, 0 to 5 percent slopes	NHEL
BrA	Brookman loam, 0 to 1 percent slopes, frequently flooded	NHEL
ChA	Centenary-Hurricane complex, 0 to 3 percent slopes	NHEL
CnB	Cowarts, moderately wet-Uchee complex, 2 to 5 percent slopes	PHEL
CnD	Cowarts, moderately wet-Uchee complex, 5 to 12 percent slopes	HEL
СРА	Croatan, Pamlico and Surrency soils, ponded, 0 to 1 percent slopes, rarely flooded	NHEL
DDA	Dasher-Dorovan-Croatan association, ponded, 0 to 1 percent slopes	NHEL
FaA	Florala loamy fine sand, 0 to 2 percent slopes	NHEL
FaB	Florala loamy fine sand, 2 to 5 percent slopes	NHEL
FoB	Foxworth sand, 0 to 5 percent slopes	NHEL
FuB	Fuquay loamy sand, 0 to 5 percent slopes	NHEL
HuA	Hurricane fine sand, 0 to 2 percent slopes	NHEL
KJA	Kinston and Johnston soils, 0 to 2 percent slopes, frequently flooded	NHEL
KOA	Kinston and Osier soils, 0 to 2 percent slopes, frequently flooded	NHEL
LaC	Lakeland sand, 0 to 8 percent slopes	NHEL
LdA	Leefield loamy sand, 0 to 3 percent slopes	NHEL
LeA	Leon fine sand, 0 to 2 percent slopes	NHEL
LoA	Leon fine sand, ponded, 0 to 2 percent slopes	NHEL
LvA	Lynn Haven fine sand, 0 to 2 percent slopes	NHEL
LYA	Lynn Haven, Allanton and Kingsferry soils, ponded, 0 to 1 percent slopes	NHEL
MaA	Mandarin fine sand, 0 to 2 percent slopes	NHEL
McA	Mascotte fine sand, 0 to 2 percent slopes	NHEL
MdA	Mascotte mucky fine sand, ponded, 0 to 1 percent slopes	NHEL
MeA	Meggett loamy fine sand, 0 to 1 percent slopes, frequently flooded	NHEL
MrB	Meldrim sand, 0 to 5 percent slopes	NHEL
ОрА	Ogeechee-Pelham complex, 0 to 3 percent slopes, occasionally flooded	NHEL
OsA	Olustee loamy fine sand, 0 to 2 percent slopes	NHEL
OuB	Ousley fine sand, 0 to 5 percent slopes, occasionally flooded	NHEL
PbA	Pottsburg fine sand, 0 to 2 percent slopes	NHEL
PhA	Pelham fine sand, ponded, 0 to 2 percent slopes	NHEL
PmB	Pelham loamy fine sand, 0 to 5 percent slopes	NHEL
Pt	Pits, borrow	
RhA	Ridgewood-Hurricane complex, 0 to 2 percent slopes	NHEL
RoA	Rigdon-Olustee complex, 0 to 2 percent slopes	NHEL
StA	Stilson loamy sand, 0 to 3 percent slopes	NHEL
SuA	Surrency mucky sand, ponded, 0 to 1 percent slopes	NHEL

- All HEL Classes were assigned using USLE factor values that were in effect as of January 1, 1990.
- This HEL list is based on the final correlation of survey data and supersedes any previous HEL list.

Technical Guide – Section II Correlation Date: April 2012 HEL List Amended: May 1, 2017

# HIGHLY ERODIBLE LAND Brantley County, Georgia

Map Symbol	Map Unit Name	HEL Class
TfB	Tifton loamy sand, 2 to 5 percent slopes	NHEL
TnA	Tifton-Florala complex, 0 to 2 percent slopes	NHEL
UdD	Udorthents, 0 to 10 percent slopes	*
UsE	Udorthents, sanitary landfill, 2 to 25 percent slopes	
W	Water	
WaA	Wahee-Ogeechee complex, 0 to 3 percent slopes, occasionally flooded	NHEL

<sup>\*</sup> Properties of this mapunit are highly variable. Site evaluation is needed to determine HEL Class.

HEL Highly Erodible Land

PHEL Potentially Highly Erodible Land

NHEL Not Highly Erodible Land

- All HEL Classes were assigned using USLE factor values that were in effect as of January 1, 1990.
- This HEL list is based on the final correlation of survey data and supersedes any previous HEL list.

#### **DOCUMENTATION FOR:**

Correlation Date: April 2012 HEL List Amended: May 1, 2017

Technical Guide – Section II

HIGHLY ERODIBLE LAND Brantley County, Georgia

### R Factor = 375

	Mapunit	Component									
Мар	<u>'</u>		% of				Min	Max	8T/	HEL	
Sym	Name	Name	MU	Slope	Т	K	LS	LS	RK	Class	
AbA	Albany-Leefield complex,	Albany	63	0-3	5	.1	0.11	0.3	1.07	NHEL	
	0 to 3 percent slopes	Leefield	15	0-3	5	.1	0.11	0.3	1.07	NHEL	
AqA	Aquents, ponded, 0 to 1 percent slopes	Aquents	98	0-1	5	.1	0.11	0.15	1.07	NHEL	
BdA	Bladen loamy fine sand, 0 to 2 percent slopes, occasionally flooded	Bladen	91	0-2	5	.1	0.11	0.23	1.07	NHEL	
ВоВ	Bonifay loamy sand, 0 to 5 percent slopes	Bonifay	75	0-5	5	.15	0.11	0.59	0.71	NHEL	
BrA	Brookman loam, 0 to 1 percent slopes, frequently flooded	Brookman	100	0-1	4	.24	0.11	0.15	0.36	NHEL	
ChA	Centenary-Hurricane complex, 0 to 3 percent slopes	Centenary	50	0-3	5	.1	0.11	0.3	1.07	NHEL	
ChA	Centenary-Hurricane complex, 0 to 3 percent slopes	Hurricane	50	0-3	5	.1	0.11	0.3	1.07	NHEL	
CnB	Cowarts, moderately wet-	Cowarts	47	2-5	3	.15	0.23	0.59	0.43	PHEL	
	Uchee complex, 2 to 5 percent slopes	Uchee	26	2-5	4	.1	0.23	0.59	0.85	NHEL	
CnD	Cowarts, moderately wet-	Cowarts	45	5-12	3	.15	0.59	1.61	0.43	HEL	
	Uchee complex, 5 to 12 percent slopes	Uchee	30	5-12	4	.1	0.59	1.61	0.85	PHEL	
CPA	Croatan, Pamlico and	Croatan	40	0-1	-					NHEL	
	Surrency soils, ponded, 0	Pamlico	30	0-1	-					NHEL	
	to 1 percent slopes, rarely flooded	Surrency	24	0-1	5	.1	0.11	0.15	1.07	NHEL	
DDA	Dasher-Dorovan-Croatan	Dasher	55	0-1	-					NHEL	
	association, ponded, 0 to	Dorovan	17	0-1	-					NHEL	
	1 percent slopes	Croatan	10	0-1	-					NHEL	
FaA	Florala loamy fine sand, 0 to 2 percent slopes	Florala	88	0-2	5	.15	0.11	0.23	0.71	NHEL	
FaB	Florala loamy fine sand, 2 to 5 percent slopes	Florala	84	2-5	5	.15	0.23	0.59	0.71	NHEL	
FoB	Foxworth sand, 0 to 5 percent slopes	Foxworth	84	0-5	5	.1	0.11	0.59	1.07	NHEL	
FuB	Fuquay loamy sand, 0 to 5 percent slopes	Fuquay	67	0-5	5	.15	0.11	0.59	0.71	NHEL	

- All HEL Classes were assigned using USLE factor values that were in effect as of January 1, 1990.
- This HEL list is based on the final correlation of survey data and supersedes any previous HEL list.

#### **DOCUMENTATION FOR:**

Correlation Date: April 2012 HEL List Amended: May 1, 2017

Technical Guide – Section II

HIGHLY ERODIBLE LAND Brantley County, Georgia

R Factor = 375

	Mapunit	Component								
Мар			% of				Min	Max	8T/	HEL
Sym	Name	Name	MU	Slope	Т	K	LS	LS	RK	Class
HuA	Hurricane fine sand, 0 to 2 percent slopes	Hurricane	87	0-2	5	.1	0.11	0.23	1.07	NHEL
KJA	Kinston and Johnston	Kinston	60	0-2	5	.24	0.11	0.23	0.44	NHEL
	soils, 0 to 2 percent slopes, frequently flooded	Johnston	40	0-2	5	.17	0.11	0.23	0.63	NHEL
KOA	Kinston and Osier soils, 0	Kinston	65	0-2	5	.24	0.11	0.23	0.44	NHEL
	to 2 percent slopes, frequently flooded	Osier	25	0-2	5	.1	0.11	0.23	1.07	NHEL
LaC	Lakeland sand, 0 to 8 percent slopes	Lakeland	90	0-8	5	.1	0.11	0.99	1.07	NHEL
LdA	Leefield loamy sand, 0 to 3 percent slopes	Leefield	80	0-3	5	.1	0.11	0.3	1.07	NHEL
LeA	Leon fine sand, 0 to 2 percent slopes	Leon	94	0-2	5	.1	0.11	0.23	1.07	NHEL
LoA	Leon fine sand, ponded, 0 to 2 percent slopes	Leon	92	0-2	5	.1	0.11	0.23	1.07	NHEL
LvA	Lynn Haven fine sand, 0 to 2 percent slopes	Lynn Haven	90	0-2	5	.1	0.11	0.23	1.07	NHEL
LYA	Lynn Haven, Allanton and	Lynn Haven	45	0-1	5	.1	0.11	0.15	1.07	NHEL
	Kingsferry soils, ponded, 0	Allanton	28	0-1	5	.1	0.11	0.15	1.07	NHEL
	to 1 percent slopes	Kingsferry	24	0-1	5	.1	0.11	0.15	1.07	NHEL
MaA	Mandarin fine sand, 0 to 2 percent slopes	Mandarin	92	0-2	5	.1	0.11	0.23	1.07	NHEL
McA	Mascotte fine sand, 0 to 2 percent slopes	Mascotte	81	0-2	5	.1	0.11	0.23	1.07	NHEL
MdA	Mascotte mucky fine sand, ponded, 0 to 1 percent slopes	Mascotte	95	0-1	5	.1	0.11	0.15	1.07	NHEL
MeA	Meggett loamy fine sand, 0 to 1 percent slopes, frequently flooded	Meggett	100	0-1	5	.24	0.11	0.15	0.44	NHEL
MrB	Meldrim sand, 0 to 5 percent slopes	Meldrim	75	0-5	5	.1	0.11	0.59	1.07	NHEL
ОрА	Ogeechee-Pelham	Ogeechee	76	0-3	5	.1	0.11	0.3	1.07	NHEL
	complex, 0 to 3 percent slopes, occasionally flooded	Pelham	20	0-3	5	.1	0.11	0.3	1.07	NHEL
OsA	Olustee loamy fine sand, 0 to 2 percent slopes	Olustee	94	0-2	5	.1	0.11	0.23	1.07	NHEL

- All HEL Classes were assigned using USLE factor values that were in effect as of January 1, 1990.
- This HEL list is based on the final correlation of survey data and supersedes any previous HEL list.

### **DOCUMENTATION FOR:**

Brantley County, Georgia

Correlation Date: April 2012 HEL List Amended: May 1, 2017

Technical Guide - Section II

HIGHLY ERODIBLE LAND

### R Factor = 375

	Mapunit	Component								
Мар	·		% of				Min	Max	8T/	HEL
Sym	Name	Name	MU	Slope	Т	K	LS	LS	RK	Class
OuB	Ousley fine sand, 0 to 5 percent slopes, occasionally flooded	Ousley	93	0-5	5	.1	0.11	0.59	1.07	NHEL
PbA	Pottsburg fine sand, 0 to 2 percent slopes	Pottsburg	85	0-2	5	.1	0.11	0.23	1.07	NHEL
PhA	Pelham fine sand, ponded, 0 to 2 percent slopes	Pelham	96	0-2	5	.1	0.11	0.23	1.07	NHEL
PmB	Pelham loamy fine sand, 0 to 5 percent slopes	Pelham	95	0-5	5	.1	0.11	0.59	1.07	NHEL
Pt	Pits, borrow	Pits	100		-					
RhA	Ridgewood-Hurricane	Ridgewood	60	0-2	5	.1	0.11	0.23	1.07	NHEL
	complex, 0 to 2 percent slopes	Hurricane	27	0-2	5	.1	0.11	0.23	1.07	NHEL
RoA	Rigdon-Olustee complex,	Rigdon	70	0-2	5	.1	0.11	0.23	1.07	NHEL
	0 to 2 percent slopes	Olustee	30	0-2	5	.1	0.11	0.23	1.07	NHEL
StA	Stilson loamy sand, 0 to 3 percent slopes	Stilson	75	0-3	5	.1	0.11	0.3	1.07	NHEL
SuA	Surrency mucky sand, ponded, 0 to 1 percent slopes	Surrency	97	0-1	5	.1	0.11	0.15	1.07	NHEL
TfB	Tifton loamy sand, 2 to 5 percent slopes	Tifton	76	2-5	4	.1	0.23	0.59	0.85	NHEL
TnA	Tifton-Florala complex, 0	Tifton	64	0-2	4	.1	0.11	0.23	0.85	NHEL
	to 2 percent slopes	Florala	24	0-2	5	.15	0.11	0.23	0.71	NHEL
UdD	Udorthents, 0 to 10 percent slopes	Udorthents	100	0-10	-					
UsE	Udorthents, sanitary landfill, 2 to 25 percent slopes	Udorthents	95	2-25	-					
W	Water	Water	100		-					
WaA	Wahee-Ogeechee	Wahee	40	0-3	5	.15	0.11	0.3	0.71	NHEL
	complex, 0 to 3 percent	Ogeechee	20	0-3	5	.1	0.11	0.3	1.07	NHEL
	slopes, occasionally flooded	Rains	20	0-3	5	.15	0.11	0.3	0.71	NHEL

HEL Highly Erodible Land

PHEL Potentially Highly Erodible Land

NHEL Not Highly Erodible Land

- All HEL Classes were assigned using USLE factor values that were in effect as of January 1, 1990.
- This HEL list is based on the final correlation of survey data and supersedes any previous HEL list.