

NATURAL RESOURCES CONSERVATION SERVICE
CONSERVATION PRACTICE STANDARD
CONSERVATION CROP ROTATION

(Ac.)

CODE 328

DEFINITION

A planned sequence of crops grown on the same ground over a period of time (i.e. the rotation cycle).

PURPOSES

This practice is applied to support one or more of the following purposes:

- Reduce sheet, rill and wind erosion.
- Maintain or increase soil health and organic matter content.
- Reduce water quality degradation due to excess nutrients.
- Improve soil moisture efficiency.
- Reduce plant pest pressures.
- Provide feed and forage for domestic livestock.
- Provide food and cover habitat for wildlife, including pollinator forage, and nesting.

CONDITIONS WHERE PRACTICE APPLIES

This practice applies to all cropland where at least one annually-planted crop is included in the crop rotation.

CRITERIA

General Criteria Applicable to All Purposes

Crops shall be grown in a planned sequence as outlined in Plans and Specifications. The crop rotation shall include a minimum of two different crops. For purposes of these criteria a cover crop is considered a different crop.

However, rotations among crops planted during the same season and other practices

noted in the Considerations section are likely to be more effective than just rotating crops between seasons when the purpose is promoting soil health or managing some plant pests.

Where applicable, plan suitable crop substitutions when the planned crop cannot be planted due to weather, soil conditions, or other local situations.

Additional Criteria to Reduce Sheet, Rill and Wind Erosion

Select crops, a tillage system, and cropping sequences that will produce sufficient and timely quantities of biomass or crop residue which, in conjunction with other practices in the management system, will reduce sheet, rill and wind erosion to the planned soil loss objective.

Determine the amount of biomass or crop residue needed by using RUSLE2 or the current approved erosion prediction technology. Compare the current soil loss and the planned erosion reduction to the actual soil loss obtained as result of implementing the practice.

Additional Criteria to Maintain or Increase Soil Health and Organic Matter Content

Grow crops that will produce a positive trend in the Organic Matter (OM) subfactor value over the life of the rotation, as determined by the Soil Conditioning Index (SCI) obtained from RUSLE2 or current soil erosion software. Make appropriate adjustments for additions to or subtractions from biomass during the rotation. Compare the current and planned soil health benefit to the actual result of implementing the practice.

Conservation practice standards are reviewed periodically and updated if needed. To obtain the current version of this standard, contact your Natural Resources Conservation Service [State Office](#) or visit the [Field Office Technical Guide](#).

NRCS, Georgia
October 2015

Additional Criteria to Reduce Water Quality Degradation Due to Excess Nutrients

Cover crops are frequently used to recover excess nutrients remaining in the soil profile from a previous crop. The benefit of using a cover crop to manage excess nutrients is the time of planting is usually immediately following the crop. See the Cover Crop (340) Standard for more information including species and planting information.

Cover crops and most subsequent crops are characterized by:

- Quick germination and root system formation,
- A rooting depth sufficient to reach the nutrients not removed by the previous crop, and
- Nutrient requirements that readily utilize the excess nutrients
- Credit nutrients provided by legumes and manure/compost as noted in the Cover Crop (340) and Nutrient Management (590) Standards.

Additional Criteria to Improve Soil Moisture Efficiency

Select crops, varieties of crops, and the sequence of crops based on local climate patterns, soil conditions, irrigation water availability, and an approved water balance procedure in the Georgia NRCS Irrigation Water Management Standard (Code 449).

Additional Criteria to Reduce Plant Pest Pressures

Design the crop sequence to suppress the pest(s) lifecycle of concern, which may include weeds, insects, and pathogens. Use a crop rotation approved by UGA or industry standard to determine a suitable crop sequence.

Rotations also allow for the use of multiple herbicide options to increase weed control and reduce the possibility of herbicide resistance occurring in weeds.

Additional Criteria to Provide Feed and Forage for Domestic Livestock

Select crops that balance the feed supply with livestock numbers. Determine the required amount of selected crops using the graze

spreadsheet <http://efotg.sc.egov.usda.gov/toc.aspx?CatID=2918> .

Additional Criteria to Provide Food and Habitat for Wildlife, Including Pollinator Forage, and Nesting

Select the crops and crop management activities that provide either food or cover for the targeted wildlife species using the Georgia NRCS wildlife suitability index (Wildlife HIS) for cropland (<http://efotg.sc.egov.usda.gov/toc.aspx?CatID=2918>).

CONSIDERATIONS

When used in combination with NRCS Conservation Practice (CPS) Stripcropping (Code 585), the crop sequence should be consistent with the stripcropping design.

Soil compaction can be reduced by adjusting crop rotations to include deep rooting crops with deep roots that extend to and penetrate compacted soil layers.

Where improving water use efficiency on deep soils is a concern, rotating or combining deep-rooted crops with shallow rooted crops can help utilize all available water in the soil profile.

Select crops that have the potential to provide larger amounts of biologically fixed nitrogen.

Considerations to Reduce Water Quality Degradation Due to Excess Nutrients:

- Include perennial or annual legume crops in the rotation to provide nitrogen for the non-legume crops, especially in fields where manure applications are restricted by high or excessive soil phosphorus or potassium levels. Legumes will also use residual nitrogen in the soil, if available, rather than produce it.
- Cereal grains, pasture grasses and sorghum-sudangrass are examples of crops that have high carbon/nitrogen ratios. The use of these plants as cover crops will minimize leaching and make nitrogen temporarily unavailable to subsequent crops.

Considerations to Increase Cropping System Diversity

Minimize the fallow years in the rotation and where the climate and soils are favorable establish cover crops during the fallow periods.

For crop diversity, the planned crop sequence should contain different crop types; for example a mix of the following: warm season grass; warm season broadleaf; cool season grass; cool season broadleaf.

- A two-crop sequence that contains a warm season and a cool season crop;
- A three-crop sequence that contains warm and cool season crops. The same crop species should not be grown in successive years in the same field.
- A four-crop sequence that contains two different crop types, neither should occupy more than half of the sequence.
- Longer crop sequences (four or more years) are more effective with no more than two consecutive years with the same crop.

Additional Considerations to Reduce Sheet and Rill or Wind Erosion.

When used in combination with NRCS CPSs for Residue and Tillage Management (Codes 329 and 345), selection of high-residue producing crops and varieties, use of cover crops and adjustment of plant density and row spacing can enhance production of the kind, amount, and distribution of residue needed. The use of a cover crop is required only with no- and strip-tillage.

When used in combination with NRCS CPSs Stripcropping (Code 585) or Contour Buffer Strips (Code 332) on steeper slopes, the effectiveness of each practice is significantly enhanced by inclusion of the other practice(s) in the conservation system.

Crop damage by wind erosion can be reduced with this practice by selecting crops that are tolerant to abrasion from windblown soil or tolerant to high wind velocity.

If crops sensitive to wind erosion damage are grown, the potential for plant damage can be reduced by crop residue management, field

windbreaks, intercropping, or other methods of wind erosion control.

Additional Considerations to Improve Soil Health

Consider including perennial sod crops with deep or extensive fibrous root systems to build organic matter throughout the soil profile. These crops produce large quantities of residue and provide cover to the soil throughout the year. As a result, sod-based rotations will result in high SCI values based upon a positive OM subfactor and a low field operations subfactor (FO). Furthermore, the Soil Tillage Intensity Rating (STIR) value, another independent factor generated with RULSE2 software, is a measure of the soil disturbance. STIR values will be low with no-till and other crop production systems that include perennial crops.

The effects of this practice can be enhanced by utilizing animal wastes, green manure crops (cover crops), or applying non-synthetic mulches to supplement the biomass produced by crops in the rotation.

Other considerations for soil health/organic matter management include:

- For at least one-third of the crop sequence (time basis) include high-biomass annual or perennial crops.
- Utilize cover crops and high residue production crops comprising at least one-half of the rotation sequence.
- For rotations dominated by low-residue crops, such as vegetables, include sufficient cover crops and high residue crops for one-half the rotation.

Additional Considerations to Reduce Plant Pest Pressures

Consider lengthening the rotation to include several years of perennial cover to break pest life cycles.

Use a mix of crops from at least three different plant families, and allow three years or longer between successive plantings of production crops within the same family.

Enhance biological pest control by designing the crop rotation to:

- Include flowering annuals or perennials that provide food and habitat for beneficial insects, such as buckwheat or clovers.
- Include plant species that release into the soil natural substances that suppress plant pathogens, nematodes or pests (biofumigation).
- Include crops in the rotation that provide habitat for natural enemies of pests.
- Retain bolting or flowering crops after harvest to provide food for beneficial insects.

Additional Considerations to Provide Food and Cover Habitat for Wildlife, Including Pollinator Forage, and Nesting

Crop residues may be a valuable food source for wintering wildlife where winter browse is sparse. Leaving several rows unharvested around the edges of the field, or planting borders of various forbs will provide protection and/or food for overwintering wildlife and for beneficial insects and pollinators.

Crop plantings may be developed to benefit particular communities, species, or life stages of wildlife. Food plots or crops for wildlife can provide part of a habitat restoration, an initial food and cover for wildlife until food and cover producing vegetation becomes established.

Retaining bolting or flowering crops after harvest may provide beneficial insects with an important food source.

Careful consideration should be given to pesticides applied to crops raised for wildlife, particularly if nesting habitat or pollinator forage species are present.

When insect-pollinated crops are part of the rotation, planting the insect-pollinated crop no more than 800 feet from their previous location may help maintain local populations of native bees that have become established because of the presence of that crop.

To maintain stable pollinator and beneficial insect populations, ensure that the same overall density of floral resources is maintained from year-to-year. For example two years of flower-rich plantings, followed by a year of only grasses, will cause a rapid decline in pollinator

populations. Such a scenario is undesirable. A list of pollinator plants is available in the Conservation Cover (327) Jobsheet through this link <http://efotg.sc.egov.usda.gov/toc.aspx?CatID=2918>.

PLANS AND SPECIFICATIONS

Develop plans and specifications for each field or treatment unit according to the Criteria and Operation and Maintenance requirements of this standard. Specifications shall describe the requirements to apply this practice to achieve the intended purpose. Record the following specification components in an approved Conservation Crop Rotation, 328, Implementation Requirements document, including the jobsheet. The following items will be documented as a minimum.

- Field number and acres,
- Purpose(s) of the crop rotation,
- The sequence of crops to be grown,
- The crop types to be grown,
- Tillage type and times,
- Length of time each crop/crop type will be grown in the rotation, and
- Total length of rotation
- Suitable crop substitutions to address weather, soil conditions, market, or other situations that may prevent the planned crop from being planted.

OPERATION AND MAINTENANCE

Rotations shall provide for acceptable substitute crops in case of crop failure or shift in planting intentions for weather related or economic reasons. Acceptable substitutes are crops having similar properties that will accomplish the purpose of the original crop.

Evaluate the rotation and the crop sequence to determine if the planned system is meeting the planned purposes.

REFERENCES FROM NRCS, NHCP SEPTEMBER 2014

Green, B., D. Kaminski, B. Rapp, M. Celetti, D. Derksen, L. Juras, and D. Kelner. 2005.

NRCS, Georgia

October 2015

Principles and practices of crop rotation.
Saskatchewan Agriculture and Food.

Karlen, D.L., E.G. Hurley, S.S. Andrews, C.A. Cambardella, D.W. Meek, M.D. Duffy, and A.P. Mallorino. 2006. Crop rotation effects on soil quality at three northern corn/soybean belt locations. *Agron. J.* 98:484-495.

Liebig, M.A., D.L. Tanaka, J.M. Krupinsky, S.D. Merrill, and J.D. Hanson. 2007. Dynamic cropping systems: Contributions to improve agroecosystem sustainability. *Agron. J.* 99:899-903.

Sherrod, L.A., G.A. Peterson, D.G. Westfall, and L.R. Ahuja. 2003. Cropping intensity enhances soil organic carbon and nitrogen in a no-till agroecosystem. *Agron. J.* 67:1533-1543.

USDA-AMS National Organic Program Final Rule 7 CFR Part 205. <http://www.ams.usda.gov/AMSV1.0/nop>.

USDA, NRCS. 2014. Revised Universal Soil Loss Equation Version 2 (RUSLE2) website: Washington, DC. <http://www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/tools/rusle2/>

USDA, NRCS. 2014. Wind Erosion Prediction System (WEPS) website: Washington, DC. <http://www.nrcs.usda.gov/wps/portal/nrcs/main/national/technical/tools/weeps/>

USDA, NRCS. 2011. National Agronomy Manual, 4th Ed. Washington, DC. <http://directives.sc.egov.usda.gov/viewerFS.aspx?hid=29606>

USDA, NRCS. 2014. Preventing or mitigating potential negative impacts of pesticides on pollinators using IPM and other conservation practices. Nat. Agron. Tech Note 9. Washington, DC. <http://directives.sc.egov.usda.gov/OpenNonWebContent.aspx?content=34828.wba>