

# Fertilizer Guide

## for Commercial Vegetable Growers

UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN • COLLEGE OF AGRICULTURE • COOPERATIVE EXTENSION SERVICE  
CIRCULAR 1185

COMMERCIAL VEGETABLE PRODUCERS must depend on chemical fertilizers and organic materials along with crop rotation to supply the nutrients essential to their vegetable crops. Although many Illinois soils are highly fertile naturally, they cannot support continuous intensive cropping of vegetables without additional fertilizer. Nutrients can also be replenished by alternating vegetables with green manure or cover crops and by adding manure or other organic materials. Regardless of how the nutrients are added, efficient commercial production of vegetables requires soil testing on a regular basis. Using fertilizer materials efficiently provides greater profits and minimizes the release of potential pollutants into the environment.

### Testing Your Soil

To determine how much of each nutrient must be added for optimum crop production, collect soil samples every 2 or 3 years and have them analyzed for pH and for phosphorus and potassium concentrations. The test results and your knowledge of the field's cropping and fertilization history will provide the information you need to develop a fertilization plan for the crop to be grown.

Samples should be collected in the late fall when the soil is relatively dry but not yet frozen. Separate samples should be tested for every field that differs in color, slope, drainage, or previous fertilization and cropping. Each sample should represent no more than 4 acres and should consist of several subsamples collected at random locations throughout the field. Check with your soil testing laboratory for more specific instructions.

### Determining Fertilizer Application Rates

The soil test results will be reported in terms of the amount of elemental phosphorus (P) and elemental potassium (K) per acre. Referring to Table 1, determine which of the four soil fertility groups (A through D) your soil's phosphorus test level fits into. Then do the same for the potassium test level. These groups can then be used in con-

Table 1. Soil Fertility Groups for Phosphorus and Potassium

| Fertility group | Nutrient, pounds per acre      |           |
|-----------------|--------------------------------|-----------|
|                 | Bray P <sub>1</sub> phosphorus | Potassium |
| A               | 0-25                           | 0-100     |
| B               | 26-50                          | 101-250   |
| C               | 51-75                          | 251-350   |
| D               | Above 75                       | Above 350 |

junction with Table 2 to determine how much of the two nutrients you should add for the crop you plan to grow.

Locate the crop in Table 2, and then find the column under the phosphorus heading that corresponds to your soil's fertility group for phosphorus. The numbers in the column below the fertility group will tell you how many pounds of P<sub>2</sub>O<sub>5</sub> you need to apply per acre to increase the phosphorus content to a satisfactory level. Follow the same procedure using the numbers in the potassium column to determine how much K<sub>2</sub>O you should apply. If soil test results are not available, use the amounts of phosphorus and potassium recommended for fertility group B.

Since soil tests for nitrogen are of little value, the nitrogen recommendations in Table 2 are based on the needs of the various crops, but in developing a fertilization program you should also take into account the field's cropping history and the type of soil.

If the crop grown in the field during the previous year was a legume (soybeans or alfalfa), the amount of nitrogen applied can be 25 to 30 pounds per acre less than that recommended in Table 2. The nitrogen status of most vegetable crops can be determined by the color of the foliage. A pale green or slightly yellow color may indicate a need to apply additional nitrogen.

Unless otherwise indicated, the fertilizer recommendations given in this circular are for the mineral soils that predominate in Illinois. Vegetable crops grown on sandy soils usually require greater amounts of nitrogen and potassium. Splitting the nitrogen fertilizer between two separate applications will result in greater efficiency and production on sandy soils that are irrigated or that receive heavy rainfall.

Plantings made early in the season in cool, wet soils may respond well to the application of band-placed phosphorus or a starter solution in addition to the nutrients recommended in Table 2.

### Adjusting for Additional Organic Matter

Do not adjust for cover crops because they supply little additional nutrition. Cover crops primarily affect the structure rather than the fertility of the soil. When wheat or rye is plowed down, however, apply a chemical fertilizer to encourage rapid decomposition because decaying wheat and rye may remove 20 to 30 pounds of nitrogen per acre. About 30 percent of this nitrogen will be released to the crop later in the season. If the previous crop was a green manure or a cropped legume, like soybeans or alfalfa, the recommendation for adding nitrogen may be reduced by 25 to 30 pounds per acre. The adjustment for manure is provided in Tables 3 and 4.

**Table 2. Fertilizer Recommendations for Vegetable Crops**

| Crop  | Recommended application rate based on soil tests |   |     |     |     |                              |     |     |    | Suggested application method   |
|---|--|---|-----|-----|-----|------------------------------|-----|-----|----|--|
|   | Nitrogen (N)                                     | Phosphorus (P <sub>2</sub> O <sub>5</sub> ) |     |     |     | Potassium (K <sub>2</sub> O) |     |     |    |  |
|   |  | Fertility group                             |     |     |     | Fertility group              |     |     |    |  |
|   |  | A   | B   | C   | D   | A                            | B   | C   | D  |  |
| <b>ASPARAGUS</b>  |  |   |     |     |     |                              |     |     |    |  |
|   |  | <i>pounds per acre</i>                      |     |     |     |                              |     |     |    |  |
| New plantings   | 50   | 150   | 50  | 25  | 0   | 150                          | 50  | 25  | 0  | Broadcast and plow down<br>Apply in trench before setting<br>Side-dress at first cultivation |
|   | 0  | 50  | 50  | 25  | 25  | 0                            | 0   | 0   | 0  |  |
|   | 30   | 0   | 0   | 0   | 0   | 50                           | 50  | 25  | 25 |  |
| Total   | 80   | 200   | 100 | 50  | 25  | 200                          | 100 | 50  | 25 |  |
| <b>Cutting beds</b>   |  |   |     |     |     |                              |     |     |    |  |
| Nonhybrids  | 50   | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50 | Broadcast and disk   |
| Hybrids   | 75   | 200   | 150 | 100 | 50  | 300                          | 225 | 150 | 75 | Broadcast and disk   |
| In sandy soils an additional 50 pounds of nitrogen per acre can be applied as a sidedress after cutting. In new beds build up organic matter with cover crops and manure 1 or 2 years before planting crowns.   |  |   |     |     |     |                              |     |     |    |  |
| <b>BEANS</b>  |  |   |     |     |     |                              |     |     |    |  |
| Lima  | 20   | 150   | 100 | 50  | 0   | 150                          | 100 | 50  | 0  | Broadcast and plow down<br>Band 2 in. × 2 in. at seeding                                     |
|   | 40   | 40  | 40  | 20  | 20  | 40                           | 40  | 20  | 20 |  |
| Total   | 60   | 190   | 140 | 70  | 20  | 190                          | 140 | 70  | 20 |  |
| Snap  | 0  | 150   | 100 | 50  | 0   | 100                          | 50  | 25  | 0  | Broadcast and plow down<br>Band 2 in. × 2 in. at seeding                                     |
|   | 40   | 40  | 40  | 20  | 20  | 40                           | 40  | 20  | 20 |  |
| Total   | 40   | 190   | 140 | 70  | 20  | 140                          | 90  | 45  | 20 |  |
| Snap, second crop   | 30   | 20  | 20  | 20  | 20  | 40                           | 40  | 20  | 20 | Band 2 in. × 2 in. at seeding  |
| In sandy soils an additional 25 pounds of nitrogen per acre can be applied as a sidedress when two or three true leaves have appeared. If the soil pH is greater than 6.8, apply 2 pounds of zinc and 1 pound of manganese per acre at planting.  |  |   |     |     |     |                              |     |     |    |  |
| <b>BEETS</b>  |  |   |     |     |     |                              |     |     |    |  |
|   | 75   | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50 | Broadcast and disk   |
|   | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 4 to 6 weeks after planting   |
| Total   | 125  | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50 |  |
| Apply 3 pounds of boron per acre on clay loams and 1 pound per acre on sandy soils.   |  |   |     |     |     |                              |     |     |    |  |
| <b>BROCCOLI AND CAULIFLOWER</b>   |  |   |     |     |     |                              |     |     |    |  |
|   | 100  | 200   | 150 | 75  | 50  | 200                          | 150 | 75  | 50 | Broadcast and disk   |
|   | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 2 to 3 weeks after transplanting  |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 5 to 6 weeks after transplanting if required                                      |
| Total   | 175  | 200   | 150 | 75  | 50  | 200                          | 150 | 75  | 50 |  |
| If the pH of the soil is greater than 6.5, apply 2 pounds of boron per acre on clay loams and 1 pound per acre on sandy soils. Also, a foliar application of 8 ounces per acre in 100 gallons of water of sodium or ammonium molybdate may avoid molybdenum deficiency. Early plantings in cold soil may respond to a high-phosphorus starter solution. |  |   |     |     |     |                              |     |     |    |  |
| <b>CABBAGE, COLLARDS, AND KALE</b>  |  |   |     |     |     |                              |     |     |    |  |
|   | 75   | 150   | 100 | 50  | 25  | 150                          | 100 | 50  | 25 | Broadcast and disk   |
|   | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 4 weeks after planting  |
| Total   | 125  | 150   | 100 | 50  | 25  | 150                          | 100 | 50  | 25 |  |
| Early plantings in cold soil may respond well to a high-phosphorus starter solution.  |  |   |     |     |     |                              |     |     |    |  |
| <b>CARROTS</b>  |  |   |     |     |     |                              |     |     |    |  |
|   | 50   | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50 | Broadcast and disk   |
|   | 30   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 4 weeks after seeding   |
| Total   | 80   | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50 |  |
| In sandy soils an additional 30 pounds of nitrogen per acre may be applied as a sidedress between 7 and 8 weeks after seeding.  |  |   |     |     |     |                              |     |     |    |  |
| <b>CELERY</b>   |  |   |     |     |     |                              |     |     |    |  |
|   | 100  | 250   | 200 | 150 | 100 | 300                          | 200 | 100 | 50 | Broadcast and disk   |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 4 weeks after planting  |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0  | Side-dress 8 weeks after planting  |
| Total   | 150  | 250   | 200 | 150 | 100 | 300                          | 200 | 100 | 50 |  |
| Because celery has a high moisture requirement, irrigation is essential for commercial production. Use of a starter solution is recommended when transplanting celery.  |  |   |     |     |     |                              |     |     |    |  |

**Table 2. Fertilizer Recommendations for Vegetable Crops (continued)**

| Crop  | Recommended application rate based on soil tests |   |     |     |     |                              |     |     |     | Suggested application method  |
|---|--|---|-----|-----|-----|------------------------------|-----|-----|-----|---|
|   | Nitrogen (N)                                     | Phosphorus (P <sub>2</sub> O <sub>5</sub> ) |     |     |     | Potassium (K <sub>2</sub> O) |     |     |     |   |
|   |  | Fertility group                             |     |     |     | Fertility group              |     |     |     |   |
|   |  | A   | B   | C   | D   | A                            | B   | C   | D   |   |
| <i>pounds per acre</i>  |  |   |     |     |     |                              |     |     |     |   |
| <b>RHUBARB</b>  |  |   |     |     |     |                              |     |     |     |   |
| New plantings   | 50   | 250   | 200 | 150 | 100 | 250                          | 200 | 150 | 50  | Broadcast and plow down<br>Side-dress around each hill 2 weeks<br>after growth starts                                       |
|   | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 100  | 250   | 200 | 150 | 100 | 250                          | 200 | 150 | 50  |   |
| Cutting beds  | 50   | 200   | 150 | 100 | 50  | 250                          | 150 | 100 | 50  | Side-dress each hill in early spring<br>Side-dress at last harvest  |
|   | 50   | 50  | 50  | 50  | 50  | 50                           | 50  | 50  | 50  |   |
| Total   | 100  | 250   | 200 | 150 | 100 | 300                          | 200 | 150 | 100 |   |
| <b>SPINACH</b>  | 100  | 200   | 150 | 100 | 50  | 200                          | 150 | 100 | 50  | Broadcast and disk<br>Side-dress 4 to 5 weeks after planting  |
|   | 20   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 120  | 200   | 150 | 100 | 50  | 200                          | 150 | 100 | 50  |   |
| <b>SQUASH</b>   | 75   | 150   | 100 | 50  | 25  | 200                          | 100 | 50  | 25  | Broadcast and disk<br>Side-dress when vines start to run  |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 100  | 150   | 100 | 50  | 25  | 200                          | 100 | 50  | 25  |   |
| <b>SWEET CORN</b>   | 100  | 150   | 100 | 75  | 50  | 200                          | 100 | 75  | 50  | Broadcast and disk<br>Side-dress when corn is 12 in. tall   |
|   | 30   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 130  | 150   | 100 | 75  | 50  | 200                          | 100 | 75  | 50  |   |
| On soils low in phosphorus or for early plantings band (2 inches × 2 inches) 30 pounds of phosphorus per acre at seeding. |  |   |     |     |     |                              |     |     |     |   |
| <b>SWEET POTATO</b>   | 60   | 200   | 150 | 100 | 50  | 200                          | 150 | 50  | 0   | Broadcast and plow down   |
| <b>TOMATOES — Fresh Market</b>  |  |   |     |     |     |                              |     |     |     |   |
| On sandy soils  | 100  | 250   | 200 | 100 | 50  | 200                          | 150 | 75  | 50  | Broadcast and plow down<br>Side-dress at first cultivation<br>Side-dress after first fruit set                              |
|   | 25   | 0   | 0   | 0   | 0   | 50                           | 50  | 25  | 0   |   |
|   | 25   | 0   | 0   | 0   | 0   | 50                           | 0   | 0   | 0   |   |
| Total   | 150  | 250   | 200 | 100 | 50  | 300                          | 200 | 100 | 50  |   |
| The second sidedress may not be required on early or semideterminate tomatoes.  |  |   |     |     |     |                              |     |     |     |   |
| On loams  | 100  | 250   | 200 | 100 | 50  | 250                          | 150 | 100 | 50  | Broadcast and plow down<br>Side-dress at first cultivation  |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 125  | 250   | 200 | 100 | 50  | 250                          | 150 | 100 | 50  |   |
| — Processing: Transplants/Direct seeded   |  |   |     |     |     |                              |     |     |     |   |
| On sandy soils  | 60/50  | 250   | 200 | 100 | 50  | 200                          | 150 | 50  | 0   | Broadcast and disk<br>Side-dress transplants 4 weeks after<br>planting and direct seeded tomatoes<br>6 weeks after planting |
|   | 30/25  | 0   | 0   | 0   | 0   | 50                           | 50  | 50  | 50  |   |
|   | 0/0  | 0   | 0   | 0   | 0   | 50                           | 0   | 0   | 0   |   |
| Total   | 90/75  | 250   | 200 | 100 | 50  | 300                          | 200 | 100 | 50  |   |
| On loams  | 50/30  | 250   | 200 | 100 | 50  | 200                          | 100 | 50  | 50  | Broadcast and disk<br>Side-dress transplants 4 weeks after<br>planting and direct seeded tomatoes<br>6 weeks after planting |
|   | 25/30  | 0   | 0   | 0   | 0   | 50                           | 50  | 50  | 0   |   |
| Total   | 75/60  | 250   | 200 | 100 | 50  | 250                          | 150 | 100 | 50  |   |
| The use of a starter solution is recommended when transplanting tomatoes.   |  |   |     |     |     |                              |     |     |     |   |
| <b>TURNIPS</b>  | 50   | 100   | 75  | 50  | 0   | 100                          | 50  | 0   | 0   | Broadcast and disk  |
| Apply 2 pounds of boron per acre on clay loams and 1 pound per acre on sandy soils.                                       |  |   |     |     |     |                              |     |     |     |   |
| <b>WATERMELONS</b>  | 50   | 150   | 100 | 50  | 0   | 150                          | 100 | 50  | 0   | Broadcast and plow down<br>Band 2 in. × 2 in. at seeding<br>Side-dress when vines start to run                              |
|   | 25   | 50  | 50  | 50  | 50  | 50                           | 50  | 50  | 50  |   |
|   | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0   |   |
| Total   | 100  | 200   | 150 | 100 | 50  | 200                          | 150 | 100 | 50  |   |
| On sandy soils increase the nitrogen sidedressing to 50 pounds of nitrogen per acre when vines start to run.              |  |   |     |     |     |                              |     |     |     |   |

**Table 2. Fertilizer Recommendations for Vegetable Crops (continued)**

| Crop   | Recommended application rate based on soil tests |   |     |     |     |                              |     |     | Suggested application method |  |
|--|--|---|-----|-----|-----|------------------------------|-----|-----|------------------------------|--|
|  | Nitrogen (N)                                     | Phosphorus (P <sub>2</sub> O <sub>5</sub> ) |     |     |     | Potassium (K <sub>2</sub> O) |     |     |                              |  |
|  |  | Fertility group                             |     |     |     | Fertility group              |     |     |                              |  |
|  |  | A   | B   | C   | D   | A                            | B   | C   | D                            |  |
| <i>pounds per acre</i>   |  |   |     |     |     |                              |     |     |                              |  |
| <b>CUCUMBERS</b>   | 50   | 100   | 50  | 0   | 0   | 150                          | 100 | 50  | 0                            | Broadcast and plow down                |
|  | 25   | 50  | 50  | 50  | 25  | 50                           | 50  | 50  | 50                           | Band 2 in. × 2 in. at seeding          |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress when vines start to run     |
|  | Total  | 100   | 150 | 100 | 50  | 25                           | 200 | 150 | 100                          | 50                                     |
| <b>EGGPLANT</b>  | 75   | 200   | 150 | 100 | 50  | 250                          | 150 | 100 | 50                           | Broadcast and disk                     |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress 4 weeks after planting      |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress 8 weeks after planting      |
|  | Total  | 125   | 200 | 150 | 100 | 50                           | 250 | 150 | 100                          | 50                                     |
| Use of a starter solution is recommended when transplanting eggplant.  |  |   |     |     |     |                              |     |     |                              |  |
| <b>HORSERADISH</b>   | 150  | 250   | 200 | 100 | 50  | 250                          | 200 | 100 | 50                           | Broadcast and disk                     |
| <b>LETTUCE (LEAF), EN-DIVE, AND ESCAROLE</b>   | 80   | 200   | 100 | 50  | 25  | 200                          | 100 | 50  | 25                           | Broadcast and disk                     |
|  | 40   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress 3 to 5 weeks after planting |
|  | Total  | 120   | 200 | 100 | 50  | 25                           | 200 | 100 | 50                           | 25                                     |
| Apply only 75 pounds of nitrogen per acre to head lettuce because excessive nitrogen may result in loose heads.  |  |   |     |     |     |                              |     |     |                              |  |
| <b>MUSKMELON</b>   | 50   | 150   | 100 | 50  | 0   | 150                          | 100 | 50  | 0                            | Broadcast and plow down                |
|  | 25   | 50  | 50  | 50  | 50  | 50                           | 50  | 50  | 50                           | Band 2 in. × 2 in. at seeding          |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress when vines start to run     |
|  | Total  | 100   | 200 | 150 | 100 | 50                           | 200 | 150 | 100                          | 50                                     |
| On sandy soils increase the nitrogen sidedressing to 50 pounds of nitrogen per acre when vines start to run.   |  |   |     |     |     |                              |     |     |                              |  |
| <b>ONIONS</b>  | 75   | 200   | 100 | 50  | 25  | 200                          | 100 | 50  | 25                           | Broadcast and disk                     |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress 4 to 5 weeks after planting |
|  | Total  | 100   | 200 | 100 | 50  | 25                           | 200 | 100 | 50                           | 25                                     |
| For green onions an additional 25 pounds of nitrogen per acre can be applied as a sidedress from 4 to 5 weeks before harvest.  |  |   |     |     |     |                              |     |     |                              |  |
| <b>PARSLEY</b>   | 75   | 200   | 150 | 100 | 50  | 200                          | 150 | 100 | 50                           | Broadcast and disk                     |
|  | 25   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress at each cutting             |
|  | Total  | 100   | 200 | 150 | 100 | 50                           | 200 | 150 | 100                          | 50                                     |
| <b>PEAS</b>  | 0  | 100   | 50  | 25  | 0   | 100                          | 75  | 50  | 25                           | Broadcast and disk                     |
|  | 50   | 50  | 50  | 50  | 50  | 50                           | 25  | 25  | 25                           | Band 2 in. × 2 in. at seeding          |
|  | Total  | 50  | 150 | 100 | 75  | 50                           | 150 | 100 | 75                           | 50                                     |
| After periods of heavy rains in the spring an additional 30 pounds of nitrogen per acre can be applied when peas are 4 to 6 inches tall. Apply only when the plants are dry to avoid burning the foliage.  |  |   |     |     |     |                              |     |     |                              |  |
| <b>PEPPERS</b>   | 100  | 200   | 150 | 100 | 50  | 250                          | 150 | 100 | 50                           | Broadcast and disk                     |
|  | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress after first fruit set       |
|  | Total  | 150   | 200 | 150 | 100 | 50                           | 250 | 150 | 100                          | 50                                     |
| In sandy soils an additional 25 pounds of nitrogen may be applied as a sidedress after the first harvest. The use of a starter solution is recommended when transplanting peppers.   |  |   |     |     |     |                              |     |     |                              |  |
| <b>POTATOES</b>  | 0  | 150   | 100 | 0   | 0   | 200                          | 150 | 100 | 50                           | Broadcast and disk                     |
|  | 100  | 100   | 100 | 100 | 100 | 100                          | 100 | 100 | 50                           | Band-place at planting                 |
|  | 50   | 0   | 0   | 0   | 0   | 0                            | 0   | 0   | 0                            | Side-dress at emergence                |
|  | Total  | 150   | 250 | 200 | 100 | 100                          | 300 | 250 | 200                          | 100                                    |
| In sandy soils an additional 30 pounds of nitrogen can be applied as a sidedress before the plants are about 8 or 10 inches tall. On soils with magnesium test values less than 250 pounds per acre, apply 50 pounds magnesium (MgO) per acre at planting. |  |   |     |     |     |                              |     |     |                              |  |
| <b>PUMPKINS</b>  | 90   | 150   | 100 | 50  | 25  | 200                          | 150 | 100 | 50                           | Broadcast and disk                     |
|  | 30   | 0   | 0   | 0   | 0   | 50                           | 0   | 0   | 0                            | Side-dress when vines start to run     |
|  | Total  | 120   | 150 | 100 | 50  | 25                           | 250 | 150 | 100                          | 50                                     |
| Excessive use of nitrogen may result in thin walls and a flat side on jack-o'-lanterns.  |  |   |     |     |     |                              |     |     |                              |  |

**Table 3. Reduction in Fertilizer Application for Applied Liquid Manures<sup>a</sup>**

| Manure source                | Nitrogen            |                  |                            | Phosphorus | Potassium |
|------------------------------|---------------------|------------------|----------------------------|------------|-----------|
|                              | Time of application |                  |                            |            |           |
|                              | Fall or winter      | Spring uncovered | Spring covered or injected |            |           |
|                              |                     |                  | <i>pounds per acre</i>     |            |           |
| Cattle, mixed livestock..... | 5                   | 10               | 12                         | 4          | 16        |
| Poultry .....                | 23                  | 46               | 58                         | 22         | 26        |
| Swine .....                  | 8                   | 15               | 19                         | 7          | 14        |

<sup>a</sup> Assuming an application of 900 gallons per acre.

**Table 4. Reduction in Fertilizer Application for Applied Solid Manure<sup>a</sup>**

| Manure source                | Nitrogen            |                  |                            | Phosphorus | Potassium |
|------------------------------|---------------------|------------------|----------------------------|------------|-----------|
|                              | Time of application |                  |                            |            |           |
|                              | Fall or winter      | Spring uncovered | Spring covered or injected |            |           |
|                              |                     |                  | <i>pounds per acre</i>     |            |           |
| Cattle, mixed livestock..... | 12                  | 24               | 30                         | 10         | 44        |
| Poultry .....                | 70                  | 140              | 175                        | 75         | 96        |
| Swine .....                  | 15                  | 30               | 38                         | 20         | 26        |

<sup>a</sup> Assuming an application of 4.5 tons per acre.

### Other Nutrient Requirements

**Calcium and magnesium.** Calcium and magnesium are dominant cations in most Illinois soils. Most vegetable crops need relatively small amounts of calcium. A calcium deficiency is extremely rare in soils with a pH value of 5.0 and above and unlikely even in soils that require lime to correct acidity. Soil test levels for calcium of 1000 pounds per acre on sandy soils and of 1500 pounds per acre on silt loams are adequate for most vegetable crops.

It is more probable that you will encounter a magnesium deficiency while growing vegetables, especially when the soil is acid and sandy or when large amounts of calcitic limestone or marl have been applied to soils already low in magnesium. Excessive potash fertilizer may also cause a magnesium deficiency on some sandy soils. Soil test values for magnesium of 300 pounds per acre on sandy soil and 500 pounds per acre on silt loams are adequate. Apply dolomitic limestone on acid sandy soils that need magnesium. Other suitable magnesium fertilizers include magnesium sulfate (Epsom salts) and potassium-magnesium-sulfate. Foliar application of 20 pounds of magnesium sulfate in 60 gallons of water is a quick remedy for magnesium deficiency in several vegetables. Vegetables that respond to magnesium include cauliflower, muskmelons, peppers, potatoes, and pumpkins.

**Sulfur.** Vegetable crop soils in Illinois are not usually known to be deficient in sulfur, and field tests have not shown a significant yield response to applications of this nutrient. Environmental sources of sulfur usually satisfy the needs of plants. These sources include mineralized organic matter, atmospheric sulfur (released from emissions of fossil fuels and returned to the soil in precipitation), and fertilizer by-products. With greater control of air pollution and in-

creased use of high analysis fertilizers, the amount of sulfur obtained from fertilizer by-products and emissions of fossil fuels is declining. As a result sulfur deficiencies may develop in the next five to fifteen years and are most likely to occur in sandy soils or in soils low in organic matter located upwind from urban industrial areas.

Soil tests for sulfur are not very reliable because sulfate ions are highly mobile in the soil. Low soil tests for sulfur should be confirmed by tissue analysis. Most vegetable crops require as much sulfur as phosphorus. Either elemental or sulfate forms of sulfur fertilizers can be used on vegetable crops. Sulfate-sulfur reacts more quickly than elemental sulfur in supplying the needs of plants. In general, the practice of applying sulfur involves little risk. Applying too much, that is, enough to reduce yields, would be difficult. It should be noted, however, that applying moderate to high levels of elemental sulfur or using ammonium sulfate leaves an acid residue in the soil. For most crops an application of 10 to 20 pounds of sulfur per acre should correct a sulfur deficiency.

**Micronutrients.** The elements boron, manganese, iron, molybdenum, copper, zinc, and chlorine are essential for plant growth but needed in very small amounts. Green plants, for example, require about 150,000 times more nitrogen than molybdenum. Deficiencies, therefore, are quite rare and usually occur in fairly predictable situations. Maintenance of soil pH between 6.2 and 6.8 is the best means of avoiding both excesses and deficiencies of micronutrients. For information on when micronutrient deficiencies are likely to occur and how to avoid or correct them, see *Horticulture Facts VC-17-81*, "Micronutrient Applications for Vegetable Crops," available from the Department of Horticulture. Preventative applications on a routine basis are not recommended for most vegetables and soils.

## Crop Rotations and Organic Matter

Soil organic matter breaks down very rapidly in soils used for vegetable production because of the intensive cultivation and the frequent irrigation. Including grasses or legumes in a crop rotation at least once every three years helps maintain organic matter as well as the good structure and friability of the soil. Clay or silt loam soils with good structure will have improved drainage, and sands or sandy loams with good organic matter will hold water better.

Seeding a cover crop in the fall is a common practice on sandy soils that are subject to blowing. Loamy soils can also benefit from cover crops since root growth and the organic residues from these crops can improve the structure of the soil as well as reduce erosion. The most commonly used cover crops are oats, wheat, and rye.

1. Oats should be seeded at 60 to 100 pounds per acre during September or October to provide a good cover that will die out over the winter.
2. Winter wheat or rye should be seeded at 80 to 110 pounds per acre in September or October. These should be plowed down in the spring before they become too vegetative.

Green manure crops are grown during the same season as a cash crop for the purpose of improving the tilth of the soil. Most commercial vegetable operations cannot allow land to be nonproductive, but small farmers may use green manures as a source of nitrogen for subsequent crops. Common green manure crops are sweet clover, soybeans, sudan grass, and a mixture of peas and oats.

1. Sweet clover should be seeded at 15 to 25 pounds per acre in the spring. This may add as much as 100 pounds of nitrogen per acre during the next several years.

2. Soybeans should be inoculated and seeded at 120 pounds per acre.
3. Sudan grass may be seeded at 35 pounds per acre.
4. A mixture of peas at 65 pounds per acre and oats at 100 pounds per acre may be sown early in the spring and plowed down later to allow for a fall crop.

Generally, 30 to 40 pounds of nitrogen and all of the phosphorus and potassium required for the next crop should be applied before the cover crop or green manure. Approximately 30 percent of the nitrogen will be recovered by the first vegetable crop.

Animal manure is the most common type of organic matter and should be used whenever available. Manure can supply an appreciable amount of nitrogen and phosphorus. More nitrogen will be available if manure is applied in the spring, especially if it is immediately covered with soil to prevent the loss of ammonia nitrogen. Manure can also be injected to prevent the loss of nitrogen and to reduce the likelihood of the runoff of phosphorus into streams.

When more than one crop is to be grown in a field it is necessary to adjust the fertilizer application rates so that the nutrients needed by all of the crops are supplied. Tailoring a fertilizer program for such situations is difficult because the amount of a nutrient that is considered adequate for one crop may be undesirably low for another. For assistance, consult *Horticulture Facts* No. VC-7-80, "Fertilizer Guide for Market Gardeners." Other publications in this series that may be of interest to you are No. VC-8-80, "Conversion Tables for Fertilizer Calculations," and No. VC-18-81, "Liming Vegetable Crops." You can obtain these publications from your county Cooperative Extension Service adviser or from the Department of Horticulture, 124 Mumford Hall, 1301 West Gregory Drive, University of Illinois, Urbana, Illinois 61801.

---

**This circular was prepared by John M. Gerber, Extension Specialist in Vegetable Crops, and John M. Swiader, Assistant Professor of Horticulture, University of Illinois at Urbana-Champaign.**

---

Issued in furtherance of Cooperative Extension Work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. WILLIAM R. OSCHWALD, Director, Cooperative Extension Service, University of Illinois at Urbana-Champaign. The Illinois Cooperative Extension Service provides equal opportunities in programs and employment.