

Establishing a stand of grass requires proper planning and attention to detail. Perennial grasses differ in establishment requirements compared to annual grain crops. Five keys to successful grass seeding and establishment are presented in the following narrative. Adhering to these guidelines will greatly improve your chances of a successful grass stand.

Key #1—Seeding Date

Grasses should be seeded when soil moisture and temperature are optimum for germination. Grasses are designated either “cool” or “warm” season based on their growth cycle. Cool-season grasses can be planted when temperatures are cooler and day lengths shorter. Warm-season grasses need warmer temperatures and longer day lengths to grow. Following are recommended planting dates for cool-season and warm-season grasses in Indiana.

Note Table 1: Refer to appropriate technical standard for more specific recommendations

Table 1 Indiana Recommended Planting Dates	
Cool Season Grasses	Warm Season Grasses
Feb 1—May 1 Aug 1-Nov 1	Apr 15—May 30

Key #2—Seedbed

A proper seedbed is firm and free of competing vegetation. Correct firmness is when an adult footprint is only slightly visible on the prepared bed prior to the seeding operation. The seedbed can be firmed, if needed, by pull-



ing a commercial or homemade packer or roller. A firm seedbed is essential for proper seeding depth. A loose, fluffy bed will place seeds too deep for proper germination. Seed requiring light for germination will be hindered by a deep planting depth. Seed that germinates but does not have enough nutrient reserve for the shoot to reach the surface is also hindered by a deep planting depth. Most species should be planted at a shallow depth of ¼ to ½ inch. Larger seeds can be planted up to 1 inch deep. Most seedings are too deep if you cannot see a few seeds on the soil surface.

Grasses can be successfully seeded into a tilled or no-tilled seedbed, provided weeds are controlled and residue is managed prior to planting. Weeds compete with seedlings for moisture and light. Optimum control comes with several years of weed management prior to seeding. At seeding time, there should be no actively growing weeds. Weeds can be controlled with tillage and/or herbicides applied before or just after seeding. Like a weed, companion crops can compete with the seeded species for water and light. Unless erosion is a problem, companion crops are generally not recommended in grass seedings. Residue affects seeding depth and seed soil contact. Tillage, fire, and mowing can be used to manage residue prior to seeding. Tolerable residue amounts are dependent on seeding equipment to be used. Residue should be harrowed to spread extra chaff and straw. Late summer and dormant seedings are best planted into standing stubble.

Table 2 Indiana Grass Seeding Rates					
Applicable Technical Standards					
	Conservation Cover 327	Filter Strip 393	Pasture and Hayland 512	Critical Area Planting 342	Recommended Planting Depth (inches)
Recommended Seeding Rates (lb PLS/acre)					
Cool Season Grasses					
Creeping red fescue				10-20	0.25-0.5
Kentucky bluegrass	2-10	1	2-4		0.25-0.5
Orchardgrass	4-10	3-8	4-6	13-20	0.25-0.5
Perennial ryegrass	15-20			5-10	0.25-0.5
Red top	3-6	0.4-2		1-2	0.5
Reed canarygrass			3-5		0.25-0.5
Smooth bromegrass	5-8	10-40	5-7		0.5-1
Tall fescue	8-15	10-25	6-10	25-50	0.25-0.5
Timothy	2-6	0.5-1	2-4		0.25-0.5
Wildrye spp.		4			0.25-0.5
Warm Season Grasses					
Big bluestem	8				0.25-0.5
Indiangrass	8				0.25-0.5
Little bluestem		4-6			0.25-0.5
Side-oats grama		1-5			0.25-0.5
Switchgrass	4	3-8		8-10	0.25-0.5

Note: Seeding rate ranges under each standard are based on the proportion of each species in a given seed mixture. Refer to listed technical standards for recommended seeding mixes.

Key #3—Seed Placement

The seeding equipment should provide proper seed depth, uniform seeding rate, and good seed to soil contact. Grass seed can be broadly categorized into three types: fluffy or chaffy, smooth small seed, and smooth large seed. Grass drills are equipped with separate boxes to properly place and meter each of the three seed types. Picker wheels and agitators in the fluffy/chaffy box and oversized feeder tubes keep rough coated seed flowing evenly. Depth bands on grass drills are essential for planting depth control. Press/packer wheels contribute to close seed/soil contact. Free flowing grass seed (e.g. wheatgrass) can be successfully planted with a small grain drill if proper, shallow, and consistent seeding depth is maintained.



Picker wheel



Agitators



Press/packer wheel

Drills should be calibrated to determine seeding rate. Seeding rate can be determined by counting dropped seeds after traveling a given distance on a hard surface, collecting seed from openers after traveling a given distance, or turning the drive wheel on the drill and collecting seed from openers. Contact the local NRCS office for additional information. See Table 2 for seeding rate ranges and planting depth of recommended grass species.

Key #4—Seed Quality

All seed must meet the requirement of the States' seed laws. The seed should be tested for purity and germination. Purity specifies any weeds and inert matter in the seed lot. Germination is an indication of the percentage of seed that will sprout and grow. Seed is usually purchased and planted on a Pure Live Seed (PLS) basis. This

is calculated by multiplying purity by germination (including dormant). A high PLS usually indicates high quality seed. Seed of adapted species and recommended cultivars within the species should be planted. It is best to select cultivars whose origin is closest to the planting site when seeding warm-season grasses. Cool-season species are more broadly adapted. Your local NRCS office can provide information on adapted species, varieties, and seeding rates. See Table 2 for seeding rate ranges and planting depth of recommended grass species.

Seed with awns or other appendages is called “fluffy” or bearded. Debearded seed has part or all of the appendages removed and is more flowable. Flowability depends on degree of debearding.

Key #5—Weed Control

Weeds compete for moisture and light with young seedlings. Competitive weeds can be controlled mechanically by clipping or chemically. Dense residue clippings should be removed from the seeded area. Weeds should be controlled with herbicides before they reach 4 inches tall. Carefully read and follow all instructions on pesticide labels.

The two main reasons grass seedlings fail are planting too deep and lack of weed con-

For more information, contact:
USDA-NRCS

Rose Lake Plant Materials Center
7472 Stoll Road

East Lansing, MI 48823

Phone: (517) 641-6300

Fax: (517) 641-4421

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202)720-2600 (voice and TDD). To file a complaint, write USDA, Director, Office of Civil Rights, Room 326W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202)720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

February 2005



United States Department of Agriculture
Natural Resources Conservation Service
Rose Lake Plant Materials Center
East Lansing, Michigan

Five Keys to Successful Grass Seeding in Indiana

☞ Seeding Date

☞ Seedbed

☞ Seed Placement

☞ Seed Quality

☞ Weed Control

