

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

Map Unit Symbol	Map unit name	Corn	Soy-beans	Winter Wheat	Grass-Legume Hay	Pasture
		Bu	Bu	Bu/Ac	Tons	AUM
AddA	Avonburg silt loam, 0 to 2 percent slopes	145	48	65	4.8	9.6
AddB2	Avonburg silt loam, 2 to 4 percent slopes, eroded	140	47	60	4.4	8.8
AfsB	Alvin-Princeton fine sandy loams, 2 to 6 percent slopes	124	43	48	3.9	7.9
AfsC2	Alvin-Princeton fine sandy loams, 6 to 12 percent slopes, eroded	111	39	43	3.6	7.2
AmkA	Ayrshire fine sandy loam, 0 to 2 percent slopes	150	49	60	5.0	10.0
BbhA	Bartle silt loam, 0 to 2 percent slopes	145	49	59	4.8	9.6
BbiB	Bartle-Pekin silt loams, 2 to 6 percent slopes	143	49	58	4.8	9.5
BcrAW	Beanblossom silt loam, 1 to 3 percent slopes, occasionally flooded, very brief duration	106	37	15	0.0	0.0
BdhAH	Bellcreek silty clay loam, 0 to 1 percent slopes, frequently flooded, brief duration	125	33	0	0.0	0.0
BfbAH	Bellcreek silt loam, 0 to 1 percent slopes, frequently flooded, brief duration	125	33	0	0.0	0.0
BgeAHU	Birds silt loam, undrained, 0 to 1 percent slopes, frequently flooded, brief duration	0	0	0	0.0	0.0
BgeAW	Birds silt loam, 0 to 1 percent slopes, occasionally flooded, very brief duration	145	40	20	0.0	0.0
BlgC2	Blocher-Cincinnati silt loams, 6 to 12 percent slopes, eroded	126	44	55	4.3	8.6
BlgC3	Blocher-Cincinnati silt loams, 6 to 12 percent slopes, severely eroded	118	41	49	3.8	7.6
BlhD2	Blocher-Bonnell silt loams, 12 to 25 percent slopes, eroded	64	22	35	2.8	5.6

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

BluC	Bloomfield-Alvin loamy sands, 6 to 12 percent slopes	92	32	35	2.9	5.8
BnuD3	Bonnell-Hickory-Blocher complex, 12 to 25 percent slopes, severely eroded	35	12	5	0.4	0.7
BobE5	Bonnell-Hickory clay loams, 15 to 30 percent slopes, gullied	---	---	---	0.0	0.1
BodAV	Bonnie silt loam, 0 to 1 percent slopes, frequently flooded, very brief duration	130	36	20	0.0	0.0
CldB2	Cincinnati-Blocher silt loams, 2 to 6 percent slopes, eroded	130	46	59	4.3	8.6
ClfA	Cobbsfork silt loam, 0 to 1 percent slopes	139	43	61	5.0	10.1
CmbAW	Cohoctah loam, 0 to 1 percent slopes, occasionally flooded, very brief duration	155	44	22	0.0	0.0
CmzA	Cliftycreek silt loam, 0 to 2 percent slopes	145	51	58	4.8	9.6
CmzB2	Cliftycreek silt loam, 2 to 6 percent slopes, eroded	140	49	56	4.6	9.2
CmzC2	Cliftycreek silt loam, 6 to 12 percent slopes, eroded	130	46	52	4.3	8.6
ColD2	Coolville-Rarden-Stonehead silt loams, 12 to 18 percent slopes, eroded	80	28	31	2.5	5.0
ConC3	Coolville-Rarden complex, 6 to 12 percent slopes, severely eroded	75	26	32	2.7	5.3
CudA	Crosby silt loam, 0 to 2 percent slopes	143	46	63	4.6	9.2
CulB	Crosby-Williamstown silt loams, 2 to 4 percent slopes	142	47	63	4.6	9.3
CxdA	Cyclone silty clay loam, 0 to 2 percent slopes	185	56	77	6.3	12.6
DbqE	Deam silt loam, very deep, 15 to 30 percent slopes	95	33	38	3.1	6.2

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

EcyAH	Eel loam, 0 to 2 percent slopes, frequently flooded, brief duration	120	42	0	0.0	0.0
EcyAW	Eel loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	135	47	19	0.0	0.0
EdeAW	Eel silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	135	47	19	0.0	0.0
EepAQ	Elkinsville silt loam, 0 to 2 percent slopes, rarely flooded	150	52	60	5.0	10.0
FdbA	Fincastle silt loam, 0 to 2 percent slopes	166	54	74	5.4	10.8
FdqB	Fincastle-Xenia silt loams, 2 to 4 percent slopes	161	54	73	5.4	10.7
FexA	Fox loam, 0 to 2 percent slopes	109	38	50	3.3	6.7
FexAQ	Fox loam, 0 to 2 percent slopes, rarely flooded	100	35	50	3.3	6.6
FexB2	Fox loam, 2 to 6 percent slopes, eroded	102	35	48	3.1	6.3
FgqC3	Fox-Casco sandy loams, 6 to 12 percent slopes, severely eroded	69	24	36	2.4	4.8
GccAH	Genesee loam, 0 to 2 percent slopes, frequently flooded, brief duration	119	42	0	0.0	0.0
GccAW	Genesee loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	135	47	19	0.0	0.0
GcpAW	Genesee silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	135	47	19	0.0	0.0
GgbG	Gilwood-Brownstown silt loams, 25 to 75 percent slopes	---	---	---	0.1	0.1
GgfD2	Gilwood-Wrays silt loams, 12 to 25 percent slopes, eroded	84	29	32	2.6	5.3
HcgAW	Haymond silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	134	47	19	0.0	0.0

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

HctAW	Haymond-Wirt silt loams, 0 to 2 percent slopes, occasionally flooded, very brief duration	128	45	18	0.0	0.0
HeoF	Hickory silt loam, 25 to 50 percent slopes	0	0	0	0.0	0.0
HleAW	Holton silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	131	39	18	0.0	0.0
KugG	Kurtz-Gnawbone silt loams, 20 to 60 percent slopes	0	0	0	0.0	0.0
LeaA	Lauer silt loam, 0 to 2 percent slopes	170	56	68	5.6	11.2
MecAQ	Martinsville loam, 0 to 2 percent slopes, rarely flooded	145	51	58	4.8	9.6
MecB	Martinsville loam, 2 to 6 percent slopes	145	51	58	4.8	9.6
MfwA	Martinsville loam, sandy substratum, 0 to 2 percent slopes	145	51	73	4.8	9.6
MfwAQ	Martinsville loam, sandy substratum, 0 to 2 percent slopes, rarely flooded	145	51	73	4.8	9.6
MfwB2	Martinsville loam, sandy substratum, 2 to 6 percent slopes, eroded	140	49	70	4.6	9.2
MfxA	Martinsville sandy loam, sandy substratum, 0 to 2 percent slopes	125	44	63	4.1	8.2
MhuA	McGary silt loam, 0 to 2 percent slopes	140	46	56	4.6	9.2
MhyB	Medora silt loam, 2 to 6 percent slopes	130	46	52	4.3	8.6
MhyC2	Medora silt loam, 6 to 12 percent slopes, eroded	115	40	46	3.8	7.6
MjjAH	Medway silty clay loam, 0 to 2 percent slopes, frequently flooded, brief duration	130	46	0	0.0	0.0
MmoC3	Miami clay loam, 6 to 12 percent slopes, severely eroded	121	42	54	4.0	8.0

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

MmoD3	Miami clay loam, 12 to 18 percent slopes, severely eroded	110	39	50	3.6	7.2
MnpB2	Miami silt loam, 2 to 6 percent slopes, eroded	142	49	63	4.6	9.2
MnpC2	Miami silt loam, 6 to 12 percent slopes, eroded	132	46	58	4.3	8.6
MnpD2	Miami silt loam, 12 to 18 percent slopes, eroded	116	41	52	3.8	7.6
MqbA	Milton silt loam, 0 to 2 percent slopes	100	35	40	3.3	6.6
MqbA	Milton silt loam, 0 to 2 percent slopes	102	35	40	3.3	6.6
MqbB2	Milton silt loam, 2 to 6 percent slopes, eroded	95	33	38	3.1	6.2
MqbC2	Milton silt loam, 6 to 12 percent slopes, eroded	85	30	34	2.8	5.6
MrbF	Milton-Rock outcrop complex, 25 to 40 percent slopes	0	0	0	0.0	0.0
NaaB2	Nabb silt loam, 2 to 6 percent slopes, eroded	117	41	50	3.6	7.3
NpcA	Nineveh gravelly sandy loam, 0 to 2 percent slopes	105	37	53	3.5	7.0
NpcAQ	Nineveh gravelly sandy loam, 0 to 2 percent slopes, rarely flooded	105	37	53	3.5	7.0
NpeA	Nineveh sandy loam, 0 to 2 percent slopes	105	37	53	3.5	7.0
NpeAQ	Nineveh sandy loam, 0 to 2 percent slopes, rarely flooded	105	37	53	3.5	7.0
NpeB2	Nineveh sandy loam, 2 to 6 percent slopes, eroded	100	35	50	3.3	6.6
ObaA	Ockley loam, 0 to 2 percent slopes	132	46	67	4.5	8.9
OfaAW	Oldenburg silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	125	43	18	0.0	0.0
Omz	Orthents, earthen dam	0	0	0	0.0	0.0

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

PcrB2	Pekin silt loam, 2 to 6 percent slopes, eroded	136	48	54	4.5	9.0
PcrC2	Pekin silt loam, 6 to 12 percent slopes, eroded	124	43	49	3.9	7.9
PcrC3	Pekin silt loam, 6 to 12 percent slopes, severely eroded	120	42	48	4.0	8.0
PhaA	Peoga silt loam, 0 to 1 percent slopes	134	41	59	4.9	9.8
PlpAV	Piopolis silty clay loam, 0 to 1 percent slopes, frequently flooded, very brief duration	135	37	19	0.0	0.0
Pml	Pits, quarry	0	0	0	0.0	0.0
PnnD	Pike-Chetwynd silt loams, 12 to 20 percent slopes	117	41	48	4.0	7.9
PnnF	Pike-Chetwynd silt loams, 20 to 50 percent slopes	0	0	0	0.0	0.0
Ppu	Pits, sand and gravel	0	0	0	0.0	0.0
RctD3	Rarden-Coolville complex, 12 to 22 percent slopes, severely eroded	56	20	21	1.7	3.5
RehA	Rensselaer-Treaty silt loams, 0 to 1 percent slopes	175	51	70	5.8	11.6
ReyA	Rensselaer loam, 0 to 1 percent slopes	175	51	70	5.8	11.6
ReyAQ	Rensselaer loam, 0 to 1 percent slopes, rarely flooded	175	51	70	5.8	11.6
RqaG	Rodman sandy loam, 25 to 50 percent slopes	0	0	0	0.0	0.0
RtxAH	Rosburg silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	130	46	0	0.0	0.0
RtxAK	Rosburg silt loam, 0 to 2 percent slopes, occasionally flooded, brief duration	135	47	19	0.0	0.0
RywB2	Russell silt loam, 2 to 6 percent slopes, eroded	150	52	67	5.0	10.0

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

SfyA	Shircliff silt loam, 0 to 2 percent slopes	130	46	52	4.3	8.6
SifE	Senachwine loam, 18 to 25 percent slopes	---	---	---	0.0	0.0
SifG	Senachwine loam, 25 to 70 percent slopes	---	---	---	0.0	0.0
SldAH	Shoals silt loam, 0 to 2 percent slopes, frequently flooded, brief duration	130	42	1	0.0	0.0
SldAW	Shoals silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	145	47	20	0.0	0.0
SnfA	Sleeth loam, 0 to 2 percent slopes	145	47	73	4.8	9.6
SoaB	Spickert silt loam, 2 to 6 percent slopes	125	44	50	4.1	8.2
SocAH	Sloan silty clay loam, 0 to 1 percent slopes, frequently flooded, brief duration	140	39	0	0.0	0.0
SocAW	Sloan silty clay loam, 0 to 1 percent slopes, occasionally flooded, very brief duration	155	44	22	0.0	0.0
SoeC2	Spickert-Wrays silt loams, 6 to 18 percent slopes, eroded	106	37	44	3.6	7.1
SolC2	Spickert-Wrays silt loams, 6 to 12 percent slopes, eroded	104	36	41	3.3	6.7
SolC3	Spickert-Wrays silt loams, 6 to 12 percent slopes, severely eroded	99	35	39	3.2	6.4
StaAV	Steff silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	130	46	18	0.0	0.0
StdAQ	Stendal silt loam, 0 to 2 percent slopes, rarely flooded	150	49	60	5.0	10.0
StdAV	Stendal silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	135	44	19	0.0	0.0
StmB	Stonehead silt loam, 2 to 6 percent slopes	145	51	58	4.8	9.6
SucC2	Stonehead-Coolville silt loams, 6 to 12 percent slopes, eroded	123	43	51	4.2	8.5

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

SujD5	Stonehead silt loam, 10 to 20 percent slopes, gullied	0	0	0	0.0	0.0
SulC2	Stonehead-Wellrock silt loams, 6 to 15 percent slopes, eroded	122	43	51	4.2	8.4
SuoAH	Stonelick fine sandy loam, 0 to 2 percent slopes, frequently flooded, brief duration	90	32	0	0.0	0.0
Uaz	Udorthents, sandy	0	0	0	0.0	0.0
Uby	Udorthents, loamy	0	0	0	0.0	0.0
UemB	Urban land-Alvin-Princeton complex, 2 to 6 percent slopes	0	0	0	0.0	0.0
UemC	Urban land-Alvin-Princeton complex, 6 to 12 percent slopes	0	0	0	0.0	0.0
UenA	Urban land-Fox complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UenB	Urban land-Fox complex, 2 to 6 percent slopes	0	0	0	0.0	0.0
UepC	Urban land-Fox-Casco complex, 6 to 12 percent slopes	0	0	0	0.0	0.0
UfcB	Urban land-Cincinnati-Nabb complex, 2 to 12 percent slopes	0	0	0	0.0	0.0
UfdA	Urban land-Cobbsfork-Avonburg complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UfnA	Urban land-Crosby-Treaty complex, fine loamy subsoil, 0 to 2 percent slopes	0	0	0	0.0	0.0
UfoA	Urban land-Cyclone complex, 0 to 1 percent slopes	0	0	0	0.0	0.0
UfxA	Urban land-Fincastle complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UfyB	Urban land-Fincastle-Russell complex, 2 to 6 percent slopes	0	0	0	0.0	0.0

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

UhyA	Urban land-Martinsville, sandy substratum, complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UkbC	Urban land-Miami complex, 6 to 12 percent slopes	0	0	0	0.0	0.0
UkpA	Urban land-Ockley complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UkqA	Urban land-Nineveh complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UkqB	Urban land-Nineveh complex, 2 to 6 percent slopes	0	0	0	0.0	0.0
UmqA	Urban land-Sleeth complex, 0 to 2 percent slopes	0	0	0	0.0	0.0
UnnA	Urban land-Westland complex, 0 to 1 percent slopes	0	0	0	0.0	0.0
Usl	Udorthents, rubbish	0	0	0	0.0	0.0
W	Water	0	0	0	0.0	0.0
WaaAV	Wakeland silt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	140	42	20	0.0	0.0
WaaAW	Wakeland silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	146	44	20	0.0	0.0
WacAW	Wakeland-Birds silt loams, 0 to 2 percent slopes, occasionally flooded, very brief duration	145	43	20	0.0	0.0
WbiAW	Wilbur-Wakeland silt loams, 0 to 2 percent slopes, occasionally flooded, very brief duration	142	47	20	0.0	0.0
WdlC2	Wawaka loam, 6 to 12 percent slopes, eroded	130	46	52	4.3	8.6
WdrB2	Wawaka silt loam, 2 to 6 percent slopes, eroded	140	49	56	4.6	9.2

NON-IRRIGATED MAP UNIT CROP YIELDS (RV)  
 Bartholomew County, Indiana

WokAW	Wilbur silt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	140	48	20	0.0	0.0
WolAV	Wilhite silty clay, 0 to 1 percent slopes, frequently flooded, very brief duration	120	32	17	0.0	0.0
WprAV	Wirt loam, 0 to 2 percent slopes, frequently flooded, very brief duration	117	41	16	0.0	0.0
WprAW	Wirt loam, 0 to 2 percent slopes, occasionally flooded, very brief duration	121	42	17	0.0	0.0
WqlA	Westland clay loam, 0 to 1 percent slopes	165	47	83	5.4	10.8
WqlAQ	Westland clay loam, 0 to 1 percent slopes, rarely flooded	165	47	83	5.4	10.8
WsuA	Whitaker loam, 0 to 2 percent slopes	155	51	62	5.1	10.2
WsyAQ	Whitaker sandy loam, 0 to 2 percent slopes, rarely flooded	150	49	60	5.0	10.0
WufB2	Williamstown silt loam, 2 to 6 percent slopes, eroded	142	49	63	4.6	9.2
XabB2	Xenia silt loam, 2 to 6 percent slopes, eroded	153	53	67	5.0	10.0
XfuB2	Miami-Rainsville silt loams, 2 to 6 percent slopes, eroded	140	49	63	4.6	9.2
XrbC2	Miami-Rainsville loams, 6 to 12 percent slopes, eroded	134	47	59	4.3	8.7
XrkD2	Miami-Kendallville loams, 12 to 18 percent slopes, eroded	119	42	50	3.8	7.7
ZboA	Zipp silty clay loam, 0 to 1 percent slopes	150	42	60	5.0	10.0