

NON-IRRIGATED COMPONENT CROP YIELDS (RV)
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
AvA	Avonburg silt loam, 0 to 2 percent slopes	Avonburg	85	145	49	65	4.8	9.6
		Cobbsfork	10	150	46	60	5.0	10.0
		Nabb	5	130	46	59	4.3	8.6
Ba	Bartle silt loam, 0 to 3 percent slopes	Bartle	90	140	47	63	4.6	9.2
		poorly drained aqualfs	10	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
BbhA	Bartle silt loam, 0 to 2 percent slopes	Bartle	83	145	49	58	4.8	9.6
		Peoga	10	135	40	54	4.5	9.0
		Pekin	5	115	40	46	3.8	7.6
		Bartle	2	120	40	48	4.0	8.0
BbiB	Bartle-Pekin silt loams, 2 to 6 percent slopes	Bartle	55	145	49	58	4.8	9.6
		Pekin	35	140	49	56	4.6	9.2
		Elkinsville	5	135	47	54	4.5	9.0
		Peoga	5	135	40	54	4.5	9.0
BcrAW	Beanblossom silt loam, 1 to 3 percent slopes, occasionally flooded, very brief duration	Beanblossom	89	110	39	15	3.6	7.2
		Wilbur	5	140	49	20	4.6	9.2
		Beanblossom	3	105	37	15	3.5	7.0
		A deep, loamy soil	3	145	44	20	4.8	9.6
Be	Beanblossom channery silt loam, occasionally flooded	Beanblossom	100	100	35	14	3.3	6.6
BeG	Berks channery silt loam, 25 to 75 percent slopes	Berks	100	---	---	---	0.0	0.0

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
BgF	Berks-Trevlac-Wellston complex, 20 to 70 percent slopes	Berks	50	---	---	---	0.0	0.0
		Trevlac	25	---	---	---	0.0	0.0
		Wellston	25	75	26	30	2.5	5.0
BlgC2	Blocher-Cincinnati silt loams, 6 to 12 percent slopes, eroded	Blocher	54	135	47	54	4.5	9.0
		Cincinnati	35	120	42	54	4.0	8.0
		Blocher	5	115	40	46	3.8	7.6
		Cincinnati	3	95	33	43	3.1	6.2
		Wakeland	2	135	40	19	4.5	9.0
		Bonnell	1	---	---	---	2.5	5.0
BlhD2	Blocher-Bonnell silt loams, 12 to 25 percent slopes, eroded	Blocher	40	120	42	48	4.0	8.0
		Bonnell	30	---	---	---	2.6	5.2
		Hickory	12	---	---	---	3.1	6.2
		Blocher	6	115	40	46	3.8	7.6
		Cincinnati	5	100	35	45	3.3	6.6
		Bonnell, severely eroded	4	---	---	---	2.1	4.2
		Wakeland	3	135	40	19	4.5	9.0
BnD2	Bonnell silt loam, 12 to 20 percent slopes, eroded	Bonnell	100	100	35	40	3.3	6.6
BpD3	Bonnell clay loam, 12 to 20 percent slopes, gullied	Bonnell, very severely eroded	100	0	0	0	0.0	0.0
CdD2	Chetwynd loam, 12 to 20 percent slopes, eroded	Chetwynd	100	105	37	42	3.5	7.0
CdF	Chetwynd loam, 20 to 50 percent slopes	Chetwynd	100	---	---	---	0.0	0.0

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
CnC2	Cincinnati silt loam, Wabash Lowland, 6 to 12 percent slopes, eroded	Cincinnati, eroded	90	105	37	47	3.5	7.0
		Cincinnati, severely eroded	5	100	35	45	3.3	6.6
		Stendal, rarely, very brief	5	150	49	60	5.0	10.0
CwB	Crosby silt loam, 1 to 5 percent slopes	Crosby	85	155	51	70	5.1	10.2
		Miami, eroded	5	140	49	63	4.6	9.2
		Treaty, drained	5	185	54	83	6.1	12.2
		Williamstown, eroded	5	140	49	63	4.6	9.2
FdbA	Fincastle silt loam, 0 to 2 percent slopes	Fincastle	84	165	54	74	5.4	10.8
		Typic Argiaquolls	10	Not Rated	54	74	Not Rated	Not Rated
		Williamstown	6	130	46	59	4.3	8.6
FdqB	Fincastle-Xenia silt loams, 2 to 4 percent slopes	Fincastle	50	165	54	74	5.4	10.8
		Xenia	40	155	54	70	5.1	10.2
		Typic Argiaquolls	10	Not Rated	54	74	Not Rated	Not Rated
GgbG	Gilwood-Brownstown silt loams, 25 to 75 percent slopes	Gilwood	45	---	---	---	0.0	0.0
		Brownstown	35	---	---	---	0.0	0.0
		Wrays	10	100	35	40	3.3	6.6
		Gilwood	3	80	28	32	2.6	5.2
		Shallow, loamy soil	3	---	---	---	0.0	0.0
		Beanblossom	2	105	37	15	3.5	7.0
		Rock outcrop	2	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
GgFD2	Gilwood-Wrays silt loams, 12 to 25 percent slopes, eroded	Gilwood	40	75	26	30	2.5	5.0
		Wrays	32	95	33	38	3.1	6.2
		Gilwood, severely eroded	10	70	25	28	2.3	4.6
		Wrays, severely eroded	10	90	32	36	3.0	6.0
		Brownstown	4	---	---	---	0.0	0.0
		Beanblossom	2	105	37	15	3.5	7.0
		Spickert	2	100	35	40	3.3	6.6
GnF	Gilpin silt loam, 25 to 55 percent slopes	Gilpin	100	---	---	---	0.0	0.0
GpD	Gilpin-Wellston silt loams, 10 to 25 percent slopes	Gilpin	50	65	23	26	2.1	4.2
		Wellston	50	100	35	40	3.3	6.6
Hc	Haymond silt loam, frequently flooded	Haymond	100	120	42	0	0.0	0.0
HeoF	Hickory silt loam, 25 to 50 percent slopes	Hickory	70	---	---	---	0.0	0.0
		Blocher	10	---	---	---	0.0	0.0
		Bonnell	10	---	---	---	0.0	0.0
		Hickory	5	---	---	---	0.0	0.0
		Holton	5	120	35	17	4.0	8.0
HkD2	Hickory silt loam, 12 to 20 percent slopes, eroded	Hickory	100	115	40	46	3.8	7.6
HkF	Hickory silt loam, 20 to 70 percent slopes	Hickory	100	---	---	---	0.0	0.0

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
KugG	Kurtz-Gnawbone silt loams, 20 to 60 percent slopes	Kurtz	53	---	---	---	0.0	0.0
		Gnawbone	32	---	---	---	0.0	0.0
		Beanblossom	3	100	35	14	3.3	6.6
		Kurtz-Gnawbone	3	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
		Brownstown	2	---	---	---	0.0	0.0
		Coolville	2	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
		Very deep, well drained colluvial soil	2	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
		Wellrock	2	100	35	40	3.3	6.6
		Stonehead	1	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
MnC2	Miami loam, 6 to 15 percent slopes, eroded	Miami	100	130	46	59	4.3	8.6
Omz	Orthents, earthen dam	Orthents	100	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
PcrC2	Pekin silt loam, 6 to 12 percent slopes, eroded	Pekin	72	125	44	50	4.1	8.2
		Pekin, severely eroded	14	95	33	38	3.1	6.2
		Elkinsville	5	120	42	48	4.0	8.0
		Pekin	5	85	30	34	2.8	5.6
		Stendal	4	135	44	19	4.5	9.0
PeB	Pekin silt loam, 2 to 6 percent slopes	Pekin	95	130	46	59	4.3	8.6
		Bartle	3	135	46	61	4.5	9.0
		Peoga	2	155	47	62	5.1	10.2

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
PeC2	Pekin silt loam, 6 to 12 percent slopes, eroded	Pekin, eroded	95	115	40	52	3.8	7.6
		Elkinsville, eroded	3	135	47	54	4.5	9.0
		Bartle	1	135	46	61	4.5	9.0
		Peoga	1	155	47	62	5.1	10.2
RoB2	Nabb silt loam, 2 to 6 percent slopes, eroded	Nabb, eroded	78	115	40	52	3.8	7.6
		Cincinnati, eroded	10	115	40	52	3.8	7.6
		Avonburg	8	130	44	59	4.3	8.6
		Wakeland	4	145	44	20	4.8	9.6
Sf	Steff silt loam, frequently flooded	Steff	100	110	39	0	0.0	0.0
SifG	Senachwine loam, 25 to 70 percent slopes	Senachwine	90	---	---	---	0.0	0.0
		Senachwine, severely eroded	5	---	---	---	0.0	0.0
		Shoals	5	135	44	19	4.5	9.0
SoaB	Spickert silt loam, 2 to 6 percent slopes	Spickert	85	125	44	50	4.1	8.2
		Wrays	10	110	39	44	3.6	7.2
		Very deep, somewhat poorly drained, silty soil	5	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
SoeC2	Spickert-Wrays silt loams, 6 to 18 percent slopes, eroded	Spickert	60	110	39	44	3.6	7.2
		Wrays	25	95	33	38	3.1	6.2
		Gilwood	5	75	26	30	2.5	5.0
		Spickert	5	95	33	38	3.1	6.2
		Wrays	5	105	37	42	3.5	7.0

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
St	Stendal silt loam, frequently flooded	Stendal	90	125	40	0	0.0	0.0
		poorly drained aquents	3	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
StmB	Stonehead silt loam, 2 to 6 percent slopes	Stonehead	85	145	51	58	4.8	9.6
		Coolville	5	95	33	38	3.1	6.2
		Stonehead	5	120	42	48	4.0	8.0
		Weddel	5	125	44	50	4.1	8.2
SulC2	Stonehead-Wellrock silt loams, 6 to 15 percent slopes, eroded	Stonehead	55	130	46	52	4.3	8.6
		Wellrock	25	105	37	42	3.5	7.0
		Coolville	7	105	37	42	3.5	7.0
		Gnawbone	5	---	---	---	0.0	0.0
		Kurtz	5	---	---	---	0.0	0.0
		Stonehead	3	145	51	58	4.8	9.6
Sv	Stendal silt loam, frequently flooded, very long duration	Stendal	100	50	14	0	0.0	0.0
SwC2	Stonehead silt loam, 6 to 10 percent slopes, eroded	Stonehead	100	130	46	52	4.3	8.6
SwD3	Stonehead silt loam, 10 to 20 percent slopes, gullied	Stonehead, very severely eroded	100	0	0	0	0.0	0.0
SxD2	Stonehead-Trevlac silt loams, 10 to 20 percent slopes, eroded	Stonehead	65	105	37	42	3.5	7.0
		Trevlac	35	75	26	30	2.5	5.0
TlB	Tilsit silt loam, 2 to 6 percent slopes	Tilsit	100	125	44	56	4.1	8.2

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
Ud	Udorthents, loamy	Udorthents, loamy	100	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
W	Water	Water	100	Not Rated	Not Rated	Not Rated	Not Rated	Not Rated
WacAW	Wakeland-Birds silt loams, 0 to 2 percent slopes, occasionally flooded, very brief duration	Wakeland	59	145	44	20	4.8	9.6
		Birds	29	145	40	20	4.8	9.6
		Wilbur	10	130	46	18	4.3	8.6
		Wakeland	2	130	39	18	4.3	8.6
WaD	Wellston-Berks-Trevlac complex, 6 to 20 percent slopes	Wellston	40	110	39	44	3.6	7.2
		Berks	35	50	18	20	1.7	3.4
		Trevlac	25	50	18	20	1.7	3.4
WbiAW	Wilbur-Wakeland silt loams, 0 to 2 percent slopes, occasionally flooded, very brief duration	Wilbur	54	140	49	20	4.6	9.2
		Wakeland	30	145	44	20	4.8	9.6
		Birds	4	145	40	20	4.8	9.6
		Beanblossom	3	100	35	14	3.3	6.6
		Haymond	3	125	44	18	4.1	8.2
		Wakeland	3	130	39	18	4.3	8.6
		Wilbur	3	125	44	18	4.1	8.2
WeC2	Wellston-Gilpin silt loams, 6 to 20 percent slopes, eroded	Wellston	58	120	42	48	4.0	8.0
		Gilpin	42	75	26	30	2.5	5.0
WeD2	Wellston silt loam, 12 to 18 percent slopes, eroded	Wellston	100	95	33	38	3.1	6.2
Wt	Wilbur silt loam, frequently flooded	Wilbur	100	125	44	0	0.0	0.0

COMPONENT CROP YIELDS (RV)--Continued
 Brown County, Indiana

Map Unit Symbol	Map unit name	Component	Comp. Pct.	Corn Bu/Ac	Soybeans Bu/Ac	Winter Wheat Bu/Ac	Grass-Legume Hay Tons/AC	Pasture AUM
XfuB2	Miami-Rainsville silt loams, 2 to 6 percent slopes, eroded	Miami	60	140	49	63	4.6	9.2
		Rainsville	30	140	49	63	4.6	9.2
		Crosby	3	140	46	63	4.6	9.2
		Fincastle	3	165	54	74	5.4	10.8
		Miami, severely eroded	3	135	47	61	4.5	9.0
		Typic Argiaquolls	1	Not Rated	54	74	Not Rated	Not Rated
XrbC2	Miami-Rainsville loams, 6 to 12 percent slopes, eroded	Miami	55	130	46	59	4.3	8.6
		Rainsville	40	140	49	63	4.6	9.2
		Typic Argiaquolls	5	Not Rated	54	74	Not Rated	Not Rated
XrkD2	Miami-Kendallville loams, 12 to 18 percent slopes, eroded	Miami	60	115	40	52	3.8	7.6
		Kendallville	40	125	44	50	4.1	8.2