

NON-IRRIGATED COMPONENT CROP YIELDS (RV)
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|--|------------------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| AlB2 | Alford silt loam, 2 to 5 percent slopes, eroded | Alford, eroded | 95 | 155 | 54 | 62 | 5.1 | 10.2 |
| | | Hosmer, eroded | 5 | 120 | 42 | 54 | 4.0 | 8.0 |
| AlC2 | Alford silt loam, 5 to 10 percent slopes, eroded | Alford, eroded | 90 | 145 | 51 | 58 | 4.8 | 9.6 |
| | | Hosmer, eroded | 6 | 110 | 39 | 50 | 3.6 | 7.2 |
| | | Alvin | 2 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Wakeland, frequently flooded | 2 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| AnB | Alvin-Bloomfield complex, 2 to 6 percent slopes | Alvin | 60 | 105 | 37 | 42 | 3.5 | 7.0 |
| | | Bloomfield | 40 | 90 | 32 | 36 | 3.0 | 6.0 |
| AnC | Alvin-Bloomfield complex, 6 to 12 percent slopes | Alvin | 55 | 95 | 33 | 38 | 3.1 | 6.2 |
| | | Bloomfield | 45 | 85 | 30 | 34 | 2.8 | 5.6 |
| Ao | Ambraw sandy clay loam, rarely flooded | Ambraw | 100 | 170 | 49 | 68 | 5.6 | 11.2 |
| Ar | Armiesburg silt loam, occasionally flooded | Armiesburg | 100 | 135 | 47 | 19 | 4.5 | 9.0 |
| AvB2 | Ava silt loam, 2 to 6 percent slopes, eroded | Ava, eroded | 95 | 115 | 40 | 52 | 3.8 | 7.6 |
| | | Iva | 3 | 165 | 54 | 66 | 5.4 | 10.8 |
| | | Shakamak | 2 | 130 | 46 | 59 | 4.3 | 8.6 |
| Ay | Ayrshire sandy loam | Ayrshire | 97 | 150 | 49 | 60 | 5.0 | 10.0 |
| | | Rensselaer | 3 | 165 | 47 | 66 | 5.4 | 10.8 |
| Bb | Bartle silt loam, 0 to 2 percent slopes | Bartle | 95 | 135 | 46 | 61 | 4.5 | 9.0 |
| | | Peoga | 3 | 150 | 46 | 60 | 5.0 | 10.0 |
| | | Pekin | 2 | 130 | 46 | 59 | 4.3 | 8.6 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|---|------------|------------|----------------|--------------------|--------------------------|-------------|
| BcF | Berks-Ebal complex, 15 to 60 percent slopes | Berks | 65 | --- | --- | --- | 0.0 | 0.0 |
| | | Ebal | 35 | --- | --- | --- | 2.8 | 5.6 |
| BfG | Berks-Rock outcrop complex, 45 to 70 percent slopes | Berks | 75 | 0 | 0 | 0 | 0.0 | 0.0 |
| | | Rock outcrop | 25 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| BlE | Bloomfield sand, 15 to 25 percent slopes | Bloomfield | 100 | --- | --- | --- | 1.8 | 3.6 |
| BlG | Bloomfield sand, 35 to 60 percent slopes | Bloomfield | 100 | --- | --- | --- | 0.0 | 0.0 |
| Bo | Bonnie silt loam, frequently flooded | Bonnie | 100 | 125 | 33 | 0 | 0.0 | 0.0 |
| Br | Booker clay | Booker | 100 | 150 | 42 | 68 | 5.0 | 10.0 |
| Bs | Booker mucky clay | Booker | 100 | 150 | 42 | 68 | 5.0 | 10.0 |
| CcE2 | Chetwynd silt loam, 18 to 25 percent slopes, eroded | Chetwynd | 100 | --- | --- | --- | 3.1 | 6.2 |
| CcF | Chetwynd silt loam, 25 to 60 percent slopes | Chetwynd | 100 | --- | --- | --- | 0.0 | 0.0 |
| Cfc2 | Cincinnati silt loam, Wabash Lowland, 6 to 12 percent slopes, eroded | Cincinnati, eroded | 90 | 105 | 37 | 47 | 3.5 | 7.0 |
| | | Cincinnati, severely eroded | 5 | 100 | 35 | 45 | 3.3 | 6.6 |
| | | Stendal, rarely, very brief | 5 | 150 | 49 | 60 | 5.0 | 10.0 |
| Cfc3 | Cincinnati silt loam, Wabash Lowland, 6 to 12 percent slopes, severely eroded | Cincinnati, severely eroded | 95 | 100 | 35 | 45 | 3.3 | 6.6 |
| | | Stendal, occasionally flooded, very brief | 5 | 140 | 46 | 20 | 4.6 | 9.2 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|---|------------|------------|----------------|--------------------|--------------------------|-------------|
| CfD2 | Cincinnati silt loam, 12 to 18 percent slopes, eroded | Cincinnati | 100 | 110 | 39 | 44 | 3.6 | 7.2 |
| CfD3 | Cincinnati silt loam, 12 to 18 percent slopes, severely eroded | Cincinnati, severely eroded | 85 | 100 | 35 | 45 | 3.3 | 6.6 |
| | | Cincinnati, eroded | 5 | 90 | 32 | 41 | 3.0 | 6.0 |
| | | Hickory, eroded | 5 | --- | --- | --- | 3.3 | 6.6 |
| | | Stendal, occasionally flooded, very brief | 5 | 140 | 46 | 20 | 4.6 | 9.2 |
| ChC2 | Cincinnati silt loam, channery substratum, 6 to 12 percent slopes, eroded | Cincinnati, channery substratum | 100 | 115 | 40 | 52 | 3.8 | 7.6 |
| Cu | Cuba silt loam, frequently flooded | Cuba | 100 | 110 | 39 | 0 | 0.0 | 0.0 |
| EcD | Ebal-Gilpin silt loams, 12 to 18 percent slopes | Ebal | 70 | 100 | 35 | 40 | 3.3 | 6.6 |
| | | Gilpin | 30 | 100 | 35 | 40 | 3.3 | 6.6 |
| EfD2 | Ebal-Wellston silt loams, 10 to 18 percent slopes, eroded | Ebal | 60 | 90 | 32 | 36 | 3.0 | 6.0 |
| | | Wellston | 40 | 105 | 37 | 42 | 3.5 | 7.0 |
| EnA | Elston loam, 0 to 2 percent slopes | Elston | 97 | 130 | 46 | 65 | 4.3 | 8.6 |
| | | Rensselaer | 3 | 165 | 47 | 66 | 5.4 | 10.8 |
| Ev | Evansville silt loam, rarely flooded | Evansville | 100 | 160 | 46 | 64 | 5.3 | 10.6 |
| FaB | Fairpoint silt loam, reclaimed, 2 to 6 percent slopes | Fairpoint | 100 | 145 | 51 | 58 | 4.8 | 9.6 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| FcC | Fairpoint parachannery clay loam, 2 to 12 percent slopes | Fairpoint | 100 | 100 | 35 | 40 | 3.3 | 6.6 |
| FcE | Fairpoint parachannery clay loam, 18 to 35 percent slopes | Fairpoint | 100 | --- | --- | --- | 0.0 | 0.0 |
| FcG | Fairpoint very parachannery loam, 35 to 90 percent slopes | Fairpoint | 100 | --- | --- | --- | 0.0 | 0.0 |
| GcE2 | Gilpin silt loam, 18 to 25 percent slopes, eroded | Gilpin | 100 | --- | --- | --- | 2.0 | 4.0 |
| GfF | Gilpin-Berks complex 30 to 60 percent slopes | Gilpin | 60 | --- | --- | --- | 0.0 | 0.0 |
| | | Berks | 40 | --- | --- | --- | 0.0 | 0.0 |
| GgE | Gilpin-Ebal silt loams, 18 to 30 percent slopes | Gilpin | 60 | --- | --- | --- | 0.0 | 0.0 |
| | | Ebal | 40 | --- | --- | --- | 2.8 | 5.6 |
| GmE | Gilpin-Wellston silt loams, 18 to 25 percent slopes | Gilpin | 60 | --- | --- | --- | 2.8 | 5.6 |
| | | Wellston | 40 | --- | --- | --- | 3.1 | 6.2 |
| HaE2 | Hagerstown silt loam, 18 to 25 percent slopes, eroded | Hagerstown | 100 | --- | --- | --- | 2.6 | 5.2 |
| Hb | Haymond silt loam, frequently flooded | Haymond | 100 | 120 | 42 | 0 | 0.0 | 0.0 |
| Hc | Haymond silt loam, rarely flooded | Haymond | 100 | 145 | 51 | 58 | 4.8 | 9.6 |
| HdA | Henshaw silt loam, 1 to 3 percent slopes | Henshaw | 97 | 155 | 51 | 62 | 5.1 | 10.2 |
| | | Patton | 3 | 150 | 42 | 60 | 5.0 | 10.0 |
| HeD2 | Hickory silt loam, 12 to 18 percent slopes, eroded | Hickory | 100 | 110 | 39 | 44 | 3.6 | 7.2 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|-----------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| HeE | Hickory silt loam, 18 to 25 percent slopes | Hickory | 100 | --- | --- | --- | 3.3 | 6.6 |
| HeG | Hickory loam, 30 to 60 percent slopes | Hickory | 100 | --- | --- | --- | 0.0 | 0.0 |
| MbB2 | Markland silty clay loam, 2 to 6 percent slopes, eroded | Markland | 100 | 125 | 44 | 50 | 4.1 | 8.2 |
| MgA | McGary silt loam, 0 to 2 percent slopes | McGary | 90 | 140 | 46 | 56 | 4.6 | 9.2 |
| | | Shircliff | 7 | 120 | 42 | 48 | 4.0 | 8.0 |
| | | Zipp | 3 | 140 | 39 | 56 | 4.6 | 9.2 |
| Mo | Montgomery silty clay loam | Montgomery | 82 | 160 | 46 | 64 | 5.3 | 10.6 |
| | | Montgomery, undrained | 5 | 0 | 0 | 0 | 0.0 | 0.0 |
| | | Montgomery, overwash | 5 | 150 | 42 | 60 | 5.0 | 10.0 |
| | | Zipp | 5 | 140 | 39 | 56 | 4.6 | 9.2 |
| | | McGary | 3 | 130 | 42 | 52 | 4.3 | 8.6 |
| Mu | Muskego muck | Muskego, drained | 100 | 135 | 33 | 54 | 4.5 | 9.0 |
| Ne | Newark loam, frequently flooded | Newark | 94 | 130 | 42 | 0 | 0.0 | 0.0 |
| | | Petrolia | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Wilhite | 3 | 100 | 25 | 0 | 0.0 | 0.0 |
| No | Nolin silt loam, occasionally flooded | Nolin | 94 | 125 | 44 | 18 | 4.1 | 8.2 |
| | | Newark | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Petrolia | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|---------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| Nr | Nolin silt loam, rarely flooded | Nolin | 94 | 145 | 51 | 58 | 4.8 | 9.6 |
| | | Newark | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Petrolia | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| Pbc2 | Parke silt loam, 6 to 12 percent slopes, eroded | Parke | 100 | 130 | 46 | 52 | 4.3 | 8.6 |
| Pbd2 | Parke silt loam, 12 to 18 percent slopes, eroded | Parke | 100 | 115 | 40 | 46 | 3.8 | 7.6 |
| Pc | Patton silty clay loam, 0 to 1 percent slopes | Patton, drained | 90 | 160 | 46 | 64 | 5.3 | 10.6 |
| | | Henshaw, drained | 5 | 155 | 51 | 62 | 5.1 | 10.2 |
| | | Montgomery, drained | 5 | 160 | 46 | 64 | 5.3 | 10.6 |
| PdB2 | Pekin silt loam, 2 to 6 percent slopes, eroded | Pekin | 100 | 125 | 44 | 56 | 4.1 | 8.2 |
| Pf | Peoga silt loam | Peoga | 100 | 155 | 47 | 62 | 5.1 | 10.2 |
| Pg | Piankeshaw silt loam, frequently flooded | Piankeshaw | 100 | 130 | 46 | 18 | 4.3 | 8.6 |
| PkB2 | Pike silt loam, 2 to 6 percent slopes, eroded | Pike | 100 | 130 | 46 | 52 | 4.3 | 8.6 |
| PkC2 | Pike silt loam, 6 to 12 percent slopes, eroded | Pike | 100 | 120 | 42 | 48 | 4.0 | 8.0 |
| PrB | Princeton fine sandy loam, 2 to 6 percent slopes | Princeton | 100 | 135 | 47 | 54 | 4.5 | 9.0 |
| PrC | Princeton fine sandy loam, 6 to 12 percent slopes | Princeton | 100 | 125 | 44 | 50 | 4.1 | 8.2 |
| RaA | Reesville silt loam, 0 to 2 percent slopes | Reesville | 100 | 170 | 56 | 68 | 5.6 | 11.2 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|---|---------------------------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| Rb | Rensselaer sandy loam | Rensselaer | 100 | 170 | 49 | 68 | 5.6 | 11.2 |
| Rd | Rensselaer loam | Rensselaer | 100 | 175 | 51 | 70 | 5.8 | 11.6 |
| RmA | Roby sandy loam, 0 to 2 percent slopes | Roby | 97 | 125 | 37 | 50 | 4.1 | 8.2 |
| | | Rensselaer | 3 | 165 | 47 | 66 | 5.4 | 10.8 |
| ScA | Shakamak silt loam, 1 to 3 percent slopes | Shakamak | 100 | 130 | 46 | 59 | 4.3 | 8.6 |
| So | Steff silt loam, rarely flooded | Steff | 100 | 145 | 51 | 58 | 4.8 | 9.6 |
| Sr | Steff silt loam, frequently flooded | Steff | 100 | 120 | 42 | 0 | 0.0 | 0.0 |
| St | Stendal silt loam, frequently flooded | Stendal | 97 | 125 | 37 | 0 | 0.0 | 0.0 |
| | | Bonnie | 3 | 120 | 32 | 0 | 0.0 | 0.0 |
| Ud | Udorthents, loamy | Udorthents | 100 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| UnE | Uniontown silt loam, 18 to 30 percent slopes | Uniontown | 100 | --- | --- | --- | 0.0 | 0.0 |
| VgA | Vigo silt loam, 0 to 2 percent slopes | Vigo | 97 | 155 | 51 | 62 | 5.1 | 10.2 |
| | | Very deep, poorly drained, silty soil | 3 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| W | Water | Water | 100 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| WcA | Waupecan silt loam, 0 to 2 percent slopes, rarely flooded | Waupecan | 100 | 160 | 56 | 64 | 5.3 | 10.6 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|--|-------------------------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| WeD2 | Wellston silt loam, 12 to 18 percent slopes, eroded | Wellston | 85 | 105 | 37 | 42 | 3.5 | 7.0 |
| | | Zanesville, soft bedrock substratum | 6 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Tulip | 5 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| | | Adyeville | 4 | Not Rated | Not Rated | Not Rated | Not Rated | Not Rated |
| WeD3 | Wellston silt loam, 12 to 18 percent slopes, severely eroded | Wellston, severely eroded | 90 | 105 | 37 | 42 | 3.5 | 7.0 |
| | | Ebal, severely eroded | 5 | 75 | 26 | 34 | 2.5 | 5.0 |
| | | Gilpin, severely eroded | 5 | 70 | 25 | 28 | 2.3 | 4.6 |
| WgD2 | Wellston silt loam, karst, 6 to 18 percent slopes, eroded | Wellston | 100 | 115 | 40 | 46 | 3.8 | 7.6 |
| Wm | Wilhite silty clay, frequently flooded | Wilhite | 100 | 110 | 28 | 0 | 0.0 | 0.0 |
| Wt | Wirt very fine sandy loam, frequently flooded | Wirt | 100 | 105 | 37 | 0 | 0.0 | 0.0 |
| ZaA | Zanesville silt loam, 1 to 3 percent slopes | Zanesville | 100 | 140 | 49 | 63 | 4.6 | 9.2 |
| ZaB2 | Apalona-Zanesville silt loams, 2 to 6 percent slopes, eroded | Apalona, eroded | 45 | 135 | 47 | 61 | 4.5 | 9.0 |
| | | Zanesville, eroded | 40 | 130 | 46 | 59 | 4.3 | 8.6 |
| | | Wellston, eroded | 10 | 140 | 49 | 56 | 4.6 | 9.2 |
| | | Johnsburg | 5 | 150 | 51 | 68 | 5.0 | 10.0 |

COMPONENT CROP YIELDS (RV)--Continued
 Greene County, Indiana

| Map Unit Symbol | Map unit name | Component | Comp. Pct. | Corn Bu/Ac | Soybeans Bu/Ac | Winter Wheat Bu/Ac | Grass-Legume Hay Tons/AC | Pasture AUM |
|-----------------|--|-----------------------------|------------|------------|----------------|--------------------|--------------------------|-------------|
| ZaC2 | Apalona-Zanesville silt loams, 6 to 12 percent slopes, eroded | Apalona, eroded | 45 | 125 | 44 | 56 | 4.1 | 8.2 |
| | | Zanesville, eroded | 40 | 110 | 39 | 50 | 3.6 | 7.2 |
| | | Deuchars, eroded | 10 | 140 | 49 | 56 | 4.6 | 9.2 |
| | | Wellston, eroded | 5 | 130 | 46 | 52 | 4.3 | 8.6 |
| ZaC3 | Apalona-Zanesville silt loams, 6 to 12 percent slopes, severely eroded | Apalona, severely eroded | 45 | 115 | 40 | 52 | 3.8 | 7.6 |
| | | Zanesville, severely eroded | 40 | 110 | 39 | 50 | 3.6 | 7.2 |
| | | Deuchars, severely eroded | 10 | 125 | 44 | 50 | 4.1 | 8.2 |
| | | Wellston, severely eroded | 5 | 120 | 42 | 48 | 4.0 | 8.0 |
| Zp | Zipp silty clay | Zipp | 100 | 135 | 37 | 61 | 4.5 | 9.0 |